

# TURF PRIDE® 92 Weed & Feed

For St. Augustine & Centipede Lawns

WEED CONTROL WITH PREMIUM LAWN FOOD

50 POUND BAG COVERS 10,000 Sq.Ft. of Lawn

Controls weeds; Florida Pusley, Lippia, Pigweed and certain broadleaf Weeds as listed. Prevents Crabgrass, Paspalums,

Spotted Spurge and other weeds as listed.

ONLY FOR USE ON ST. AUGUSTINE AND CENTIPEDE LAWNS

## ACTIVE INGREDIENT:

Atrazine: CAS [1912-24-9] - - - - 0.92%


[2 Chloro-4-ethylamino-6- (propylamino) s triazine]

OTHER INGREDIENTS: - - - - 99.08%

Total 100.00%

Contains 18.4 lbs. of Atrazine per ton.

when used according to label directions applies 2 lbs. a.i. per acre.



## KEEP OUT OF REACH OF CHILDREN CAUTION

### FIRST AID

IF SWALLOWED - Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES - Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN - Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment.

SEE ADDITIONAL PRECAUTIONARY STATEMENTS ON THIS LABEL

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid inhalation or contact with eyes. Wash after handling or using. Keep away from feed or food products.

### ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on turf being grown for sale, or other commercial uses, such as for commercial seed production, or for research purposes. For use on St. Augustine and Centipede turfgrass being grown in parks, on golf courses and lawns.

### READ ENTIRE LABEL BEFORE USING

Read the entire Directions for Use, CONDITIONS OF SALE, Disclaimer of Warranties, and Limitation of Liability, before using this product.

## WHERE TO USE

This material is for use only on St. Augustine and Centipede lawns. On newly seeded centipede, delay application until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within 6 months after application. Do not plant any crop (flower or vegetable plants) to treated areas for 18 months or injury may result. Apply this product only when grass foliage is dry.

DO NOT USE ON BAHIA, BENT, BERMUDA, BLUEGRASS, DICHONDRA LAWNS, FESCUE, OR RYE, GRASSES, BECAUSE SERIOUS TURF DAMAGE CAN OCCUR.

## WHEN TO APPLY

Best weed control will result if applied before weeds sprout since the product acts as a Pre-Emergent and also as a Post Emergent weed killer. Established weeds are harder to kill and may require an additional application in the fall or spring. Apply January 1st up to March 1st for best control otherwise product may be applied up to June 15th with good weed control. A fall application will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit Tobacco, Florida Betony and Evening Primrose.

In case of heavy weed infestation or low soil moisture, a repeated application may be necessary. Time for complete kill of mature weeds may be 4 to 5 weeks. Wait 2 months for repeat application. Apply when air is calm to prevent particle wind drift.

## HOW TO APPLY

This product may be applied with either a drop type or broadcast type spreader. Apply evenly across the entire lawn at a rate of 50 pounds to treat 10,000 Sq.Ft. — 40 lbs. to treat 8,000 sq.ft. of lawn. (equivalent to 5 lbs. per 1,000 Sq.Ft.) Do not exceed recommended rates. Sprinkle the area immediately after application with water to wash granules off the grass blades and down to soil level.

## USE PRECAUTIONS

Do not apply this product on or over the root zone of ornamental trees, shrubs, bedding plants, flowers, or garden plants because injury to them may result. Do not apply to lawns containing grasses or mixtures of grasses other than St. Augustine or Centipede grasses as recommended on this label, because serious injury to those other grasses may result.

### SPREADER SETTINGS for 5 lbs./ 1,000 sq.ft.

CYCLONE 4.5 SCOTT 7

**NOTE:** Spreader rate varies with walking speed. Check spreader over a known square foot area.

(35-46-acceptd-8-3-01-p1of2-9-13-01)

**HOWARD FERTILIZER and CHEMICAL COMPANY**  
P.O. BOX 628202 - ORLANDO, FL 32862

2/2

**TURF PRIDE® 92 Weed & Feed****For St. Augustine & Centipede Lawns****EPA REG. NO. 35512-46****EPA EST. NO. 35512-FL-1****NET WEIGHT 50 POUNDS****WEEDS CONTROLLED  
OR SUPPRESSED**

Annual Bluegrass (Poa Annua)	Henbit
Annual Sowthistle	Knotweed
Clover	Lambsquarter
Black Medic	Lespedeza
Blue Toadflax	Lippia (Matchweed)
Bull Paspalum	Moneywort
Buckhorn Plantain	Morning Glory
Bur Clover	Narrowleaf Vetch
Carolina Geranium (Cranesbill)	Parsley-Piert
Chickweed (Common and Mouse Ear)	Oxalis (Common and Yellow Woodsorrel)
Cocklebur	Pigweed
Common Dandelion	Purslane
Common Purslane	Ragweed
Crabgrass	Red Sorrel
Crowfoot Grass	Rescuegrass
Cudweed	Sandspur
Cutleaf Evening Primrose	Sedges*
Dichondra	Shepards Purse
Dollar Weed	Sicklepod
Florida Beggarweed	Spotted Spurge
Florida Betony	Spurweed
(Rattlesnake Weed)	Stinging Nettle
Florida Pusley	Swinecrest
Fragrant Cudweed	Venus Looking Glass
(Rabbit Tobacco)	Virginia Creeper
	Wild Mustard
	Wood Sorrel

*\*Repeat application are required for control of Nutsedge grasses.***GUARANTEED FERTILIZER ANALYSIS**

Total Nitrogen (N) - - - - -	variable%
Nitrate Nitrogen	variable%
Ammoniacal Nitrogen	variable%
Water Soluble Nitrogen	variable%
(and/or Urea Nitrogen)	
Water Insoluble Nitrogen	variable%
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) - - - - -	variable%
Soluble Potash (K <sub>2</sub> O) - - - - -	variable%
Chlorine not more than	variable%

Derived from Sulfate of Ammonia, Ammonium Phosphate,  
Muriate of Potash, Sulphate-Potash-Magnesium. Potential Acidity Equivalent  
to 1245 lbs. of Calcium Carbonate per ton.

Statement of Secondary Plant Foods

Total Magnesium as Mg - 1.20%; Iron as Fe - 1.00%; Water Soluble Magnesium Mg - 1.20%; Zinc as Zn - 0.05% Manganese as Mn - 0.50%; Boron as B - 0.02% Copper as Cu - 0.02%; Molybdenum as Mo - 0.002% Derived from: Manganese Sulphate, Iron Oxide, Manganese Oxide, Copper Oxide, Zinc Oxide, Borax, Sodium Molybdate.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store product in its original labeled container in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment, then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. **HOUSEHOLD:** Do not reuse empty bag. Wrap in several layers of newspaper and discard in trash.

**CONDITIONS OF SALE**

If the terms are not acceptable, return the unopened container at once. By using this product, user and buyer accept the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and should be followed carefully. However it is impossible eliminate all risks associated with the use of this product. Injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Howard Fertilizer and Chemical Co. All such risks shall be assumed by the user or buyer.

**Disclaimer of Warranties:** There are no warranties express or implied, of merchantability or of fitness for a particular purpose or otherwise, which extend beyond the statements made on this label. No agent of Howard Fertilizer and Chemical Co. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Howard Fertilizer and Chemical Co. disclaims any liability whatsoever for special, incidental, or consequential damages resulting from the use or handling of this product.

**LIMITATION OF WARRANTIES:** The exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price, or at Howard Fertilizer and Chemical Co.'s option, the replacement of the product.

**HOWARD**  
**FERTILIZER and CHEMICAL**  
**COMPANY**

**P.O. BOX 628202 - ORLANDO, FL 32862**