7-30-2003

1/2

Please read instructions	en r	everse before comp	eting form.			Form Ap	proved	. OMB No.	. 207 0-	0060	Approve	expires 2-28-9	
United States Environmental Protect Washington, DC 2					tion Agency			Registration Amendment Other			OPP Identifier Number		
			Applica	tior	for Pestici	de - Sec	tion	l					
1. Company/Product Number 35512-48					2. EPA Product Manager Tompkins					3. Proposed Classification			
4. Company/Product (Name) Turf Pride 115 Weed & Feed					РМ# 25					ت			
5. Name and Address of Applicant (Include ZIP Code) Howard Fertilizer & Chemical Co., Inc. P.O. Box 628202 Orlando, FL 32862-8202 Check if this is a new address					6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. Product Name								
					Section - l	<u> </u>							
Amendment - Explain below. Resubmission in response to Agency letter date Notification - Explain below.				Agency let				pplication.				CATION 0 2003	
NOTIFICATION OF ADDI 18 LB BAG COVERS 4,51 Apply evenly across the e This notification is consistent with the Fertilizer & Chemical Company Inc. 98-10 and 40 CFR 152-46, this prod	00 S intire e prov under uct mi	SQ. FT. 8 lawn at a rate of 18 visions of PR Notice 98-10 and stands that it is a violation of 16 say be in violation of FIFRA and	pounds to treater EPA regulations at 4 1 U.S.C. Sec. 1001 I	at 4,5 10 CFR 10 willfu	500 sq. ft. 3 152.46, and no other cha- illy make any false stateme	nges have been mi nt to EPA, and furt subject to enforcer	ade to the	labeling or the Co	onfidential (notification	Statemer is not co	insistent with the to		
1. Material This Product 1	Will	Be Packaged In:			T			T	<u> </u>				
Child-Resistant Packaging Yes No	Jnit Packaging Yes No No No. per			Water Soluble Packaging Yes No If "Yes" No. per			2. Type of Contains Metal Plastic Glass Paper						
* Certification must be submitted		Unit Packaging wg			Package wgt contain					Other (Specify)			
3. Location of Net Conten	4. Size(s) F	4. Size(a) Retail Container 5. Loc 18LB, 40LB, 50LB					cation of Label Directions On Label						
6. Manner in Which Label is Affixed to Product				Lithograph Other Stanciled									
					Section - I	/							
1. Contact Point (Comple	te i	tems directly below	for identifice	tion	of individu a l to be	contected,	if nece	asary, to p	100 033	this e	pplication.)		
Name Rebecca M. Horton									none No. (Include Area Code) 75-8826				
Certific I certify that the statements I have made on this form an I acknowledge that any knowlingly false or misleading stooth under applicable law.				nd all attachments thereto are true, accurate and complete.					- 1	6. Date Application Received (Stamped)			
2. Signature Ribicou M. Hotton				3. Title Consultant/Agent							新 (水域の) 1 () () () 2 () () () () () () () () () (
4. Typed Name Rebecca M. Horton				5.	Date ク-15-	03						or E. G. C. C. G. C.	

For St. Augustine & Centipede Lawns Weed Control with Premium Lawn Food 50 POUND BAG TREATS 12,500 Sq. Ft.

(18 POUND BAG TREATS 4,500 Sq. Ft.)

Controls weeds; Florida Pusley, Lippiea, Pigweed and certain Broadleaf Weeds as listed. Prevents Crabgrass, Paspalums, and Spotted Spurge and other weeds as listed.

ONLY FOR USE ON ST. AUGUSTINE AND CENTIPEDE LAWNS

ACTIVE INGREDIENT:

Atrazine: CAS [1912-24-9]..... OTHER INGREDIENTS:

98.85% TOTAL 100.00%

1.15%

Contains 23 lbs. of a.i. Atrazine per ton and when used according to label directions applies 2 lbs. a.i. per acre.

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

IF SWALLOWED - Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person

IF IN EYES - Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN - Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Cal a poison control center or doctor for treatment advice.

IF INHALED - Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment

READ ENTIRE LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid inhalation or contact with eyes. Wash after handling or using. Keep away from feed or food products

ENVIRONMENTAL HAZARDS

Arrazine can travel (seep or leach) through soil and can enter ground water, which may be used as drinking water Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well drained Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water

This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertibal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

READ ENTIRE LABEL BEFORE USING

Read the entire Directions for Use, Conditions of Sale, Disclaimer of Warranties and Limitation of Liability before using this product.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on turf being grown for sale, or other commercial uses, for commercial seed production, or for research purposes. For use on St Augustine and Centipede turfgrass being grown in parks, on golf courses and lawns

WHERE TO USE

<u>This material is for use only on St. Augustine and Centipede (awns.</u> On newly seeded centipede, delay application until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within 6 months after application. Do not plant any crop (flower or vegetable plants) to treated areas for 18 months or injury may result. Apply this product only when grass foliage is dry.

DO NOT USE ON BAHIA, BENT, BERMUDA, BLUEGRASS, DICHONDRA LAWNS, FESCUE, OR RYE,

GRASSES, BECAUSE SERIOUS TURF DAMAGE CAN OCCUR.

WHEN TO APPLY

Best weed control will result if applied before weeds sprout since the product acts as a Pre-Emergent and also as a Post Emergent weed killer. Established weeds are harder to kill and may require an additional application in the fall or spring. Apply January 1* up to March 1* for best control otherwise product may be applied up to June 15* with good weed control. A fall application will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit Tobacco, Florida Belony and Evening Primrose.

In case of heavy weed intestation or low soil moisture, a repeated application may be necessary. Time for complete kill of mature weeds may be 4 to 5 weeks. Wait 2 months for repeat application. Apply when air is calm to prevent particle wind drift

HOW TO APPLY

This product may be applied with either a drop type or broadcast type spreader. Apply evenly across the entire lawn at a rate of 50 pounds to treat 12.500 Sq. Ft. — 40 lbs. to treat 10.000 sq. ft. — 18 lbs. to treat 4.500 sq. ft. of lawn, (equivalent to 4 lbs. per 1,000 Sq. Ft.) Do not exceed recommended rates. Sprinkle the area immediately after

WEEDS CONTROLLED OR SUPPRESSED

Annual himeorass (Poa Annua) **Annual Sowthistle** Black Medic Blue Toadflax **Bull Pasnalum Buckhom Plantain Bur Clover** Carolina Geranium (Cranesbill) Chickweed (Common and Mouse Earl Cocklebur Common Dandelion Common Pursiane Crabgrass Crowfoot Grass Cudweed Cuteat Evening Primrose Dichondra **Dollar Weed** Florida Beggerweed Florida Retony (Rattlesnake weed) Florida Pursley Fragrant Cudweed

(Rabbit Tobacco)

Henbit Knotwood NOTIFICATION Lambaquarte Lespedeza JUL 3 0 2003 Lippis (Matchweed) Moneywort **Morning Glory** Narrowleaf Vetch Parsley-Piert Oxalis (Common and Yellow Woodsorrel) Pigweed Purslane Ragweed Red Sorrel Rescuegrass Sandspur Sedges' Shephards Purse Sicklepod Spotted Spurge Spurweed Stinging Nettle Swinecrest Venus Lookino Glass Virginia Creeper Wild Mustard Wood Sorrel

*Repeat application are required for control of Nutsedge grasses.

SPREADER SETTINGS for 4lbs. / 1,000 sq. ft.

CYCLONE B-1 Rotary.....# 4 SCOTTS Model 35.....# 6

NOTE: Spreader rate varies with walking speed. Check spreader over a known square food area.

STORAGE AND DISPOSAL

Do not contaminate water, lood or feed by storage or disposal.

STORAGE: Store product in its original tabeled container in a cool, dry, locked place out of reach of

(Commercial Use)

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. <u>CONTAINER DISPOSAL</u>: Completely empty bag into application equipment, then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. [Residential Use]

DISPOSAL: If empty - Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled - Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	variable %
Nitrate Nitrogen	variable %
Ammoniacal Nitrogen	
Water Soluble Nitrogen	variable %
(And/Or Urea Nitrogen)	
Water Insoluble Nitrogen	varisble %
Available Phosphate Acid (P205)	variable %
Soluble Potash (K20)	
Chlorine, not more than	

Derived from: Sulfate of Ammonia, Ammonium Phosphate, Muriate of Potash, Sulphate-Potash-Maganesium. Potential Acidity Equivalent to 1245 lbs. of Calcium Carbonate per ton.

 $Statement of Secondary Plant Foods: \\ Total Magnesium as Mg = 1.20\%; Iron as Fe = 1.00\%; Water Soluble Magnesium Mg = 1.20\%; Zinc as Zn = 1.00\%; Water Soluble Mg = 1.$ 0.05% Manganese as Mo - 0.50%; Boron as B 0.02% Copper as Cu - 0.02%; Nolybdenum as Mo -0.002% Derived from: Manganese Sulphate, Iron Oxide, Manganese Oxide, Copper Oxide, Zinc Oxide, Borax, Sodium Molybdate

CONDITIONS: The directions for use of this product are believed to be adacustosand should be followed. carefully. However it is impossible eliminate all risks associated with this use of this product. Idury, ineffectiveness or other unintended consequences may result because of grith factors is visabler conditions. presence of other materials, or the manner of use or application, all of which a wovered the control of Howard Fertilizer & Chemical Co., Inc. All such risks shall be assumed by the user or buyer.

6 3

....

00006

0 0