



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
35512-56

Date of Issuance:
MAR 14 2005

Term of Issuance:
Conditional

Name of Pesticide Product:
Turf Pride® Lawn Food
with Trimec® Herbicide

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Howard Fertilizer & Chemical Co., Inc.
P.O. Box 628202
Orlando, FL 32862-8202

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

- 1. Add the phrase "EPA Registration No. 35512-56" to the label before you release the product for shipment.

COMMENTS CONTINUED ON PAGE 2 OF THIS NOTICE OF REGISTRATION

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product under the enclosed stamped copy of the label constitutes acceptance of these conditions.

Enclosure

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Signature of Approving Official:

Joanne I. Miller

Date:

MAR 14 2005

Page 2
EPA Reg. No. 35512-56
Comments Continued:

2. Submit the following data required for the registration of this pesticide product within 1 year from the date of this Notice of Registration:

<u>EPA Guideline Data Number</u>	<u>Guideline Descriptor</u>
830.6317	Storage Stability Study
830.6320	Corrosion Characteristics Study
870.2400	Acute Eye Irritation Study

The eye irritation study must be conducted with the highest free level of nitrogen, as described in PR Notice 2001-2.

3. Revise the "Container Disposal" statement to read:
If empty: Do not reuse this container. Place in trash or offer for recycling if available.
If partially filled: Call your local solid waste agency or toll free number (1-800-_____ for disposal instructions. Never place unused product down any indoor or outdoor drain.
4. Revise the "Storage" statements to read: "Store in original container and keep bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs".
5. Add the contact phone number to the "First Aid" statements.
6. Submit one (1) copy of the final printed labeling before you release this product for shipment.
7. Submit and/or cite all data required for the registration of this product when the Agency requires all registrants of similar products to submit data; and submit acceptable responses required for re-registration of this product under FIFRA, section 4. A stamped copy of the label is enclosed for your records.

Enclosure

3 8 4
ACCEPTED
with COMMENTS
In EPA Letter Dated:
MAR 14 2005

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

35512-56

Turf Pride[®] Lawn Food with Trimec[®] Herbicide

A TRIPLE ACTION BROADLEAF HERBICIDE WITH LAWN FOOD
 50 LB TREATS UP TO 5000 SQ. FT. OF TURFGRASS

FOR USE IN CONTROL OF BROADLEAF WEEDS IN HOME LAWNS, APARTMENT AND CONDOMINIUM TURF, IN PARKS, PLAYGROUNDS AND OTHER ORNAMENTAL AND RECREATIONAL TURFGRASS

A BROADLEAF WEED KILLER FOR SOUTHERN LAWNS

BALANCED LAWN FOOD GIVES TURF FAST-GREENING WITH EXTENDED FEEDING

CONTROLS: Dandelions, Chickweed, Knotweed and many Other Weeds as listed.

For use on St. Augustinegrass, Centipedegrass, Zoysiagrass, Bahiagrass, Bermudagrass, Bluegrass, Bentgrass and Fescue varieties.

ACTIVE INGREDIENTS:

- Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid0.20%
- Dimethylamine salt of 2,4-dichlorophenoxyacetic acid.....0.11%
- Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid).....0.05%

OTHER INGREDIENTS:99.64%
TOTAL:100.00%

This product contains 3.2 LB or 0.16% (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid per ton, 1.80 LB or 0.09% 2,4-dichlorophenoxyacetic acid per ton, 0.80 LB or 0.04% dicamba (3,6-dichloro-o-anisic acid) per ton.

Isomer specific by AOAC Methods. Contains the single isomer form of Mecoprop-p. TRIMEC[®] is a registered trademark of PBI/GORDON Corporation.

GUARANTEED ANALYSIS

- | | |
|---------------------------------|-------|
| Total Nitrogen (N)..... | X-X-X |
| x.xx% Ammoniacal Nitrogen | x.xx% |
| x.xx% Nitrate Nitrogen | |
| x.xx% Urea Nitrogen | |
| x.xx% Water Insoluble Nitrogen* | |

Available Phosphate (P₂O₅)..... x.xx%
 Soluble Potash (K₂O)..... x.xx%

Chlorine (Cl), not more than x.xx%

Derived from: _____

*x.xx% slowly available nitrogen from _____

F387

Information concerning the raw materials composing this product can be obtained by writing to the Guarantor listed above and referring to the batch number found on this container

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact x-xxx-xxx-xxxx for emergency medical treatment information.

If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If swallowed	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

ENVIRONMENTAL HAZARDS: This product is extremely toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. Do not apply when weather conditions favor drift from treated areas.

READ ENTIRE LABEL BEFORE USING

Read the entire Directions for Use, Conditions, Disclaimer of Warranties, and Limitation of Liability before using this product. If the terms are not acceptable, return the unopened container at once. By using this product, user and buyer accept the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However it is impossible to eliminate all risks associated with the use of this product. Injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Howard Fertilizer & Chemical Co., Inc. All such risks shall be assumed by the user or buyer. **DISCLAIMER OF WARRANTIES:** There are no warranties express or implied, of merchantability or of fitness for a particular purpose or otherwise, which extend beyond the statements made on this label. No agent of Howard Fertilizer & Chemical Co. is authorized to make any warranties contained herein. Howard Fertilizer & Chemical Co. disclaims any liability whatsoever for special, incidental, or consequential damages resulting from the use or handling of this product. **LIMITATION OF WARRANTIES:** The exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price, or at Howard Fertilizer & Chemical Co., Inc.'s option, the replacement of the product.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until dust has settled.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Storage & Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store product in its original closed container in a cool, dry place out of reach of children and animals. In case of spill, avoid contact. Isolate spills, sweep up, and use as directed on this label.

Pesticide Disposal: Wastes resulting from a spill or from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into equipment and use as directed. Do not re-use empty bag. Dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Turf Pride[®] is a registered trademark of Howard Fertilizer & Chemical Company, Inc.
 Manufactured & Guaranteed by
HOWARD FERTILIZER & CHEMICAL COMPANY, INC.
 P.O. BOX 628202 - ORLANDO, FL 32862-8202

NET WT 50 LB (22.7 kg)

EPA Reg. No. 35512-_____
 Nov. 22, 2004

EPA Est. No. 35512-FL-2
 Page 1 of 2 Pages

USE PRECAUTIONS: Do not use this product where desirable clovers are present or allow the material to drift onto flowers, vegetables or ornamental shrubbery. Do not spread over exposed roots of trees and ornamentals. Do not exceed specified dosages for any area; be particularly careful within dripline of tree and other ornamental species. Apply when the air is calm. Do not apply to newly seeded lawns.

Do not use on dichondra. Do not over-apply. Under conditions of adequate moisture, high temperature and high humidity, over-application may cause a minor transitory discoloration on St. Augustinegrass.

WHERE TO USE: This product may be used on all varieties of Zoysiagrass, Bermudagrass, Centipedegrass, St. Augustinegrass, Bahiagrass, Bentgrass, Fescues and Bluegrass.

WHEN TO USE: Apply when weeds are young and actively growing in the spring, summer or fall. Best results will be obtained if applied in the morning when dew is on the grass and when daily temperatures range from 65° to 90°F. For lawns that are heavily over-run by hard-to-kill weeds or in a state of poor growth due to low moisture conditions, a second treatment in 30 to 45 days may be necessary.

The maximum application rate to turf is 0.5 lb. 2,4-D acid equivalent per acre per application site. The maximum number of broadcast applications per treatment site is 2 per year.

HOW TO APPLY: Mow lawn to normal height 1 to 2 days before application. Water lawn thoroughly, as needed, at least 1 day before application to sustain moisture until the next watering (1-2 days after application). Choose a calm day to ensure uniform coverage and avoid spreading granules onto non-target plants. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto non-target plants. Wait 1-2 days following application, then water treated area thoroughly.

SPREADER SETTINGS

To check for your spreader settings refer to your spreader manual details and apply 10 pounds of product to cover 1,000 sq. ft. of turfgrass. During application, close spreader openings when turning.

BROADLEAF WEEDS CONTROLLED

Annual yellow sweetclover, Aster, Austrian fieldcress, Bedstraw, Beggertick, Betony (Florida), Bindweed (field), Bird vetch, Bitter wintercress, Bittercress (hairy), Bitterweed, Black-eyed Susan, Black medic, Black mustard, Blackseed plantain, Blessed thistle, Bloodflower milkweed, Blue lettuce, Blue vervain, Bracted plantain, Brassbuttons, Bristly oxtongue, Broadleaf dock, Broadleaf plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull thistle, Bullnettle, Burclover, Burdock, Burning nettle, Burweed, Buttercup, Buttonweed, Canada thistle, Carolina geranium, Carpetweed, Catchweed bedstraw, Catnip, Catsear, Chickweed (common, mouseear), Chicory, Cinquefoil, Clover (crimson, hop, red, strawberry, sweet, white), Cockle, Cocklebur, Common mullein, Creeping jenny, Cudweed, Curly dock, Daisy (English, fleabane, oxeye), Dandelion, Dichondra, Dogbane, Dogfennel, Dollarweed, False dandelion, False flax, False sunflower, Fiddleneck, Florida pusley, Frenchweed, Galinsoga, Goathead, Goldenrod, Ground ivy, Gumweed, Hairy fleabane, Hawkweed, Healall, Heartleaf drymary, Heathbaster, Hedge bindweed, Hedge mustard, Hemp, Henbit, Hoary cress, Hoary plantain, Hoary vervain, Horsenettle, Jimsonweed, Knawel, Knorweed, Kochia, Lambsquarters, Lespedeza, Mallow, Matchweed, Mexicanweed, Milk vetch, Morningglory, Mouseear hawkweed, Mugwort, Musk thistle, Mustard, Narrowleaf plantain, Narrowleaf vetch, Nettle, Orange hawkweed, Oriental cocklebur, Oxalis, Parsley-piert, Parsnip, Pearlwort, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Pineywoods bedstraw, Plains coreopsis (tickseed), Plantain, Poison ivy, Poison oak, Pokeweed, Poorjoe, Prairie sunflower, Prickly lettuce, Prickly sida, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Prostrate vervain, Puncturevine, Purslane (common), Ragweed, Red sorrel, Redroot pigweed, Redstem filaree, Rough cinquefoil, Rough fleabane, Russian pigweed, Russian thistle, Scarlet pimpernel, Scotch thistle, Sheep sorrel, Shepherdspurse, Slender plantain, Smallflower galinsoga, Smooth dock, Smooth pigweed, Sorrel, Sowthistle, Spanishneedles, Speedwell, Spiny amaranth, Spiny cocklebur, Spiny sowthistle, Spotted catsear, Spotted spurge, Spurweed, Stinging nettle, Strawberry (India mock), Tall nettle, Tall vervain, Tansy ragwort, Tansy mustard, Tanweed, Thistle, Trailing crownvetch, Tumble mustard, Tumble pigweed, Velvetleaf, Venice mallow, Virginia buttonweed, Virginia creeper, Virginia pepperweed, Water pennywort, Wavyleaf bullthistle, Western clematis, Western salsify, White mustard, Wild aster, Wild buckwheat, Wild carrot, Wild four-o'clock, Wild garlic, Wild geranium, Wild lettuce, Wild marigold, Wild mustard, Wild onion, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild vetch, Woodsorrel, Woolly croton, Woolly morningglory, Woolly plantain, Wormseed, Yarrow, Yellow rocket, Yellowflower pepperweed, and other broadleaf weeds