

QUINCY FARM CHEMICALS, INC.

TELONE II

SOIL FUMIGANT

NET CONTENTS

Cameraman's Note
Poor Copy

GENERAL INFORMATION

Use TELONE II soil fumigant only as a **preplanting soil treatment** to control plant parasitic nematodes (root-knot, meadow lesion), citrus cyst formers (golden sugar beet, soybean), burrowing ring, spiral, sting pin, stubby root, stylet, dagger), also to control wireworms and garden centipedes (symphylans). Fumigate land to be planted to the crops listed below by applying TELONE II under the conditions and at the rates recommended under DIRECTIONS FOR USE, DOSAGE AND USE RECOMMENDATIONS, and USE PRECAUTIONS. Read the entire label before using TELONE II.

Vegetable Crops:

asparagus	cauliflower	horseradish	parsnips	shallots
beans	celery	kale	peas	spinach
beets	collards	kohlrabi	peppers	squash (summer)
blackeyed peas	corn	leeks	pimentoes	squash (winter)
broccoli	cowpeas	lettuce	potatoes	sweet potatoes
brussels sprouts	cucumbers	melons	pumpkins	swiss chard
cabbage	egg plant	mustard greens	radishes	tomatoes
cantaloupe	endive	okra	rutabaga	turnips
carrots	garlic	onions	salsify	watermelons

Field Crops:

alfalfa	flax	oats	sericeum
barley	grasses	pasture grass	soybeans
birdsfoot trefoil	hops	peanuts	sugar beets
buckwheat	lespedeza	popcorn	sugarcane
clover	millet	rice	tobacco
corn	milo	rye	vetch
cotton	mint	safflower	wheat

Citrus Fruit Tree Planting Sites:

grapefruit	kumquats	lemons	limes	orange	tangerines	tangelos
------------	----------	--------	-------	--------	------------	----------

Deciduous Fruit and Nut Tree Planting Sites:

almonds	dates	olives	plums
apples	figs	peaches	pomegranates
apricots	filberts	pears	prunes
cashews nuts	hazelnuts	pecans	quince
cherries	hickory nuts	persimmons	walnuts
chestnuts	nectarines	pineapple	

Bush and Vine Planting Sites:

blackberries	currants	huckleberries	youngberries
blueberries	dewberries	loganberries	
boysenberries	gooseberries	raspberries	
cranberries	grapes	strawberries	

DOSAGE AND USE RECOMMENDATIONS
to Control Nematodes, Symphylans, and Wireworms

Crops (consult list of individual crops under General Information)	Type of Treatment	Soil Type	Dosage	
			Gallons Per Acre ¹	Fl Oz 1000 Ft Row Per Chisel ²
Shallow Rooted Plants				

Nursery Crops including floriferous and vine and bramble fruits of

WHEN TO TREAT: Apply TELONE II For best results with annual crop (before October 15) is best such as celery, tomatoes, and crop. **Note:** Treat muck soils or (clay) soils only when they are when it is cold and/or wet.

SOIL PREPARATIONS: TELONE II through the soil and the soil's period. The soil should be in moisture at about one-half of (If undecomposed plant debris) TELONE II. Tillage deeper than or plow pan occurs at plow to simultaneously apply the fumigant.

APPLICATION: TELONE II made application equipment that will surface. For overall application part. When the fumigant is injected up to twice the application depth direction or at an angle to the application, use chisel equipment the soil where the crop is to be distribute about 1 1/2 times more chisels are used per row, application fumigant required per acre will of planting, avoid placing the chisels used per row, place the plant in the center of the area.

SEALING: Immediately after chisel application, use a roller and then compact it by float wheel by listing or by bedding fumigating listed rows, seal it.

EXPOSURE PERIOD: After an exposure period will be required.

AERATION AND PREPARATION: the soil to aerate completely. TELONE II is no longer evident time for each 10 gallons of TELONE II and shrub planting sites, a 3 to 4 inch if heavy rains or low temperature treatment zone. After row treatment possible recontamination of the to the depth of the treatment of muck soils.

quackgrass and for suppression of potatoes. Apply as an overall treatment.

Time of Treatment
Spring
Fall

Mint in Northwestern States: overall treatment in the spring.

Cameraman's Note

Poor Copy

- Citrus Fruit Tree Planting Sites:**
 grapefruit kumquats lemons limes oranges tangerines tangelos
- Deciduous Fruit and Nut Tree Planting Sites:**
 almonds dates olives plums
 apples figs peaches pomegranates
 apricots filberts pears prunes
 cashews nuts hazelnuts pecans quince
 cherries hickory, nuts persimmons walnuts
 chestnuts nectarines pineapple
- Bush and Vine Planting Sites:**
 blackberries currants huckleberries raspberries
 blueberries dewberries loganberries strawberries
 boysenberries gooseberries
 cranberries grapes

**DOSAGE AND USE RECOMMENDATIONS
 to Control Nematodes, Symphylans, and Wireworms**

Crops (consult list of individual crops under General Information)	Type of Treatment	Soil Type	Dosage		
			Gallons Per Acre ¹	Fl Oz. 1000 Ft Row Per Chisel ²	
Shallow Rooted Plants Field Crops Floral Crops Grasses and Turf Small Fruits Vegetables Ornamentals	Row (42") ³	Mineral	4.5 to 6	46 to 52	
		Muck or Peat	9 to 12	93 to 106	
	Overall (Broadcast)	Mineral	9 to 15	26 to 44	
		Muck or Peat	24 ³ to 36	71 to 106	
Strawberries	Overall	Mineral	24 to 36	71 to 106	
Sugar Beets Root-Knot Nematode Sugar Beet Nematode	Row (42")	Mineral	6	62	
	Overall		12 to 15	35 to 44	
	Row (42")		9	93	
	Overall		12 to 18	35 to 51	
Pineapple ⁴	Row	Mineral	24 to 36		
Citrus -- Florida ⁵	Overall	Mineral	36	106	
Nursery and Field Citrus Fruit Trees Deciduous Fruit Trees ⁶ Forest Trees Grapes Nut Trees Ornamentals (deep rooted)	Overall Gallons Per Acre to Penetrate Various Depths				
	Mineral Soils	3 ft	4 ft	5 ft	7 ft
	Sand	15	21	27	33
	Sandy Loam	27	30	36	48
	Silt Loam	42	51	63	75
	Clay Loam	54	69	84	102

- ¹Use the higher rates in heavier soil.
- ²For cyst forming nematodes increase dosage to 18 gallons per acre (53 fl oz / 1000 ft row per chisel).
- ³For muck soils containing less than 30% organic matter use 18 gallons per acre.
- ⁴For Hawaiian pineapple, application may be made at time of or just before planting.
- ⁵For burrowing nematode in citrus inject in 18 inch centers, 12 inches deep, keep free of citrus roots for 2 years before replanting to citrus.
- ⁶Row treatment is not recommended for potatoes in irrigated areas of western and northwestern states.
- Tree planting sites prepared by backhoeing may be treated by plowing 2 1/2" of TELONE II into the soil and planting. For best results prepare and treat sites in the fall and plant in the spring.
- Use TELONE II as an aid in the control of bacterial canker and decline of peach trees by application as a preplant or seed treatment in light (sandy) soils at the rate of 24 to 36 gallons per acre (71 to 106 fl oz / 1000 ft row per chisel) preferably in the fall when the soil is warm (50-85 F) at 6 inch depth and moist. Inject the fumigant at a depth of 10 to 12 inches with a soil injector or similar device. Follow directions for soil preparation, sealing, exposure, and aeration as specified elsewhere in this label.
- NOTE** To control symphylans (garden centipedes) use only overall at 18 or more gallons per acre and top dressing areas with manure partly fall when the soil is warm. To control wireworms use the higher dosages recommended for nematodes in very hard soil and treat treatments.

White Potatoes in Northwestern States: Use TELONE II as a spring or preferably a fall treatment to control

fumigating listed rows, seal in the
EXPOSURE PERIOD: After application exposure period will be required.
AERATION AND PREPARATION the soil to aerate completely by TELONE II is no longer evident time for each 10 gallons of TELONE II and shrub planting sites a 3 to 6 if heavy rains or low temperature treatment zone. After row treatment possible recontamination of the soil to the depth of the treatment zone of muck soils.

quackgrass and for suppression of potatoes. Apply as an overall treatment.

Time of Treatment
Spring
Fall

Mint in Northwestern States: Use overall treatment in the spring, or Verticillium wilt in disease infested weeks or until the odor of the fumigant Station or State Extension Service weed control and cultural practices. Verticillium wilt.

Important — Note Carefully. Fumigants in the soil. This is most likely if soils are either cold, wet, acid, or high in salts. Tests made after fumigation. To avoid soils do not use fertilizers containing salts above 65 F.

Certain crops including cotton, sugarcane, and alfalfa do not apply to them. When using high ammonia highly acid soils before application. Certain nursery vegetable crops such as cauliflower. To avoid this possible effect, if soil experience indicates a deficiency.

Attention: To avoid reinfestation of crop remains that could carry seeds. Since TELONE II soil fumigant is kerosene immediately after use. **Equipment made of aluminum, magnesium, or severely corrosive to such metals** will be penetrated readily by this material. Avoid contact with hands with polyethylene gloves. In outside storage, avoid bottom recessed areas.

Do not allow contamination of seed, feedstuffs, or domestic water. To avoid injury to fish and other bodies of water.

Rinse equipment and containers. Dispose of empty containers. Use this product only as specified.

NOTICE: Seller warrants that the product is safe when used in accordance with directions. **WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE** is hereby disclaimed, and the user assumes the risk of any such use.



355567

ALS, INC.

TELONE II FUMIGANT

INFORMATION

Use TELONE II to control plant parasitic nematodes (root-knot, beet, soybean) burrowing ring, spiral, sting and garden centipedes (symphylans). Fumigate with TELONE II under the conditions and at the rates and USE RECOMMENDATIONS and USE PRE-

parsnips	shallots
peas	spinach
peppers	squash (summer)
pimentos	squash (winter)
potatoes	sweet potatoes
pumpkins	swiss chard
radishes	tomatoes
rutabaga	turnips
salsify	watermelons

grass	sorghum
	soybeans
	sugar beets
	sugarcane
	tobacco
	vetch
	wheat

oranges	tangerines	tangelos
---------	------------	----------

	plums
	pomegranates
	prunes
	raisins
	walnuts

	strawberries
--	--------------

RECOMMENDATIONS

for Nematodes, and Wireworms

Crop	Dosage	
	Gallons Per Acre	Fl Oz. 1000 Ft. Row Per Chisel
Beans	4.5 to 6	46 to 64

Nursery Crops including floral plants, ornamentals, shrubs, and bushes, forest, shade, fruit and nut trees and vine and bramble fruits of all types

DIRECTIONS FOR USE

WHEN TO TREAT: Apply TELONE II either in the spring or fall, whenever soil type and conditions permit. For best results with annual crops, treat the soil each year. In northern states, late summer, or early fall treatment (before October 15) is best for land to be planted to early spring crops, especially transplanted crops such as celery, tomatoes, and nursery and orchard stock. Early fall treatment permits planting a fall cover crop. **Note:** Treat muck soils only in the early fall and plant as late as possible in the spring; treat fine textured (clay) soils only when they are near or at the wilting point. Do not use TELONE II to treat any type of soil when it is cold and or wet.

SOIL PREPARATIONS: TELONE II gives best results when conditions permit rapid diffusion of the fumigant through the soil and the soil surface can be sealed to prevent excessive fumigant loss during the exposure period. The soil should be in good seed bed condition, free of clods and undecomposed plant material, moisture at about one-half of field capacity, and temperature between 40° and 80°F at the depth of injection. If undecomposed plant debris is present, it should be plowed down and allowed to decompose before applying TELONE II. Tillage deeper than 12 inches is necessary for good fumigant penetration in soils where a hard or plow pan occurs at plow depth. Where deep tillage is used, the tillage equipment may often be modified to simultaneously apply the fumigant and thus avoid going over the field twice.

APPLICATION: TELONE II may be applied either as an overall (broadcast) or row treatment, using suitable application equipment that will ensure placement of the fumigant at least 6 to 8 inches below the final soil surface. For overall application use either plowsole or chisel equipment with the chisels spaced 12 inches apart. When the fumigant is injected at a depth of 12 inches or more (deep tillage), the chisel spacing may be up to twice the application depth but should not exceed 30 inches. Application may be made in the same direction or at an angle to the direction of the planting row, whichever is most convenient. For row application, use chisel equipment with one chisel per row or two chisels spaced 12 inches apart to treat only the soil where the crop is to be planted. When one chisel per row is used, adjust the fumigant flow rate to distribute about 1 1/2 times more fumigant per chisel than is recommended for overall application. When two chisels are used per row, apply at the same flow rate per chisel as for overall. In both cases, the amount of fumigant required per acre will decrease as the distance between rows is increased and vice versa. At time of planting, avoid placing the seed row directly over the furrow left by the applicator chisel. When a single chisel is used per row, place the seed 3 to 4 inches to one side of the chisel furrow. When two chisels are used, plant in the center of the area between the chisel furrows.

SEALING: Immediately after application, compact the soil surface to prevent excessive fumigant loss. After chisel application, use a roller, cultipacker, or similar sealing device. After plow-sole application, disk the land, then compact it by floating or rolling. Sealing after row application can be accomplished by the tractor wheel, by listing, or by bedding so that the fumigant will be 12 to 14 inches below the top of the bed. When fumigating listed rows, seal in the fumigant with ring rollers, press sealers, or by re-listing.

EXPOSURE PERIOD: After application and sealing, leave the soil undisturbed for 7 to 14 days. A longer exposure period will be required if the soil becomes excessively cold or wet during the exposure period.

AERATION AND PREPARATION OF SOIL BEFORE PLANTING: At the end of the exposure period allow the soil to aerate completely before planting the crop. Aeration is usually complete when the odor of TELONE II is no longer evident. Under optimum soil and weather conditions, allow one week of aeration time for each 10 gallons of TELONE II applied per acre. When TELONE II is used for treating deep-rooted tree and shrub planting sites, a 3 to 6 months aeration period should be allowed. To hasten aeration, especially if heavy rains or low temperatures occur during the exposure period, work the soil to the depth of the treatment zone. After row treatment use a knife-like chisel in the bed without turning the soil, thus reducing possible recontamination of the treated soil. To hasten aeration after overall treatment, plow or deep cultivate to the depth of the treatment zone. This is especially desirable in northern states after fall fumigation of muck soils.

quackgrass and for suppression of the damaging effects of Verticillium wilt in fields to be planted to white potatoes. Apply as an overall treatment according to the following directions:

Time of Treatment	Gallons per Acre	Fl. Oz./1000 Ft. Row per Chisel
Spring	17 to 25	50 to 73
Fall	25 to 34	73 to 100

Mint in Northwestern States: Use 59 gallons of TELONE II per acre (173 fl. oz./1000 ft. row per chisel) as an overall treatment in the spring, or preferably in the fall, to aid in the reduction of the damaging effects of Verticillium wilt in disease infested land to be used for mint production. After treatment allow at least 7 to 8 weeks or until the odor of the fumigant has left the soil before planting. Consult State Agricultural Experiment

DANGER

KEEP OUT OF REACH OF CHILDREN
HAZARDOUS LIQUID AND VAPOR
MAY BE FATAL IF INHALED,
ABSORBED THROUGH SKIN, OR SWALLOWED
CAUSES BURNS OF SKIN OR EYES
COMBUSTIBLE

Do Not Get in Eyes, on Skin, or on Clothing. Wear Eye and Skin Protection Necessary to Prevent Contact When Handling This Material. Wash Thoroughly after Handling and Before Eating and Smoking. Wash Contaminated Clothing and Air Contaminated Shoes Thoroughly Before Reuse. DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME. Use Only with Adequate Ventilation. Do Not Breathe Vapor.

In case of contact, immediately remove contaminated clothing and shoes and wash skin with soap and water. For eyes, flush with flowing water for at least 15 minutes and seek medical attention.

In case of spillage indoors, evacuate area and wear an approved full face mask equipped with a fresh blank canister meeting specifications of the U.S. Bureau of Mines for organic vapors. Use a full face self contained breathing apparatus or full face air supply respirator.

If inhaled, remove to fresh air and get medical attention if breathing has stopped. Get immediate medical attention. Keep patient absolutely quiet.

If swallowed, call a doctor immediately by giving information such as 2 tablespoonfuls of table salt in 1 glass of water.

Cameraman's Note

Poor Copy

ACTIVE INGREDIENT:
1,3-Dichloropropene 92%
INERT INGREDIENTS: 8%
E.P.A. Registration No. ~~XXXXXXXXXX~~ 35550-1
E.P.A. Est. No. ~~XXXXXXXXXX~~
~~XXXXXXXXXX~~ 35550-WA01
~~XXXXXXXXXX~~

AGRICULTURAL CHEMICAL

Do Not Ship or Store with
Food, Feeds, or Clothing

PRECAUCION AL USUARIO: Si usted no lee
inglés, no use este producto hasta que la eti-
queta le haya sido explicada ampliamente.

wheat
 oranges
 tangerines
 tangelos
 plums
 pomegranates
 prunes
 quince
 walnuts
 youngberries

wheat, by listing, or by covering so that the fumigant will be 12 to 14 inches below the top of the bed. When fumigating listed rows, seal in the fumigant with ring rollers, press sealers, or by re-listing

EXPOSURE PERIOD: After application and sealing, leave the soil undisturbed for 7 to 14 days. A longer exposure period will be required if the soil becomes excessively cold or wet during the exposure period.

AERATION AND PREPARATION OF SOIL BEFORE PLANTING: At the end of the exposure period allow the soil to aerate completely before planting the crop. Aeration is usually complete when the odor of TELONE II is no longer evident. Under optimum soil and weather conditions, allow one week of aeration time for each 10 gallons of TELONE II applied per acre. When TELONE II is used for treating deep-rooted tree and shrub planting sites, a 3 to 6 months aeration period should be allowed. To hasten aeration, especially if heavy rains or low temperatures occur during the exposure period, work the soil to the depth of the treatment zone. After row treatment use a knife-like chisel in the bed without turning the soil, thus reducing possible recontamination of the treated soil. To hasten aeration after overall treatment, plow or deep cultivate to the depth of the treatment zone. This is especially desirable in northern states after fall fumigation of muck soils.

quackgrass and for suppression of the damaging effects of Verticillium wilt in fields to be planted to white potatoes. Apply as an overall treatment according to the following directions:

Time of Treatment	Gallons per Acre	Fl. Oz./1000 Ft. Row per Chisel
Spring	17 to 25	50 to 73
Fall	25 to 34	73 to 100

Mint in Northwestern States: Use 59 gallons of TELONE II per acre (173 fl. oz./1000 ft. row per chisel) as an overall treatment in the spring, or preferably in the fall, to aid in the reduction of the damaging effects of Verticillium wilt in disease infested land to be used for mint production. After treatment allow at least 7 to 8 weeks or until the odor of the fumigant has left the soil before planting. Consult State Agricultural Experiment Station or State Extension Service Specialists for the use of other practices such as flaming the stubble, weed control, and cultural practices when using TELONE II as an aid to reducing damage caused by Verticillium wilt.

USE PRECAUTIONS

Important — Note Carefully. Fumigation may temporarily raise the level of ammonia nitrogen and soluble salts in the soil. This is most likely to occur when heavy rates of fertilizer and fumigant are applied to soils that are either cold, wet, acid, or high in organic matter. To avoid injury to plant roots, fertilize as indicated by soil tests made after fumigation. To avoid ammonia injury or nitrate starvation, or both, to crops on high organic soils do not use fertilizers containing nitrates until after the crop is well established and the soil temperature is above 65 F.

Certain crops including cotton, sugarcane, and pineapple are tolerant to ammonia and the above rule does not apply to them. When using high rates of TELONE II as required by certain state nursery regulations, liming of highly acid soils before fumigation may stimulate nitrification and reduce the possibility of ammonia toxicity. Certain nursery crops such as citrus seedlings, *Cornus sp.*, *Crataegus sp.*, spruce, and vegetable crops such as cauliflower have shown evidence of phosphorus deficiency following fumigation. To avoid this possible effect, it is suggested that additional phosphate fertilizer be used on soils where experience indicates a deficiency may occur.

Attention: To avoid reinfestation of treated soil do not use irrigation water, transplants, tools, seed pieces, or crop remains that could carry soilborne pests from infested land. Clean equipment carefully before using. Since TELONE II soil fumigant is corrosive under certain conditions, flush all applicators with fuel oil or kerosene immediately after use. **DO NOT USE WATER. Do not use containers, pumps, or other transfer equipment made of aluminum, magnesium or their alloys, as under certain conditions TELONE II may be severely corrosive to such metals.** Common protective clothing, such as rubber gloves and boots, etc., may be penetrated readily by this material. Polyethylene provides a good barrier. For field operations cover shoes and hands with polyethylene bags. Store TELONE II in tightly closed containers in a cool place away from dwellings. In outside storage, store drums on their sides to avoid accumulation of rain water in the top or bottom recessed areas.

Do not allow contamination of seeds, plants, fertilizers, or other pesticide chemicals. Do not contaminate food, feedstuffs, or domestic water supplies.

To avoid injury to fish and other wildlife, do not spill or empty TELONE II into streams, ponds, or other bodies of water.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Dispose of empty containers by punching holes in them and burying with wastes.

Use this product only as specified on this label.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the use of any such use.

ACTIVE INGREDIENT:
 1,3-Dichloropropene 92%

INERT INGREDIENTS: 8%

E.P.A. Registration No. ~~XXXXXXXXXXXX~~ 35550-1

E.P.A. Est. No. ~~XXXXXXXXXXXX~~ 35550-WA01

AGRICULTURAL CHEMICAL

Do Not Ship or Store with Food, Feeds, or Clothing

PRECAUCION AL USUARIO: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

TRANSLATION: (TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

DIRECTIONS
and Wireworms:

Dosage	
Gallons Per Acre ¹	Fl. Oz. 1000 Ft. Row Per Chisel ²
4.5 to 6	46 to 62
9 to 12	93 to 124
9 to 15 ²	26 to 44
24 ³ to 36	71 to 116
24 to 36	71 to 116
6	62
12 to 15	35 to 44
9	93
12 to 18	35 to 51
24 to 36	
36	106

Time to Penetrate Various Depths		
4 ft	5 ft	6 ft
21	27	33
30	36	44
51	63	75
69	84	102

¹ Gallons per acre.
² Row per chisel.
³ Row per chisel.
 Use of plants such as tree fruit, rowing, etc., in the northwestern states.
 TELONE II is the trade name for aldrin, DDT, and dieldrin.
 Use by application as a preplant, preplant, or post-plant treatment.
 4 per chisel, preferably in the fall when the soil is 12 to 14 inches with chisels mounted in the row centers.
 Use elsewhere as directed on this label.
 100 lbs per acre and apply to 100 ft. rows or as recommended for row spacing, etc., in the label.
 or preferably a fall treatment to control

"Distributed By"
QUINCY FARM CHEMICALS, INC.
 P. O. Box 307
 Quincy, Washington 98848