

35571-1

8-23-1977

1/1



CHEM PRO

ALGAE CONTROL ED LIQUID

ACTIVE INGREDIENT

Poly (oxyethylene (dimethyliminio) ethylene-(dimethyliminio) ethylene dichloride) 15.0%

INERT INGREDIENT 85.0%

This product contains 1.29 lbs. of active ingredient per gallon and weighs 8.58 lbs. per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes or clothing.

PRACTICAL TREATMENT (FIRST AID): If swallowed, promptly drink a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams or ponds. Permits may be required for discharges containing this pesticide into lakes, streams, ponds or public water. For guidance, contact the regional office of the Environmental Protection Agency.

DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Chem Pro Algae Control ED Liquid is used to control algae, bacteria, and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove algal growth, microbiological slime and other deposits. An initial slug addition of 4.0 to 10.0 fluid ounces of Chem Pro Algae Control ED Liquid per 1000 gallons of water to provide a concentration of 32 to 80 parts per million of Chem Pro Algae Control ED Liquid based on the total weight of water in the system is recommended. Repeat initial dosage until control is evident. Subsequent slug additions of 1.0 to 10.0 fluid ounces of Chem Pro Algae Control ED Liquid per 1000 gallons of water (8 to 80 parts per million of Chem Pro Algae Control ED Liquid) should be employed every 2 to 5 days or as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the microbiological problem. Slug additions should be made in the sump of the water cooling tower.

STORAGE AND DISPOSAL: Keep container closed when not in use. Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Do not reuse empty container. Triple rinse (or equivalent) and dispose of container in an incinerator or in an approved landfill, or bury in a safe place. Rinsate that cannot be used or reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Open dumping is prohibited.

Manufactured By
CHEM PRO LABORATORY, INC.

941 West 190th Street
Gardena, California 90248

EPA REG. NO. 35571-1, EPA EST. NO. 35571-A-01
NET CONTENTS: ACCEPT

23 AUG 1977

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR THE NATIONAL POISON REGISTER
EPA REG. NO. 35571-1