


Comments
 3-21-55
 3-21-55
 3-21-55



SUPER CONCENTRATED

ALGAECIDE

Helps control of ALGAE in swimming pools

ACTIVE INGREDIENTS: CHLORINE 4.0%
 DIMETHYL BENZYL AMMONIUM CHLORIDE 1.0%
 NET WEIGHT: 32 FL. OZ.

DANGER: KEEP OUT OF REACH OF CHILDREN

See additional precautions and first aid on back panel.

NET: 1 QT. (32 FL. OZ.)

EPA REG. NO. 6541-B EPA EST. NO. 15879-CA-1

SPP SUPER CONCENTRATED ALGAECIDE

SPP Super Concentrated Algaecide is compatible with all pool water treatments. It is used to control algae growth with SPP Algaecide. The chlorine demand of the pool may be significantly reduced. Apply this product only as specified on the label.

— DIRECTIONS —

1. In freshly filled pools or pools showing no visible algae growth, apply along pool edges to the water line (1) gallon of SPP Algaecide per 50,000 gallons of water to prevent algae growth.
2. If algae growth is observed, add one (1) gallon SPP Algaecide for each 20,000 gallons of water to kill and control these algae species most commonly found in swimming pools. When existing growth is dead, clean the pool, drain and refill if necessary, and add one gallon SPP Algaecide per 50,000 gallons of water.
3. Maintenance Dose: Use one (1) quart per 50,000 gallons of water every 5-7 days. When weather is hot or bathing load is heavy, use this dosage every 3 days or as necessary.

— DANGER —

WARNING: KEEP OUT OF REACH OF CHILDREN: CORROSIVE causes severe eye and skin damage. Do not get into eyes, on skin or clothing. Protect eyes and skin. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink PROMPTLY a large quantity of milk and whites, gelatin solution or if these are not available, drink large quantities of water. Avoid Alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measured against cerebral toxicity, shock, respiratory depression and convulsions may be needed. If persistent convulsions may be controlled by the cautious intravenous injection of a short acting barbiturate.

DO NOT REUSE CONTAINER — DESTROY WHEN EMPTY
 This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, rivers or public waters. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

Manufactured by CHEM-TAB CHEMICAL CORP., 1111 Beach Blvd.

ACCEPTED WITH COMMENTS