

CAUTION:

Keep Out of Reach of Children. May be Harmful if Swallowed. Skin and eye contact may cause severe irritation. Avoid contact with skin and eyes. In case of contact flush with plenty of water. Do not use ammonia or other bowl cleaners with this product. To do so will release hazardous gases.

FIRST AID

EXTERNAL—Wash with water

INTERNAL—Drink mucilage, raw egg white, milk or rice gruel. Follow with emetic to induce vomiting.

CALL A PHYSICIAN

ENVIRONMENTAL HAZARD WARNING


This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds, or public water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

SODIUM HYPOCHLORITE
CAUTION: KEEP OUT OF REACH OF CHILDREN

CONTENT

Active Ingredients: Sodium Hypochlorite 10.5%
Inert Ingredients: 89.5%

ACCEPTED
35639-1
SEP 17 1975
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO.



AQUA CHEMICAL CORPORATION
EPA Reg. No. 35639-1

GENERAL OFFICES

13501 BALTIMORE BLVD. (Rt. 1) • LAUREL, MARYLAND 20810
WASHINGTON CALL (301) 953-3788 BALTIMORE CALL (301) 792-7255

15 GALLONS

The
The
after
For
beld
men
to
to
suit
ched
stan
and
foild

SU
App
wat
be s
gall
appl
bein
3.0

RITE
CHILDREN

ACCEPTED
35639-1
SEP 17 1975
BY THE FEDERAL INSECTICIDE
AND RODENTICIDE ACT
ECONOMIC POISON REGISTER-
NUMBER NO.
ATTENTION

RYLAND 20810
LL (301) 792-7255

The chlorine in swimming pools tends to dissipate in direct sunlight. Therefore it is best to make an application in the evening, preferably after use of the pool has been discontinued for the day.

For newly filled pools add quantity that is indicated by size of pool below. 10 oz. per 10,000 gallons yields .8 ppm, additional 1 oz. increments raises level .1 ppm. Repeat dosage until a chlorine residual of 0.6 to 1.0 is obtained. Repeat dosage daily or as needed to maintain 0.6 to 1.0 ppm. chlorine residual at all times. Test water frequently with suitable test kit following the kits directions. Before entering pool, check the chlorine residual and if above 3.0 ppm. allow the pool to stand until the residual drops. Maintain pH of the water between 7.2 and 7.6 for best results. Test frequently with suitable test kit and follow directions for corrections.

SUPER CHLORINATION:

Approximately every 7 to 10 days, it is recommended that the pool water be superchlorinated. Prior to super chlorination, the pool should be swept or vacuumed to remove dead algae. Add 2 quarts per 10,000 gallons of water will raise the residual to 5.0 ppm. For best results application should be initiated in the evening with the pool filter being turned off until morning. When the residual has dropped to 3.0 ppm. or less the pool is safe for use.

