

CP CHEMICALS, INC.

BASIC COPPER TS-53-WP  
Copper Sulfate, tribasic  
Wettable Powder

For control of certain fungal and bacterial diseases of fruits, nuts, vegetables, field crops and ornamentals indicated below. Do not apply to plants which are dormant or to plants which are in flower. Do not spray during rain or when plants are wet. Do not spray directly before or after a sulfur, borax, or other sulfocarbamate fungicide.

DO NOT USE THIS PRODUCT UNDILUTED. DAMAGE TO CROP WILL RESULT.

DIRECTIONS FOR USE

(1) In a solution of Federal law use this product in a manner which complies with the labeling.

SPRAY PREPARATION: Mix 1/2 cup (125 g) of product with 1 gallon (3.8 liters) of water. For use in a sprayer, add 1/2 cup (125 g) of product to 100 gallons (3.8 kiloliters) of water. For use in a backpack sprayer, add 1/2 cup (125 g) of product to 20 gallons (75.7 liters) of water. For use in a hand sprayer, add 1/2 cup (125 g) of product to 5 gallons (18.9 liters) of water. For use in a mist blower, add 1/2 cup (125 g) of product to 10 gallons (37.9 liters) of water. For use in a pump sprayer, add 1/2 cup (125 g) of product to 10 gallons (37.9 liters) of water. For use in a backpack sprayer, add 1/2 cup (125 g) of product to 20 gallons (75.7 liters) of water. For use in a hand sprayer, add 1/2 cup (125 g) of product to 5 gallons (18.9 liters) of water. For use in a mist blower, add 1/2 cup (125 g) of product to 10 gallons (37.9 liters) of water. For use in a pump sprayer, add 1/2 cup (125 g) of product to 10 gallons (37.9 liters) of water.

Recommended Spray Schedules

Quantities are given for conventional spraying. For use in a backpack sprayer, divide quantities by 20. For use in a hand sprayer, divide quantities by 5. For use in a mist blower, divide quantities by 10. For use in a pump sprayer, divide quantities by 10.

AGRICULTURAL  
Fruits and Nuts

ALMONDS: For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

APPLES: For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

APRICOTS: For anthracnose (conspicuous blight) use 4 to 6 pounds/100 gallons with 1/2 cup (125 g) of spreader-sticker as fall dormant spray (October to November 15-Jan 15). Repeat in red bud stage. For brown rot, use 3 pounds/100 gallons at red bud, early bloom, and late bloom. For scab, use 3 pounds/100 gallons. Do not spray after petal fall during growing season while trees in leaf. For scab, apply 3 pounds/100 gallons when recommended by State Extension Service Specialists.

AVOCADOS: For anthracnose, blight (conspicuous), scab, use 3 pounds/100 gallons; spray first when blossom buds open. Make 3 applications at four-week intervals. LIMITS: 30 to 40 pounds/Acre. Do not apply more than 1 1/2 days after bloom.

BLACKBERRIES, BOYSENBERRIES, DOWDBERRIES, LOGANBERRIES, RASPBERRIES: For leaf spot (conspicuous), use 2 pounds/100 gallons. Spray first when leaf buds swell, again when leaves begin to open. For anthracnose, use 2 pounds/100 gallons. Spray when leaf buds begin to open, again when leaves begin to open. Repeat 10 to 14 days after first spray. For brown rot, use 2 pounds/100 gallons. Do not spray after petal fall during growing season while trees in leaf. For scab, apply 3 pounds/100 gallons when recommended by State Extension Service Specialists.

CHERRIES (sour) - Sweet cherry: 3 to 4 pounds/100 gallons. Apply when buds begin to bloom and late bloom stages. Leaf spot: 2 to 3 pounds/100 gallons plus 4 to 6 pounds/100 gallons of spreader-sticker. Apply at petal fall and make 1 or 2 post-bloom applications. For brown rot, use 3 pounds/100 gallons. Apply at petal fall and second stage. For scab, use 3 pounds/100 gallons. Do not spray after petal fall during growing season while trees in leaf. For scab, apply 3 pounds/100 gallons when recommended by State Extension Service Specialists.

CITRUS: For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

COCONUTS: For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

CUCUMBERS: For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

CURRANTS, GOOSEBERRIES - anthracnose, leaf spot: (a) spray with 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. (b) spray 6 pounds/100 gallons at 10-14 day intervals. (c) spray 6 pounds/100 gallons at 10-14 day intervals, and after harvest. LIMITS: 14 pounds/Acre.

GRAPEFRUIT: See Citrus.

GRAPES: For anthracnose, 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) on dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

GUAVAS, LEMON, LIME: See Citrus.

OLIVES: For leaf spot/peacock spot, cyclanoptera, 4 to 6 pounds/100 gallons applied in late October. LIMITS: 6 to 12 pounds/Acre.

NECTARINES: See Peaches.

ORANGES: See Citrus.

PEACHES, NECTARINES: For bacterial blight (the blight of peach), leaf spot (shear blight), apply dormant spray at 4 pounds/100 gallons plus 1/2 cup (125 g) of spreader-sticker (100 gal) between Nov. 15 and Dec. 15 or before leaf fall. For brown rot, anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray during growing season while trees in leaf. For scab, apply 3 pounds/100 gallons when recommended by State Extension Service Specialists.

PEARS: For blight, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at 10% bloom, repeat every five days or after each rain until end of leaf bloom. For leaf spot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at 10% bloom, repeat every five days or after each rain until end of leaf bloom. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

PEARS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

PEARS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

PEARS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

PEARS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

CARROTS: For downy mildew and leaf spots, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

CAULIFLOWERS: For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

CUCUMBERS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

EGGPLANT: For anthracnose, downy mildew, early blight, late blight, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

LETTUCE: For anthracnose, leaf spot (conspicuous), downy mildew, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

ONIONS: For downy mildew, leaf spots, use 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

PEAS: For downy mildew, leaf spots, use 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

PEPPERS: For anthracnose, bacterial spot, early & late blight, downy mildew, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

POTATOES: For early and late blights, start spraying when plants emerge, repeat at 7 to 10 day intervals (every 5 days for severe late blight). Use from 2 to 3 pounds/100 gallons. LIMITS: 12 pounds/Acre, based on plant size and severity of disease.

SEEDLINGS, VEGETABLE: See vegetable seedlings.

SPINACH: For anthracnose, late mildew (downy mildew), leaf spot (conspicuous), white rust, use 2 to 3 pounds/100 gallons. Do not spray if disease is severe. Repeat every 7 to 10 days. Apply 1/2 cup (125 g) of spreader-sticker.

TOMATOES: For anthracnose, bacterial spot, bacterial canker (leaf blight), early & late blights, leaf mold, railroad wilt, septoria and stem-end rot, leaf spots - begin protective spray in start bed or 2 to 3 pounds/100 gallons. Repeat at 4, 7, 10 or 14 day intervals as required.

VEGETABLE SEEDLINGS: To prevent or control damping-off, apply 4 pounds/100 gallons, 8 pounds/Acre, to soil surface in 2 to 4 inch beds every 7 to 10 days.

AGRICULTURAL FIELD CROPS

PEANUTS: For leaf spot (conspicuous), use 2 to 3 pounds/100 gallons. Do not spray if disease is severe. Repeat every 7 to 10 days. Apply 1/2 cup (125 g) of spreader-sticker.

SUGAR BEETS: For downy mildew, leaf spot (conspicuous) - begin spraying when plants emerge, repeat at 7 to 10 day intervals. Apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) as required to cover. Repeat at 7 to 10 day intervals.

TABACCO: For mildew, 1 pound in 100 gallons of water/100 sq. ft. plant bed. Begin after sowing seed or when plants emerge. Repeat at 5 to 10 day intervals until transplanted.

AGRICULTURAL SEED TREATMENT

WHEAT - striking smut (Bunt): 2 ounces basic copper sulfate/100 lbs. of seed. 3 ounces on winter wheat, and 4 ounces on winter wheat. Apply as a dry mix to seed. DO NOT USE TREATED SEED FOR FOOD, FEED OR OIL PURPOSES.

AGRICULTURAL ORNAMENTAL CROPS & TREES

ARBOR VITAE, ASTERS - blights, leaf spots: 3 to 6 pounds basic copper sulfate/100 gallons of dust, as well as 100 to 200 gallons in early fall and repeat in late fall.

BIRCH - leaf blights, leaf spots: 3 pounds/100 gallons/Acre. Apply at 7 to 10 day intervals.

For diseases on plants listed below, apply 4 pounds of basic copper sulfate/100 gallons, or 4 pounds in a dust form. Begin before disease appears. Repeat at 7 to 10 day intervals at 3 week intervals.

AZALEAS, BEGONIAS, RHODODENDRONS - leaf blight, leaf spot, leaf spot (and leaf gall on azaleas)

BOXWOOD - mildew

CANONAS, LILIES, TULIPS - botrytis blight

CHRYSAANTHEMUMS - botrytis blight, downy mildew, leaf spots

DANIELS - botrytis blight, leaf spots

FLOWERING SHRUBS - leaf spots

GERANIUMS - botrytis blight, downy mildew

GLADIOLI - botrytis blight, downy mildew, leaf spots

GRASSES, lawn (Florida) - gray leaf spot

HOLLYHOODS - downy mildew, leaf spots

IRIS, PANSEYS, VIOLETS, VIRGINIA CREEPER - downy mildew

IRIS - bacterial blight

NASTURTIUMS - anthracnose, leaf spots

PEONIES - botrytis blight, leaf spots

ROSES - anthracnose, leaf spots, downy mildew, leaf spots, leaf gall

SHRUBS, FLOWERING - leaf spots

SNAPDRAGONS - anthracnose, leaf spots, leaf spots

SWEET PEAS - downy mildew, leaf spots, leaf spots

ZINNAS - anthracnose, leaf spots, leaf spots

ZINNAS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

ZINNAS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

ZINNAS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

ZINNAS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

ZINNAS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

ZINNAS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

ZINNAS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

ZINNAS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

ZINNAS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

ZINNAS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

ZINNAS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

ZINNAS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

ZINNAS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

ZINNAS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

ZINNAS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.


ZINNAS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

ZINNAS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

ZINNAS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

ZINNAS (continued): For anthracnose, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) to dormant spray (November to December 15) repeat in spring at petal fall and second stage. For brown rot, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. For scab, apply 1/2 cup (125 g) plus 1/2 cup (125 g) of spreader-sticker (100 gal) at petal fall and second stage. Do not spray after bloom. Do not spray after bloom.

Made in U.S.A.



50 LBS.  
(22.7 Kg)  
NET

# CP-BASIC COPPER TS-53-WP

## WETTABLE POWDER AGRICULTURAL FUNGICIDE

Active Ingredient ..... 51%  
Copper as metal  
Inert Ingredients ..... 49%  
Insecticides, etc. (see label)

**KEEP OUT OF REACH OF CHILDREN**

**WARNING**

Hazardous to humans and animals.

IRITANT: May cause temporary eye irritation. May irritate respiratory tract, nose, throat, and skin. May cause dizziness, headache, and nausea. Avoid contact with eyes, nose, mouth, and skin. Avoid contact with food, feed, or drift to other crops or forage.

**FIRST AID**

If in contact, immediately flush with plenty of water for at least 15 minutes. For eye contact, flush with water for at least 15 minutes. If swallowed, drink plenty of water. If inhaled, get fresh air. If these are not available, drink large quantities of water. Call a doctor for a physical examination.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Keep out of lakes, streams or ponds.

**DIRECTIONS FOR USE — SEE BACK OF BAG**  
(It is a violation of Federal law to use this product in a manner inconsistent with this label.)

**STORAGE & DISPOSAL**

**Prohibitions:** Keep pesticide only in original container. Do not contaminate water, food or feed by storage or disposal. Open burning and dumping prohibited. Do not reuse empty container.

**Disposal:** Pesticide or residue that cannot be used or chemically reprocessed should be disposed of according to procedure approved by federal, state, or local disposal authorities.

**Container Disposal:** Consult federal, state, or local disposal authorities for approved procedures.



Manufactured by  
**CP CHEMICALS, INC.**



POWDER SPRINGS, GA 30075  
SEWAREN, NJ 07077, USA  
ESTABLISHMENT NO. 35896-GA-1  
35896-NJ-1  
EPA REGISTRATION NO. 35896

Net Contents  
50 lbs  
(22.7 kg)

100

**CP-BASIC COPPER TS-53-WP**



100