



FIXED COPPER FLOWABLE AGRICULTURAL FUNGICIDE

Active Ingredient
Copper (in Basic Copper Sulfate)
expressed as metallic
Inert Ingredients
(contains one pound Fixed Copper per gallon)

KEEP OUT OF REACH OF CHILDREN

WARNING

Hazards to humans and domestic animals:

IRRITANT: May cause eye damage and skin or nasal irritation. Do not get in eyes or nose, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful if swallowed. Avoid contamination of food or water, or drift to other crops or forage.

FIRST AID

In case of contact, immediately flush eyes (under lids) or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, or gelatin solution; if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds.

For control of certain fungal and/or bacterial diseases of fruits, nuts, vegetables, field crops and ornamentals indicated below. Do not apply to any crop not listed on this label. Consult State Experiment Station or State Extension Service Specialists for recommended spray schedules for your local area. Follow a regular schedule of application and spray thoroughly. Do not use with or immediately before or after lime, sulfur, ferbam, or other dithiocarbamate fungicides.

DO NOT USE THIS PRODUCT UNDILUTED. DAMAGE TO CROP WILL RESULT.

DIRECTIONS FOR USE

(It is a violation of federal law to use this product in a manner inconsistent with its labeling.)

SPRAY PREPARATION: Half-fill spray tank with water. Start agitator; add softener to very hard water to prevent curdling. Add other desired adjuvants. Stir in required amount of MICROBLU® Fixed Copper Flowable gradually to permit smooth suspension while filling tank with balance of water. Keep agitator running during filling and until tank has been emptied. Rinse tank thoroughly.

Recommended Spray Schedules

Quantities are given for conventional spray rig having 100 gal. tank. Single application limits per acre are shown; for example, 12 lb. limit allows 300 gal. of 4 lbs./100 gal. spray per acre, per application.

CITRUS: For Brown rot, septoria leaf, fruit spot, apply 1/4 to 3/4 gal. plus 4 pounds hydrated lime and 3 pounds zinc sulfate per 100 gallons; apply before fall rains or at first sign of disease. Spreader-sticker may be added. Use lowest rate in California where copper injury may be a problem or groves are fumigated with hydrogen cyanide; use highest rate elsewhere.
LIMIT: 30 gal. Acre per application

For greasy spot, melanose, scab, apply 3/4 to 1-1/2 gal. in dormant spray before flush begins; repeat at 2/3 petal fall. Same limits apply.

AGRICULTURAL VEGETABLE CROPS

CELERY: For early blight (*cercospora*), late blight (*septoria*), leaf spots: begin with about 1 gal./100 gal. in plant bed or field when plants are 6 inches high or diseases first appear. Repeat in 5 to 14 days as needed, building up to 2-1/2 gal./100 gal. as plants grow.

LIMIT: 3 to 10 gal./Acre

PEPPERS: For anthracnose, bacterial spot, early & late blight, *cercospora* leaf spot (frog-eye spot), downy mildew, spray in plant bed before diseases appear. Continue in field at 7 to 10 day intervals, especially during fruiting stage. Use 1 to 2 gal./100 gal.

LIMIT: 2 to 4 gal./Acre

TOMATOES: For anthracnose, bacterial spot, bacterial canker (Southeast), early & late blights, leaf mold, nailhead rust, septoria and stemphylium leaf spots — begin protective spray in plant bed or field before diseases appear. Use 1 to 2 gal./100 gal., up to 6 gal./Acre.

LIMIT: Repeat at 4, 7, 10, or 14 day intervals as required.

AGRICULTURAL FIELD CROP

PEANUTS: For leaf spot (*cercospora*), use 1 to 1-1/2 gal./Acre when disease first appears. Addition of sulfur at State-recommended level will improve spot and powdery mildew control. Use lower (1 gal.) rate on small plants. Repeat at 10 to 14 day intervals.*

*Consult your county agricultural extension agent for current state recommendations as to amounts of basic copper sulfate and/or sulfur to be used as spray.

WARRANTY

Product warranted to meet label specifications. Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of the use or handling not in accordance with label directions and accepts the product on these conditions. Nothing hereon should be construed as authorization to practice any unexpired patent.

STORAGE AND DISPOSAL

Prohibitions: Keep pesticide only in original container. Do not contaminate water, food or feed by storage or disposal. Open burning and dumping prohibited. Do not reuse empty container.

Disposal: Pesticide or rinsate that cannot be used or chemically reprocessed should be disposed of according to procedures approved by federal, state, or local disposal authorities.

Container Disposal: Consult federal, state, or local disposal authorities for approved procedures.



Manufactured for
CP CHEMICALS, INC.
SEWAREN, NEW JERSEY 07077, U.S.A.



EPA ESTABLISHMENT NO. 35896-NJ-1

EPA REGISTRATION NO. 35896-14

NET CONTENTS

15 GAL.

150 LBS.

LOT NO

05