



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

35915-21

Date of Issuance:

8/10/16

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Tetraconazole + Thiophanate-Methyl

Name and Address of Registrant (include ZIP Code):

Nicole O'Loughlin
Agent for Oxon Italia S.p.A.
c/o Pyxis Regulatory Consulting, Inc.
4110 136th St. Ct. NW
Gig Harbor, WA 98332

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Shaja Joyner, Product Manager 20
Fungicide & Herbicide Branch, Registration Division (7505P)

Date:

8/10/16

2. Be aware that proposed data requirements have been identified in a Final Work Plan EPA-HQ-OPP-2015-006. For more information on these proposed data requirements, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division:
<http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>
3. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. A one year study is required to satisfy these data requirements. You have 18 months from the date of registration to provide these data.
4. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 35915-21.”
5. Submit one copy of the final printed label for the record before you release the product for shipment.


Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 05/05/2016
- Alternate CSF 1 dated 05/05/2016
- Alternate CSF 2 dated 05/05/2016

If you have any questions, please contact Lisa Pahel by phone at (703)347-0459, or via email at pahel.lisa@epa.gov.

Enclosure: Product Chemistry Review dated 06/20/2016, DP#434064; Similarity Clinic Memorandum dated 05/25/2015, DP#433566



Group	1	3	Fungicides
-------	---	---	------------

Tetraconazole + Thiophanate-Methyl

For Control and/or Suppression of the listed diseases in Pecans, Peanuts, Soybeans, and Sugarbeet.

ACTIVE INGREDIENTS:

Thiophanate-methyl: dimethyl [1,2-phenylenebis(iminocarbonothioyl)bis[carbamate] 21.27%
 Tetraconazole: 1-[2-(2,4-dichlorophenyl)-3-(1,1,2,2-tetrafluoroethoxy) propyl]-1H-1,2,4-triazole. 4.20%
OTHER INGREDIENTS 74.53%
TOTAL..... 100.0%

Contains 1.98 pounds of thiophanate-methyl and 0.39 pounds of tetraconazole per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (If you do not understand the label, find someone to explain it to you in detail).

FIRST AID	
IF INHALED:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
Emergency phone numbers	(800) 424-9300 CHEMTREC (transportation and spills) (800) 222-1222 Poison Control Center

EPA Registration No. 35915-ER
 EPA Establishment No. _____

Net Contents _____ [gallons] [gal.] [liters]

Manufactured for:
OXON ITALIA S.p.A.
 Via Sempione, 195 – 20016 PERO
 MILANO – ITALY

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if inhaled. Avoid breathing spray mist. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, and viton ≥ 14 mils.

Applicators and other handlers must wear:

- Long sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves for all mixes and loaders and for applicators using hand held equipment and chemical resistant aprons for mixers, loaders, and other handlers exposed to the concentrate.

USER SAFETY REQUIREMENTS

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands thoroughly with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product may be toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms adjacent to treatment areas. Exercise care when making applications of this product, and do not apply when atmospheric conditions favor drift or runoff. Do not contaminate water when disposing of equipment washwater or rinsate.

In order to mitigate concern for reproductive effects to endangered bird and mammal species which may occur incidentally in sugarbeet growing areas, you are required to ascertain through the state Department of Agriculture, or Cooperative Extension Service, whether the treatment area may contain habitat of federally listed bird and mammal species; if so, treatment must be avoided in these areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through

drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the REI. The REI for each crop is listed in the directions for use associated with that crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

USE INFORMATION

Apply this product in a water carrier by spraying onto crop surfaces that are intended to be protected from disease. To obtain adequate coverage of typical agricultural crops, total spray volume will usually range from 20 to 150 gallons per acre (200 to 1400 liters per hectare) for dilute sprays, and 5 to 10 gallons per acre (50 to 100 liters per hectare) for concentrate ground sprays and aircraft applications. Both ground and aircraft methods of application are recommended.

Add this product to the spray tank while filling with water. Keep the agitator running when filling spray tank and during spray operations. When tank mixing this product with other pesticides, observe the more restrictive label limitations and precautions. Do not exceed any label dosage rates. This product cannot be mixed with any product containing a label prohibition against such mixing. Combination in the spray tank with other pesticides, fertilizers or surfactants is not recommended unless prior use has shown the combination to be physically compatible, effective and noninjurious under your conditions of use. When an adjuvant is to be used with this product, Advan LLC recommends the use of a Council of Producers and Distributors of Agrotechnology (CPDA) certified adjuvant.

Resistance Management Recommendations

This product contains thiophanate-methyl, a Group 1 fungicide, and tetraconazole, a Group 3 fungicide. Fungal isolates with acquired resistance to Group 1 and/or Group 3 may eventually dominate the fungal population if Group 1 and/or Group 3 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by this product and/or Group 1 and/or Group 3 fungicides.

To delay fungicide resistance consider:

- Avoiding the consecutive use of this product or other target site of action Group 1 and/or Group 3 fungicides that have a similar target site of action, on the same pathogens.
- Using tank-mixtures or premixes with fungicides from different target site of action Groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Basing fungicide use on a comprehensive IPM program.
- Monitoring treated fungal populations for loss of field efficacy.
- Contacting your local extension specialist, certified crop advisors, and/or local Advan LLC representative for fungicide resistance management and/or IPM recommendations for specific crops and resistant pathogens.

Rotational Crop Restrictions

Refer to the table below for the minimum time intervals required between the last application of this product and a new crop planting.

Crop	Rotational Interval (in days)
Corn	0
Grains, small (barley, buckwheat, millet, oats, rice, rye, triticale and wheat)	45
Grape	0
Peanut	0
Pecan	0
Soybean	0
Strawberry	0
Sugarbeet	0
Sugarcane	45
All other crops	120

CROPS

PEANUTS		
DISEASES CONTROLLED	RATE PER ACRE	APPLICATION DIRECTIONS
Early Leaf Spot <i>(Cercospora arachidicola)</i> Late Leaf Spot <i>(Cercosporidium personatum)</i> Limb Rot <i>(Rhizoctonia spp.)</i> Rust <i>(Puccinia arachidicola)</i> Web Blotch <i>(Phoma arachidicola)</i>	23 fl. oz.	Apply when conditions are favorable to disease, generally when leaf wetness first occurs, or 30 to 40 days after planting. Repeat applications on a 14-day schedule if conditions remain favorable for disease. Do not make more than 4 applications of this product per year. A multi-site mode of action fungicide, such as Echo (or other Chlorothalonil based products) should be used in combination with this product to reduce risk of resistance. Consult with your Extension Service representatives for guidance on the proper use of this product in programs which attempt to minimize the occurrence of disease resistance to fungicides.
RESTRICTIONS: <ul style="list-style-type: none"> • Pre-harvest Interval: 14 days (digging). • Restricted Entry Interval (REI): 24 hours. • Do not feed hay or threshings from treated fields to livestock. • Do not graze or feed cover crops grown in treated areas to livestock. • Do not make more than 4 applications of this product per year. • Do not apply more than 0.41 lbs of tetraconazole active ingredient and 1.4 lbs of thiophanate-methyl active ingredient per acre per year. 		

PECANS		
DISEASES CONTROLLED	RATE PER ACRE	APPLICATION DIRECTIONS
Brown Spot, (<i>Cercospora fusca</i>) Downy Spot, Leaf Blotch (<i>Mycosphaerella spp.</i>) Powdery Mildew, (<i>Microsphaera penicillata</i>) Scab, (<i>Cladosporium caryigenum</i>) Vein Spot and/or Liver Spot (<i>Gnomonia spp.</i>) Zonate Leaf Spot (<i>Cristulariella moricola</i>)	36 – 41 fl. oz.	Begin applications when conditions are favorable to disease and repeat at 14-21 day intervals until shuck split. Do not apply after shuck split. Apply the highest use rate when disease pressure is severe. Minimum recommended spray volume is 20 gallon for ground applications, 10 gallons for aerial applications. Alternate applications of this product with fungicides with a different mode of action. Consult with your Extension Service representatives for guidance on the proper use of this product in programs which attempt to minimize the occurrence of disease resistance to fungicides.
RESTRICTIONS: <ul style="list-style-type: none"> • Pre-harvest Interval: 30 days. • Restricted Entry Interval (REI): 3 days. • Do not graze or feed cover crops grown in treated areas to livestock. • Do not apply more than 0.5 lb of tetraconazole active ingredient and 2.1 lbs of thiophanate-methyl active ingredient per acre per year • Do not apply after shuck split. 		

SOYBEAN		
DISEASES CONTROLLED	RATE PER ACRE	APPLICATION DIRECTIONS
Anthracnose <i>(Colletotrichum spp.)</i> Brown Spot <i>(Septoria glycines)</i> Purple Seed Stain <i>(Cercospora kikuchii)</i> Frogeye Leaf Spot <i>(Cercospora sojina)</i> Pod and Stem Blight <i>(Diaporthe phaseolorum)</i> Powdery Mildew <i>(Microsphaera diffusa)</i> Powdery Mildew <i>(Microsphaera diffusa)</i>	20-23 fluid ounces	Make one application at early pod fill (R3 soybean growth stage). If environmental conditions are favorable to continued disease development, make a second application after 15 days dependent upon the severity of disease pressure. Apply the higher rate and reduce interval between applications when disease pressure is severe. Apply this product in a minimum of 10 gallons of spray suspension per acre by ground sprayer or in a minimum of 5 gallons of spray suspension per acre by aircraft.
White Mold/Sclerotinia Stem Rot <i>(Sclerotinia sclerotiorum)</i>	23 fluid ounces	Make one application at R2 stage (early bloom), if environmental conditions are favorable to continued disease development; make a second application after 14 days dependent upon the severity of disease pressure. Apply this product in a minimum of 10 gallons of spray suspension per acre by ground sprayer or in a minimum of 5 gallons of spray suspension per acre by aircraft. Full coverage of the flowers, stems and branches is required for disease control.
Asian Soybean Rust <i>(Phakopsora pachyrhizi)</i>	20-23 fluid ounces	Apply preventively when disease infection is likely to occur. Make a second application if conditions are favorable for disease infection before stage. Apply this product in a minimum of 10 gallons of spray suspension per acre by ground sprayer or in a minimum of 5 gallons of spray suspension per acre by aircraft.
RESTRICTIONS: <ul style="list-style-type: none"> • Do not make more than 2 applications per year. • Do not graze or feed treated forage or hay to livestock. • Do not harvest immature soybeans for consumption once plants are treated with this product. • Do not apply more than 1.4 lb of Thiophanate-methyl and 0.15 lb of tetraconazole per acre per year • Do not apply after soybean growth stage R5 (beginning seed). • Do not sure on vegetable soybean varieties grown for their immature pods. 		

SUGARBEETS		
DISEASES CONTROLLED	RATE PER ACRE	APPLICATION DIRECTIONS
Cercospora Leaf Spot <i>(Cercospora beticola)</i> Powdery Mildew (<i>Erysiphe</i>) Ramularia (<i>Ramularia beticola</i>)	32 fl. oz.	Apply when conditions are favorable for Cercospora Leaf Spot, Powdery Mildew, or Ramularia. Do not make more than one application of this product per year. To obtain adequate coverage of typical agricultural crops, total spray volume usually ranges from 20 to 150 gallons per acre for dilute sprays, and 5 to 10 gallons per acre for concentrate ground sprays and aircraft applications. For areas east of the Rocky Mountains: Do not make more than one application of this product per year. Follow resistance management guidelines.
RESTRICTIONS: <ul style="list-style-type: none"> • Pre-harvest Interval: 21 days • Restricted Entry Interval (REI): 24 hours. • Do not make more than one application of this product per year. • Do not apply more than 0.102 lb of tetraconazole active ingredient and 2.1 lbs of thiophanate-methyl active ingredient per acre per year. 		

OPTIONAL LABEL LANGUAGE: USE IN CALIFORNIA BY CHEMIGATION ONLY FOR SUGARBEETS.

CHEMIGATION INSTRUCTIONS

Apply this product only through one or more of the following types of systems: sprinkler (including center pivot), lateral move, end tow, side (wheel) roll, traveler, big gun, solid set or hand move irrigation system. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

For specific information about calibration, contact State Extension Service specialists, equipment manufacturers, or other irrigation experts.

Do not connect an irrigation system used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make the necessary adjustments should the need arise.

To prevent the movement of this product into the soil:

- Minimize pesticide contact with the soil surface by chemigating above the crop canopy.
- Stop chemigation when pesticide mixture is observed running off crop surfaces or after 0.25 inches

of water has been applied, whichever occurs first.

- Allow for sufficient time after chemigation for crop surfaces to dry prior to expected rainfall or to irrigation applied above the crop canopy.

Sprinkler Chemigation

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump motor when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

When mixing, fill nurse tank half full with water. Add this product slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. Stickers, spreaders, etc. should be added last. If compatibility is in question, use the compatibility jar test before mixing a whole tank.

Because of the wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products used in mixtures.

Add this product through a traveling irrigation system continuously or at the last 30 minutes of solid set or hand moved irrigation systems. Agitation is recommended.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a dry, temperature-controlled, secure place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on-site or at an approved waste disposal facility.

CONTAINER HANDLING:

[Containers < 5 Gallons:] Nonrefillable container. Do not reuse or refill this container. Triple rinse container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

[Containers ≥ 5 Gallons:] Nonrefillable container. Do not reuse or refill this container. Triple rinse container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

[Bulk Containers:] Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. When the container is empty, replace the cap and seal all openings that have been opened during use and return to the point of purchase or to a designated location named at the time of purchase of this product. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. Check for leaks after refilling and before transporting. Do not transport if this container is damaged or leaking. If the container is damaged or leaking, call Chem-Trec at (800) 424-9300.

If the container is damaged and leaking or material has been spilled, follow these procedures:

- Cover spill with absorbent material.
- Sweep into disposal container.
- Wash area with detergent and water and follow with clean water rinse.
- Do not allow to contaminate water supplies.
- Dispose of according to instructions.

If not returned to the point of purchase or to a designated location, clean empty container as instructed above and offer for recycling. Disposal of this container must be in compliance with state and local regulations.

THIS CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

WARRANTY AND LIMITATION OF DAMAGES

Conditions of sale: To the extent consistent with applicable law, Oxon Italia S.p.A. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to Oxon Italia S.p.A. Oxon Italia S.p.A. DISCLAIMS ALL OTHER WARRANTIES, EXPRESS OR IMPLIED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, OXON ITALIA S.P.A. SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, AND OXON ITALIA S.P.A. SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE. BUYER AND USER ACKNOWLEDGE AND ASSUME ALL RISKS AND LIABILITY RESULTING FROM HANDLING, STORAGE AND USE OF THIS PRODUCT. OXON ITALIA S.P.A. DOES NOT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTY, GUARANTEE OR REPRESENTATION CONCERNING THIS PRODUCT.

Tetraconazole + Thiophanate-Methyl (EPA Reg. No. 35915-xx) (Application 04-15-2016)