### A HARD VINYL ANTI-FOULING PAINT

Commence of the state of the st

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labelling

#### CTIVE INGREDIENT

Bis (tributyltin Admate 19 14%

NERT I'NGREDIENTS 80.86%
PA Establishment No. 35923-FL-1

A Registration No. 35923-9

## WARNING:

paints

ee side panel for additional recautionary statements.

#### REPAINT WORK

- 1 If OLD coating is a VINYL composition, in GOOD condition
  - A Clean thoroughly and let DRY;
  - 8 Apply 2 coats of TRI-TINTY
- 2 If OLD coating is not a hard, VINYL con potion, but in GOOD condition
  - A Cean thoroughly and let DRY
  - B Count ENTIRE hull with KELIGLO # 101, PARRIER PRIMER
  - Conformal instructions carefully
- $3 A_{FI} = 2$  coats of TRI-TIN. The

#### NEW FIBERGLASS HULLS

- 1. Apply 1 or 2 coats of KEL-GLO # 103 FIBERGLASS PRIMER; follow instructions.
- 2. Apply 2 coats of TRI-TIN.TM

#### NEW ALUMINUM of STEEL HULLS


- 1 Apply KEL-GLO # 102 ZINC CHROMATE PRIMER, foliowing instructions carefully.
- 2 Apply 2 cost, of TRI-TIN.TM

#### NEW WOOD HULLS

- 1 Sand to a smooth finish.
- 2 Apply 1 or 2 coats of KEL-GLO # 101 FIBERGLASS PRIMER; follow instructions.
- 3 Apply 2 coats TRI-TIN. TM

NOTE, DRYING TIME RETWEEN COATS A bound of normal conditions

KEL-GLO CORPORATION, 54 NORTHEAST 73RD STREET, MIAMI, FLORIDA 33138



# assault.

LONG LASTING NON-SELECTIVE WEED KILLER ACTIVE INGREDIENT:

Monuron [3-(p-chiorophenyl)-1.1-dimethylurea] trichloroacetate INERT INGREDIENTS:

12.47% 87.53%

100.00%

ASSAULT weed killer may be used for the control of most annual and perennial grasses, broad-leaved weeds, and general weed control on non-crop land such as: right of ways, industrial plants, lumber yards, tank farms, schools, storage areas, parking lots and similar areas. Best results will be obtained if ASSAULT is applied shortly before or at the time plant growth begins. Thorough watering after application will cause the material to move to the root zone and thus speed up control. The effectiveness and length of time that control will last depend on type of soil, amount of rain and other conditions.

# CAUTION: Keep Out Of Reach Of Children.

Harmforst swar bard directions to skin and eyes. Do not get on skin, in eyes of noticent in one of contact strong dately wash skin with a bit of water, for eyes fush with water for at least 15 mounts and set medical attention.

west

Contains 1.03 pounds of cotax ingred of per callon. Do not use or store hear heat or open flame.

on the most actioned to pro-

#### WEST CHEMICAL PRODUCTS, INC.

42 16 A E 5 C STROUTE CONTROL ON THE CONTROL OF THE CONTROL ON THE

#### DIRECTIONS FOR USE:

For treating one acre (43,560 square ft.), use 100 gallons of solution made by taking 20 gallons of ASSAULT and adding approximately 80 gallons of water. Apply uniformly throughout area. In case of exceptionally heavy vegetation, a second application may be required after 10 days. For small areas, dilute six (6 conces of ASSAULT in one 1; quart of water to treat 100 square feet.

ASSAULT should be used only in equipment recommended by your West Representative. Use only a coarse wet spray to reduce drift. For best results, excess growth hay be moved and removed before applying ASSAULT.

NOTE: Do not apply, allow to drift or drain or flush onto or near croplands. Jawie trees, or other desirable plants, or to areas where their roots may extend. Do not contaminate domestic water supplies or irrigation of thes. Do not allow contact with fertilizers, insecticides, fungicides or seeds. Application equipment should be thoroughly cleaned after use with a detergent and water to prevent corrosion. Equipment which has been used to apply this product should not be used for applying insecticides, fungicides, fertilizers or other agricultural cremics—as plant injury may result.

844

Total Files 40, 52 166

4 - 1 + 1 + 1 + 1 to 1