

The L.A. Lux Company
2285 Scranton Road
Cleveland, OH 44113

NOTE: Changes in labeling, formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.


Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Add the phrase "EPA Registration No. 35969-10" to your label before you release the product for shipment.
3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.


John H. Lee
Product Manager (31)
Disinfectants Branch
Registration Division (PS-767C)

Enclosures

ATTACHMENT IS APPLICABLE

BEST AVAILABLE COPY

SIGNATURE OF APPROVING OFFICIAL

DATE

MICROBIOCID 410

ACCEPTED
with COMMENTS
in EPA Letter Dated:

JUN 30 1987

INDUSTRIAL MICROBICID

ACTIVE INGREDIENTS:

Methylene bis(thiocyanate)
2-(Thiocyanomethylthio)benzothiazole

INERT INGREDIENTS

This product weighs 8.26 pounds per gallon.

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
Registered for the pesticide
Registered under EPA Reg. No.
35969-10 5%
5%
90%

**KEEP OUT OF REACH OF CHILDREN
DANGER**

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

DANGER: Corrosive, causes irreversible eye damage and causes skin damage. Harmful or fatal if swallowed or absorbed through the skin. This product is a potential skin sensitizer. Workmen handling this product or treated material should wear impervious gloves, goggles or face shield, and protective clothing. All protective clothing, work shoes or boots, and equipment must be left at the work site at the end of the day. Eating, drinking, or smoking during use of this product is prohibited.

FIRST AID: In case of skin contact, wash promptly and thoroughly with soap and water and finally with glycerin. If product gets in the eyes, flush immediately with copious amounts of clean, cool water for 15 to 30 minutes. Get medical attention immediately. If product is swallowed, call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Do not apply directly to water or wetlands. Permits may be required for discharges containing this pesticide into lakes, streams, ponds, or public water. For guidance, contact the regional office of the Environmental Protection Agency.

PHYSICAL AND CHEMICAL HAZARDS: Do not expose to extreme temperatures.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Do not expose to extreme temperatures. Do not stack more than five drums high. Drums should be opened in well-ventilated areas. Leaking or damaged drums should be placed in overpack drums for disposal. Spills should be absorbed in sawdust or sand and disposed of in a sanitary landfill. Keep container closed when not in use.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at your EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

BEST AVAILABLE CGPY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner

COOLING TOWERS: MICROBIOCID 410 is used on wood against soft or surface rot and internal or dry rot. It is applied by spraying or brushing on surfaces containing 1.0 to 1.4% MICROBIOCID 410. The amount applied should provide 1.2 to 1.8 lb MICROBIOCID 410 to the recirculating cooling tower. The dosage should provide 2.5 lb of MICROBIOCID 410 per 1000 ft² of wood surface. Soft or surface rot can also be inhibited by spraying MICROBIOCID 410 to the recirculating cooling tower.

The dosage should provide 2.5 lb of MICROBIOCID 410 per 1000 ft² of wood surface. Soft or surface rot can also be inhibited by spraying MICROBIOCID 410 to the recirculating cooling tower. The dosage should provide 2.5 lb of MICROBIOCID 410 per 1000 ft² of wood surface. Soft or surface rot can also be inhibited by spraying MICROBIOCID 410 to the recirculating cooling tower.

COOLING WATER: MICROBIOCID 410 is used to control bacteria and fungi in industrial recirculating cooling water systems. Before use, the system should be cleaned thoroughly to remove old algal growth, microbiological sludge, and debris. The system should then be drained, flushed, refilled with water, and treated with MICROBIOCID 410. The dosage should provide 0.4 to 2.4 fl oz per 1000 gal should be made every 1 to 5 days, depending on the severity of microbiological fouling.

DRILLING FLUIDS: To inhibit bacterial and fungal degradation of drilling of wells, MICROBIOCID 410 is incorporated into drilling fluids at concentrations of 0.1 to 0.5% based on the total wet weight of the fluid.

PETROLEUM SECONDARY RECOVERY: MICROBIOCID 410 is used to control sulfate-reducing bacteria, slime-forming bacteria and fungi in micellar floods, water-disposal systems, and other oil-field water systems. The dosage should provide 7.8 to 26.0 fl oz MICROBIOCID 410 per 1000 gal of water. The dosage should provide 7.8 to 26.0 fl oz MICROBIOCID 410 per 1000 gal of water. The dosage should provide 7.8 to 26.0 fl oz MICROBIOCID 410 per 1000 gal of water.

CRUDE AND REFINED OILS: MICROBIOCID 410 is used to control bacteria and fungi during storage. It should be added to the oil as it is being transferred to storage tanks at the rate of 1.2 to 12.0 fl oz MICROBIOCID 410 per 1000 gal of oil. Addition should be made batchwise where mixing occurs or continuous transfer pump.

Manufactured by:
THE L.A. LUX CO.
2285 SCRANTON RD.
CLEVELAND, OH 44113

EPA REG. NO.

EPA EST.

NET CONTENTS