

ACCEPTED

NOV 24 1980

Under Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. **35980-1**

DIRECTIONS FOR USE
General Classification

It is a violation of Federal Law to use this product in a manner inconsistent with the label.

Shake well before using.



PLANT GROWTH REGULATOR FOR INCREASED YIELDS

ACTIVE INGREDIENT:
Cytokinin (as kinetin) 0.001%*
*based on biological activity
INERT INGREDIENTS: 99.999%

KEEP OUT OF REACH OF CHILDREN
CAUTION

Discard used container in a safe place.

NET CONTENTS: 5 U.S. GALLONS (18.0 Liters)
Derived from: Extract of marine algae.

Distributed by: **ATLANTIC PACIFIC RESEARCH INC.**

P.O. Box 14368
North Palm Beach, Florida 33408.

E.P.A. Reg. No.: 35980-1 Establishment No.: 35980-FL-1

35980-1

DIRECTIONS
(Continued)

For use on **CORN** to increase yields:
Apply CYTEX at the rate of two gallons per acre in 25-100 gallons of water per acre, sprayed over the foliage at the 10' high stage.

For use on **WHEAT** to increase yields:
Apply CYTEX at the rate of one gallon per acre in 25-100 gallons of water per acre, sprayed over the foliage at the anthesis stage.

For use on **CELERY** to increase yields:
Apply CYTEX at the rate of one gallon per acre in 25-100 gallons of water per acre, sprayed over the foliage at the 2-leaf stage.

For use on **GRAPE, PEPPER, STRAWBERRY** to increase yields:
Apply CYTEX at the rate of one gallon per acre in 25-100 gallons of water per acre, sprayed over the foliage at first bloom stage.

For use on **CARROT** to increase yields:
Apply CYTEX at the rate of one gallon per acre in 25-100 gallons of water per acre, sprayed over the foliage at tuber initiation.

For use on **SOYBEAN** to increase yields:
Apply CYTEX at the rate of 1/2 gallon per acre in 10-50 gallons of water per acre, sprayed over the foliage at the R-5 stage.

CYTEX is a plant growth regulator which should be applied with enough water to thoroughly cover all plant surfaces and allow some run-off to the soil.

For use on **ORANGES and APPLES** to increase yields:
Apply CYTEX at the rate of one gallon per acre in 250-500 gallons of water. Apply over the tree foliage at first bloom with ground sprayer equipment.

For use on **TOMATO** to increase yields:
Apply CYTEX at the rate of one gallon per acre in 20-100 gallons of water per acre, sprayed over the foliage at first bloom stage. Aerial application (California only): Apply CYTEX at the rate of one gallon per acre in a minimum of 10 gallons of water per acre.

For use on **PEACH** to increase yields:
Apply CYTEX at the rate of one gallon per acre in 250-500 gallons of water per acre, sprayed over the foliage at pit-hardening stage.

For use on **PEANUT** to increase yields:
Apply CYTEX at the rate of one gallon per acre in 25-100 gallons of water per acre, sprayed over the foliage at pegging stage.

(Continued on side panel)

35980-1

11/24/1980

1/1