	1 A A A A A A A A A A A A A A A A A A A
	CTIONS FOR USE
ACCEPTED DEC 15 1982	General Classification It is a violation of Federal law to use this product in a manner inconsistent with the label. SHAKE WELL BEFORE USING
CYTEX ORGANIC CROP STIMULANT TO INCREASE YOUR HARVEST	Spray leaf surface until thoroughly wet, allow some run-off. Do not apply in direct sunlight. Do not store product at temperatures over 100°F or in direct sunlight. Do not store diluted product for extended periods of time. On indoor plants, remove: from living area and allow adequate ventilation. Profect surfaces
Active Ingredient:	from overspray. Spray plant, allow product to dry and
C ytokinin (as Kinetin) 0.01%* 'based on biological activity	return to living area. CAUTION: Disposal - Do not use empty container. "Wfap empty container and put in trash. Application Rate
Inert Ingredients: 99.99%	General Use - 1 oz. in 1 qt. of water, covers 350 sq. ft For trees and shrubs - 1 oz. in 2 gal. of water, covers 350 sq. ft.
KEEP OUT OF REACH OF CHILDREN CAUTION	With <u>Amway</u> Pistol Grip Sprayer - 1 tbsp. ($\frac{1}{2}$ oz.) and fill container with water (16 oz.) - covers 175 sq. ft With <u>Amway</u> First Prize Sprayer - use 1 oz. and dilute $\overline{\mathbf{y}}$
	with 1 oz. of water, apply to 350 sq. ft. With compressed air sprayer - use 1 oz. and dilute with 32 oz. of water, apply to 350 sq. ft.
NET CONTENTS: 5 FL. OZ. (145 mL)	Initial Application Time
D	<u>·Tomatoes</u> – at first bloom
Distributed by: & Pacific RESEARCH, Inc.	The following for use in all states except California
P.O. Box 14545 North Palm Beach, Florida 33408	<u>Oranges, peppers, strawberries</u> - at first bloom
	<u>Carrots</u> – at 3–5 inch stage
E.P.A. Reg. No. 35980-1 Establishment No. 35980-FL-1	· <u>Celery</u> – at 2–3 leaf stage
	<u>·Peaches</u> - at 40 days after mid-bloom
	• <u>Potatoes</u> - at first bloom or 4-5 weeks prior to harvest
	•Flowering ornamentals and houseplants - apply monthly, following initial application
	· · · ·

2