

35980-1

**DIRECTIONS FOR USE**

**General Classification**

It is a violation of Federal Law to use this product in a manner inconsistent with the label.

Shake well before using.

CYTEX is a plant growth regulator which should be applied with enough water to thoroughly cover all plant surfaces and allow some run-off to the soil. For aerial applications, apply CYTEX with a minimum of 10 gallons of water per acre. Application should not be made in direct sunlight.

**For use on CARROT to increase yields:**

Apply CYTEX at the rate of one gallon per acre in 25-100 gallons of water per acre, sprayed over the foliage at tuber initiation.

**For use on CELERY to increase yields:**

Apply CYTEX at the rate of one gallon per acre in 25-100 gallons of water per acre, sprayed over the foliage at the 2-3 leaf stage.

**For use on CITRUS to increase yields:**

Apply CYTEX at the rate of one gallon per acre in 250-500 gallons of water. Apply over the tree foliage at first bloom with ground sprayer equipment or with nutritional spray.

**For use on PEACH to increase yields:**

Apply CYTEX at the rate of one gallon per acre in 25-100 gallons of water per acre, sprayed over the foliage at pit-hardening stage.

**For use on PEPPER to increase yields:**

Apply CYTEX at the rate of one gallon per acre in 25-100 gallons of water per acre, sprayed over the foliage at first bloom stage. Repeat when branch fruit is in bloom.



(TM)

**PLANT REGULATOR FOR INCREASED YIELDS**

ACTIVE INGREDIENT

Cytokinin (as kinetin) 0.01%\*

\*Based on biological activity

INERT INGREDIENTS: 99.99%

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

Discard used container in a safe place

NET CONTENTS: 5 U.S. GALLONS (189.07 Liters)

Derived from: Extract of marine algae

Distributed by **ATLANTIC & PACIFIC RESEARCH, INC.**

P.O. Box 14846  
North Palm Beach, Florida 33408

EPA Reg. No. 69801 Establishment No. 35980 F11

**ACCEPTED**

**MAR 9 - 1983**

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 35980-1

**DIRECTIONS (Continued)**

For use on PINE to increase conelet retention and seedling growth: Apply Cytex at the rate of 2 gallons in 200 gallons of water per acre, sprayed over the foliage at cone-set

**For use on POTATOES to increase yields:**

Apply CYTEX at the rate of one gallon per acre in 25-100 gallons of water per acre, sprayed over the foliage between tuber set and 3 weeks thereafter or 4-5 weeks before harvest.

**For use on STRAWBERRY to increase yields:**

Apply CYTEX at the rate of one gallon per acre in 25-100 gallons of water per acre, sprayed over the foliage at first bloom stage.

**For use on TOMATO to increase yields:**

Apply CYTEX at the rate of one gallon per acre in 25-100 gallons of water per acre, sprayed over the foliage at first bloom stage.

Since weather, crop, soil and other conditions may vary beyond their control, neither Atlantic & Pacific Research, Inc. nor the Seller make any warranty concerning the effectiveness of the product, including any warranty of merchantability or fitness for a particular purpose. The user assumes all risk and responsibility for use, handling or storage of product not in accordance with directions.

03/09/1983

1/1