

35980-1

12/28/05

BLACK PMS347 5G

ACCEPTED

AGRICULTURE USE REQUIREMENT

Use the product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. These Standards contain requirements for the protection of agricultural workers on farms, forests, nurseries, green houses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to statements on this label about personal protective equipment (PPE) notification to workers and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated area during the restricted entry interval (REI) of 4 hours. Early entry into treated areas involving contact with anything that has been treated is permitted under the worker protection standard only if PPE, coveralls, waterproof gloves, and shoes plus socks are properly worn.

PRECAUTIONARY STATEMENTS

**HAZARD TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear long-sleeve shirt, long pants, waterproof shoes, plus socks. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separated from other laundry.

ENVIRONMENTAL HAZARDS

For terrestrial uses only, do not apply directly to water, areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposing of equipment washwaters or rinseate.

STORAGE AND DISPOSAL

STORAGE: Store away from direct sunlight. Avoid extremes in storage area temperature. Stack no higher than 3 high.
DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Waste resulting from use of this product may be disposed of on site (if allowed by local authorities) or at an approved waste disposal facility. Do not reuse this container. Triple rinse, then offer container for recycling; or puncture and dispose of in a sanitary landfill.



PLANT REGULATOR FOR INCREASED YIELDS

ACTIVE INGREDIENT: Cytokinin (as kinetin) 0.04%*

*based on biological activity
OTHER INGREDIENTS: 99.96%

NET CONTENTS: 5 U.S. GALLONS

Derived from: Extract of marine algae.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Harmful if swallowed or absorbed through the skin! Causes skin irritation! Do not breathe vapor or spray mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly with soap and water after handling.

FIRST AID

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

Call poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency medical information, you can reach a local poison control center at 1-800-222-1222.

(SEE SIDE PANES FOR ADDITIONAL PRECAUTIONARY STATEMENTS)

Manufactured by Atlantic & Pacific
P.O. Box 1336 • Hendersonville, NC 28793
E.P.A. Reg. No.: 35980-1 Establishment No.: 35980-NC-1

DIRECTIONS FOR USE

9/9/05
35980-1
It is a violation of Federal Law to use this product in a manner inconsistent with the label.
Shake well before using.

CYTEX is a plant growth regulator. Apply with enough water to thoroughly cover all plant surfaces and allow some run-off to the soil. For aerial applications, apply CYTEX with a minimum of 10 gallons of water per acre. Do not apply in direct sunlight. For best results DO NOT spray with fungicides.

For use on CARROT to increase yields:

Apply CYTEX at the rate of one gallon per acre in 25-100 gallons of water per acre, sprayed over the foliage at tuber initiation.

For use on CITRUS to increase yields:

Apply CYTEX at the rate of one gallon per acre in 250-500 gallons of water per acre. Apply over the tree foliage at first bloom with ground sprayer equipment or with nutritional spray.

For use on PEACH to increase yields:

Apply CYTEX at the rate of one gallon per acre in 25-100 gallons of water per acre, sprayed over the foliage at pit-hardening state.

For use on PEPPER to increase yields:

Apply CYTEX at the rate of one gallon per acre in 25-100 gallons of water per acre, sprayed over the foliage at first bloom state. Repeat when branch fruit is in bloom.

For use on POTATOES to increase yields:

Apply CYTEX at the rate of one gallon per acre in 25-100 gallons of water per acre, sprayed over the foliage between tuber set and 3 weeks thereafter or 4-5 weeks before harvest.

For use on STRAWBERRY to increase yields:

Apply CYTEX at the rate of one gallon per acre in 25-100 gallons of water per acre, sprayed over the foliage at first bloom stage.

For use on TOMATO to increase yields:

Apply CYTEX at the rate of one gallon per acre in 25-100 gallons of water per acre, sprayed over the foliage at first bloom stage.

Since weather, crop soil and other conditions may vary beyond their control, neither Atlantic & Pacific nor the Seller make any warranty concerning the effectiveness of the product, including any warranty or merchantability or fitness for a particular purpose. The user assumes all risk and responsibility for use, handling or storage of product not in accordance with directions.

BLACK PMS347 5G

DIE LINE BLACK GREEN PMS 347