CHILDREN

Hazards to Humans and Domestic animals. Harmful if swallowed. Avoid contact with skin. Avoid prolonged or repeated breathing of spray mist. Avoid contamination of feed and food stuffs.

CAUTION

#### **Environmental Hazards**

This product is toxic to bees and should not be applied when bees are actively visiting the area. This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

### **Physical or Chemical Hazards**

Do not use J-Chlor-2 with any spray materials which are incompatible with oil. Do not apply within 14 days of sulfurproduce application. Observe the usual precaution concerning application of oil-containing products to plants during high temperatures, since many plants are then more susceptible to oil injury.


## **Directions For Use**


It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

## STORAGE AND DISPOSAL

**STORAGE:** Keep container closed when not in use. Do not use or store near heat or open flame.

**DISPOSAL:** This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not re-use container; crush container and bury in approved landfill. If available, return containers to approved drum reconditioner.





# Buildings and

Contact and Solutions of Flies and Mosquitoes — To reduce opulations of houseflies, stable flies and mosquitoes in farm buildings, in barnyards, in alleys, on dumps and other outdoor locations, dilute 1 pt. with 2½ gallons of water and apply in space by mist blower or directly to flies by hydraulic sprayer, hand sprayer or cylindrical pressure sprayer. Care should be taken to avoid contamination of milk handling equipment, feedstuffs, feed troughs and watering receptacles. Dairy animals should not be present while spraying.

GRAIN STORAGE BINS, ELEVATOR TUNNELS (GALLERY FLOOR, HEADHOUSE), PEANUT WAREHOUSES, FREIGHT CARS, GRAIN TRUCKS, SHIPS HOLDS

Aids in control of cadelle, flat grain beetle, sawtoothed grain beetle, lesser grain borer, red flour beetle, longheaded flour beetle, rice weevil, granary weevil, confused flour beetle, foreign grain beetle and hairy fungus beetle:

Use 1 gallon in 10 gallons of water and apply 2 gallons spray per 1000 sq. ft.; treat when empty. Do not add grain to bin for at least 24 hours or until walls have dried. LOT #\_\_\_\_\_