

Ultraquat 10

MAY 22 1984

36306-10

ACTIVE INGREDIENT:
 Poly(oxyethylene (dimethyliminio) ethylene-
 (dimethyliminio) ethylene dichloride) 10.0 %

INERT INGREDIENT 90.0 %

This product contains 0.85 lb. of active ingredient per gallon and weighs 8.49 lb. per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes, or clothing.

FIRST AID If swallowed, drink promptly a large quantity of water or milk. If eye contact occurs, flush with water for at least 15 minutes. If irritation persists, seek medical attention. Avoid alcohol. Call a physician immediately.

ENVIRONMENTAL HAZARDS This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Permits may be required for discharges containing this pesticide into lakes, streams, ponds, or public water. For guidance, contact the regional office of the Environmental Protection Agency.

BEST DOCUMENT AVAILABLE

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Ultraquat 10 is used to control algae, bacteria, and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove algal growth, microbiological slime, and other deposits. An initial slug addition of 6.13 to 15.32 fluid ounces of Ultraquat 10 per 1000 gallons of water to provide a concentration of 48 to 120 parts per million of Ultraquat 10, based on the total weight of water in the system, is recommended. Repeat initial dosage until control is evident. Subsequent slug additions of 1.53 to 15.32 fluid ounces of Ultraquat 10 per 1000 gallons of water (12 to 120 parts per million of Ultraquat 10) should be employed every 2 to 5 days, or as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the microbiological problem. Slug additions should be made in the sump of water cooling towers.

Ultraquat 10 is used to control bacteria in industrial air-washing systems that maintain effective mist eliminating components. Prior to its use, systems should be cleaned to remove bacterial slime and other deposits. An initial slug dose of 24.7 to 41.2 fluid ounces of Ultraquat 10 per 1000 gallons of water is recommended. Repeat initial dosage until control is evident. Subsequent slug additions of 12.2 to 41.2 fluid ounces of Ultraquat 10 per 1000 gallons of water should be employed each 1 to 5 days, or as needed. The frequency of addition depends upon the relative amount of bleedoff and severity of the bacterial problem. Slug additions may be made to the sump or to the water collection trays of the airwash system.

STORAGE AND DISPOSAL

STORAGE Store in original container in a cool, dry place. Do not store in open containers. Do not store in areas where children or pets can access. Do not store in areas where food or feed is stored. Do not store in areas where food or feed is prepared. Do not store in areas where food or feed is consumed. Do not store in areas where food or feed is stored, prepared, or consumed. Do not store in areas where food or feed is stored, prepared, or consumed.

RESIDUE DISPOSAL Wash containers thoroughly before reuse. Do not reuse for food or feed. Do not reuse for other purposes. Do not reuse for other purposes.

CONTAINER DISPOSAL Do not reuse containers for food or feed. Do not reuse for other purposes. Do not reuse for other purposes.

Manufactured by:

EPA REG. NO. _____ **EPA EST. NO.** _____

NET CONTENTS _____

U.S. ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF PESTICIDE REGISTRATION
REGISTRATION DIVISION
WASHINGTON, D.C. 20460

EPA REGISTRATION NO.

DATE OF ISSUANCE

MAY 23 1984

FORMULA CHANGE

NOTICE OF PESTICIDE:

Under the Federal Insecticide, Fungicide, and Rodenticide Act

NAME OF PESTICIDE SUBJECT

NAME AND ADDRESS OF REGISTRANT (SEE ZIP)

NOTE Changes in labeling formula, differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an endorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.



ATTACHMENT AVAILABLE

SIGNATURE OF APPLICANT (SEE ZIP)

DATE