

EMPIRE
LABORATORIES

EM 0615 WEED KILLER

KILLS MOST VEGETATION

WEEDS CAN BE KILLED QUICKLY

WARNING: KEEP OUT REACH OF CHILDREN.

HARMFUL IF SWALLOWED OR ABSORBED THROUGH SKIN. CAUSES SKIN IRRITATION. SEE SIDE PANEL FOR FURTHER CAUTION STATEMENTS. KEEP AWAY FROM CHILDREN.

ACTIVE INGREDIENTS:

Petroleum oil	96.10%
2,4-Dichlorophenoxyacetic acid, isooctyl ester*	1.09%
Bromacil (5-bromo-3-sec-butyl-6-methyluracil)	0.61%
Pentachlorophenol	0.80%
Other chlorophenols	0.09%

INERT INGREDIENTS: 1.31%

Total 100.00%

* 2,4-Dichlorophenoxyacetic acid equivalent 0.70

CAUTION:

Do not apply liquid directly to, or otherwise permit it to come into contact with, susceptible plants, such as vegetables, flowers, fruit trees, ornamentals, cotton, grapes, or other small fruits. Do not permit it to come in contact with fruit or to them. Do not spray or drift to them. Vapor from this product may injure susceptible plants in their immediate vicinity. Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes, or do not use same spray equipment for other purposes. Do not store near fertilizers, acids, insecticides, or fungicides. Do not contaminate waters used for irrigation or domestic purposes. Do not reuse this container for other materials without thorough cleaning.

CAUTION: Harmful if swallowed or absorbed through skin! Causes skin irritation! Do not breathe vapor or spray mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly after handling. In case of contact wash skin with soap and water; for eyes, immediately flush with large amounts of water for at least 15 minutes and get medical attention. Remove and wash contaminated clothing before re-use.

CAUTION: This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

GENERAL WEED CONTROL: To control most weeds for an extended period of time in non-cropland areas such as parking lots, fence rows, loading ramps, railroad sidings, storage yards, around buildings and industrial plant sites, apply 2 to 5 gals. per 1,000 sq. ft. treated (80 to 200 gals. per acre). To control annual weeds and grasses such as foxtail, ryegrass, wild oats, crabgrass, cheatgrass, broomsedge, ragweed, lambsquarter and turkey mullein, apply 1 to 2 gals. per 1,000 sq. ft. (40 to 80 gals. per acre).

For other HARD TO-KILL perennial weeds and grasses such as Johnson-grass, saltgrass, bounding bet., bracken fern and horsetail apply 3 to 5 gals. per 1,000 sq. ft. (130 to 220 gals. per acre).

To control perennial weeds and grasses, such as smooth brome, Canada bluegrass, purpletop, quackgrass, broomsedge, aster, dandelion, dog fennel, goldenrod, plantain and wild carrot, apply 2 to 4 gals. per 1,000 sq. ft. (80 to 160 gals. per acre).

GENERAL INFORMATION:

A liquid weed killer. To be applied undiluted as a spray on non-cropland areas where bare ground is desired. The degree of weed control and duration of effect will vary with the amount of chemical applied, soil type, rainfall and other conditions.

NOTICE TO BUYER: Seller makes no warranty of any kind, express or implied, concerning the use of this product. Buyer assumes all risks of use or handling whether in accordance with directions or not.

DIRECTIONS: Apply any time of year except when the ground is frozen. Best results are obtained after predominant species are well established in late spring or early summer. Before treating, calibrate equipment to insure uniform cover of the area to be treated. Measure out the proper amount and apply. Do not apply to areas underlaid by roots of desirable trees and shrubs.

EQUIPMENT: Power sprayers, compressed air hand sprayers, knapsack sprayers or an ordinary sprinkler can be used to apply. No dilution or agitation is required.

For more information see Empire's Consultant.

U.S.D.A. REG. NO. 534

EMPIRE LABORATORIES

ATLANTA BALTIMORE CHICAGO MIAMI