

FRUIT AND NUT CROPS Rates given in pints per 100 gallons unless specified. Aircraft use equivalent rate in 10 to 15 gallons water per acre.

APPLES 1 pint red banded leafhopper. Do not apply after second cover spray or within 40 days of harvest. Do not exceed 16 pints per acre.

CITRUS (GRAPEFRUIT, HONEYMELONS, LEMONS, LIMES, ORANGES, TANGELOES, TANGRINES) 3-4 pints grasshopper. Apply to trunk and main branches. Do not apply within 30 days of harvest. Do not feed treated forage to dairy animals or animals being finished for slaughter. Do not use over 10 pints per acre.

CRANBERRIES 3 pints grasshopper, 4 pints armyworm. Do not apply after fruit begins to form.

PEARS 1 pint red banded leafhopper, 1-2 pints pear psylla. Do not apply within 40 days of harvest or after second cover spray. Do not exceed 16 pints per acre.

STRAWBERRIES 2-3 pints per acre, cutworm (climbing only), strawberry crown borer, strawberry weevil. Do not apply after fruit begins to form.

FORAGE, FIELD, SEED AND VEGETABLE CROPS Rates given in pints per acre. Do not exceed recommended rates. Ground rig use 15-100 gallons water per acre. Aircraft 5-50 gallons water per acre.

ALFALFA (Seed crop only) 2 1/2-3 1/2 pints alfalfa looper, alfalfa weevil, armyworm; 1-3 1/2 pints climbing cutworm; 3 pints alfalfa caterpillar, pea aphids, webworm; 2 pints clover leaf weevil, Lygus bug, grasshopper; 1 1/2 pints spittle bug. Do not feed treated forage to dairy animals or animals being finished for slaughter.

BARLEY, OATS, WHEAT 2 pints armyworms, cutworms, grasshopper. Do not feed treated forage to dairy animals or animals being finished for slaughter. No restriction on the use of the grain.

BEANS (Dry, Green, Lima) 4 pints armyworms, Mexican bean beetle, bean leaf beetle, cucumber beetle, blister beetle, flea beetle, cutworm, Lygus bug, bean leaf skeletonizer, garden webworm, pea weevil, 1 pint corn earworm, grasshopper, serpentine leafminer, 1 pint bean leafroller, bean leathopper, 2-6 pints thrips. No limitation on use of shelled beans as human food. Do not apply sprays to green or shelled beans after pods begin to form. Do not feed treated forage to dairy animals or animals being finished for slaughter.

BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER 4 pints cabbage looper, corn earworm, diamondback moth, 3 pints flea beetle, 2 pints green peach aphid, serpentine leafminer, 1 1/2 pints climbing cutworms, fall armyworm, thrips. Do not apply within 30 days of harvest or after edible parts start to form.

CABBAGE 4-6 pints to control insects listed under Broccoli. Do not apply after heads start to form.

CARROTS 3 pints cutworm, flea beetle, 2 pints grasshopper, 4 pints Lygus bugs. No time limitation. Do not feed treated foliage to dairy animals or animals being finished for slaughter.

CELERY 1-5 pints cutworms, fall armyworms, green peach aphid, serpentine leafminer, thrips, 1-5 pints celery leafhopper. Do not apply after celery plants start to bunch or after plants are half mature.

CORN 2-4 pints fall armyworm and corn earworm feeding in whorl, 3 pints chinch bug, cutworm, flea beetle, grasshopper, 2 1/2 pints weevil. Do not feed treated forage to dairy animals or animals being finished for slaughter in fields treated late in season. Do not graze dairy animals. Do graze meat animals on treated stover within four weeks of slaughter.

COTTON 2-4 pints armyworm, boll weevil, bollworm, cotton leafworm, cutworm (climbing), flea beetle, grasshopper, Lygus bug, thrips, 4 pints stink bug, 3 pints flea hopper. Do not graze dairy animals or animals being finished for slaughter in fields treated late in season. Do not graze treated Cotton, or Feed gin waste to dairy animals or animals being finished for slaughter.

EGGPLANT, PEPPERS; 2-3 pints armyworm, blister beetle, Colorado potato beetle, cutworm, flea beetle, leafhopper, serpentine leafminer, 3 pints cucumber beetle. Do not apply within 5 days of harvest.

LETTUCE 4 pints grasshopper, field cricket, 3 pints cucumber beetle, flea beetle, 2 pints armyworm, cutworm, 3-5 pints cabbage looper. Do not apply after seeding stage-leaf lettuce. Do not apply after heads begin to form-head lettuce.

ONIONS 2 pints thrips. Do not apply to green or spring onions.

PARSNIPS 1-2 pints grasshopper, striped flea beetle. Do not use tops for feed or food.

RICE, 2 pints armyworm, chinch bug, cutworm, grasshopper, field stink bug. Do not feed treated forage to dairy animals or animals being finished for slaughter. No limitation on use of grain.

RUTABAGAS 2 pints cutworm, 4 pints flea beetle. Do not use treated tops for feed or food.

SOYBEANS 2-4 pints armyworm, bean leaf beetle, blister beetle, corn earworm, grasshopper, velvetbean caterpillar, 2 pints cricket, 1 1/2 pints flea beetle. Do not feed treated plants or ensilage made from treated plants to poultry, dairy animals or animals being finished for slaughter. Do not feed soybean mill trash to livestock or poultry. Do not apply within 21 days of harvest.

SPINACH 2 pints armyworm, cutworm, green peach aphid, thrips, serpentine leafminer. Do not apply within 21 days of harvest. Do not apply more than once per season.

TOMATOES 2-4 pints armyworm, blister beetle, cabbage looper, Colorado potato beetle, cutworm, flea beetle, cutworm, russet mite, serpentine leafminer, tomato fruitworm, field stink bug, 2 pints thrips, 4 pints grasshoppers. Do not apply within 7 days of harvest.