



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

TRIPLE TIN 4L Fungicide can be applied as a ground or aerial spray to control fungicidal insects. Rates are for general use. The state agricultural extension or agricultural experiment station for specific applications and timing recommendations. With any spray application thorough coverage is essential.

Apply this product only as specified on this label.

PRECAUTIONS:

TRIPLE TIN 4L Fungicide can be applied in combinations with Flowable Sulfur for additional surfactants, spreaders, or stickers. Do not graze dairy or meat animals in treated areas.

GENERAL INSTRUCTIONS FOR APPLICATION

SUGAR BEETS: Cercospora Leafspot

Ground application, apply 3.8 - 9.5 fluid ounces of TRIPLE TIN 4L per acre in at least 15 gallons of water. Aerial application, (helicopter or fixed wing aircraft) apply 3.8 - 9.5 fluid ounces per acre in 5 - 10 gallons of water. Diluted spray should be directed uniformly to all parts of the plant. Use the lower rate for protective sprays and the higher rate for curative sprays. Application should begin when leafspot conditions appear or be repeated at 10 to 14 day intervals. Do not treat within 14 days of harvest. Do not graze or feed.

POTATOES: Early and Late Blight

Ground application, apply 4.7 - 9.5 fluid ounces of TRIPLE TIN 4L per acre in at least 15 gallons of water. Aerial application, apply 4.7 - 9.5 fluid ounces per acre in 5 - 10 gallons of water. Diluted spray should be directed uniformly to all parts of the plant and the application should be repeated at 7 day intervals. Do not apply with the appearance of blight weather conditions and continue on a 7 day schedule. Do not apply by ground or aerial application equipment. The lower rate of application should be used mid to late season when blight infection is in the area.

PEANUTS: Cercospora Leafspot

Ground application, apply 4.7 - 7.6 fluid ounces per acre in 10-100 gallons of water. Aerial application, apply 4.7 - 7.6 fluid ounces per acre in 3 - 10 gallons of water. Use lower rate when leafspot is light, increasing as disease increases.

Apply in sufficient water for uniform, full coverage of foliage. Use lower gallonage in early season spray volume as plant size increases. Application should begin approximately six (6) weeks after leafspot appears. Spray application should continue on a 10-14 day schedule. Do not apply with hogs to feed on peanuts in treated fields. Hulls from treated peanuts may be used in feed for livestock.

CARROTS: Cercospora Leafspot and Alternaria Blight

Apply 3.8 - 7.6 fluid ounces per acre. Use lower rate when diseases are light, increasing rate when diseases are severe. Application should be directed to all parts of the plant. Applications should begin at pre-pollin and continue at 7 day intervals or as needed for control. Do not use tops as feed for livestock.

PECANS: Scab, Brown Leafspot, Downy Spot, Powdery Mildew, Liver S Blotch.

Aerial application, apply 11.4 - 22.8 fluid ounces per acre in a minimum 20 gallons of water. Ground application, apply 11.4 - 22.8 fluid ounces per acre in 15.2 - 45.6 fluid ounces per 500 gallon tank. Diluted spray should be directed to all parts of the tree. Applications should begin at pre-pollin and continue at 7 day intervals or as needed for control. Do not use tops as feed for livestock.

WARRANTY - CONDITION OF SALE:

DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and are not intended to eliminate all risks associated with use of this product. Because such factors as material and manner of use for application are all beyond the control of Mid America Chemical Co. such things as crop injury, ineffectiveness or other unintended consequences may result.

ALL SUCH RISKS ARE ASSUMED BY THE BUYER

MACCO warrants that this product conforms to the chemical description of the label and is ready for use in the directions for use. MACCO makes no other warranties, express or implied, including FITT case shall MACCO or the Seller be liable for consequential, special or indirect damages resulting from the use of this product. The foregoing is a condition of sale by Mid America Chemical Company and is not subject to change.

ACTIVE INGREDIENT:

Triphenyltin hydroxide

INERT INGREDIENTS:

TOTAL

Contains 4 pounds Triphenyltin hydroxide per gallon

KEEP OUT OF REACH OF CHILDREN

DANGER

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. See a physician immediately.
IF SPLASHED IN EYES: Immediately flush eyes with plenty of water and get medical attention.
IF SPILLED ON SKIN: Immediately remove contaminated clothing, including shoes, and wash skin with soap and plenty of water. If irritation develops send for a physician.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

DANGER May be fatal if swallowed or absorbed through the skin. May cause skin irritation. Do not breathe spray mist. May cause irritation of nose, throat and skin. Do not allow contact of spray with skin or eyes, wear goggles or face shield when handling. Wash thoroughly after use and before smoking or eating. Change to clean clothing daily.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal or cleaning of equipment

PESTICIDE DISPOSAL:

Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state or local procedures.

CONTAINER DISPOSAL:

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or disposal of in a sanitary landfill, or by incineration if allowed by state and local authorities.

EPA Reg. No. 36480-76

Net Contents:

EPA Est. No.

Manufactured by:

MID AMERICA CHEMICAL CO., INC.
400 South Fifth Street
Lawrence, Kansas 66044

ACCEPTED

COMMENTS

EPA Letter Dated:

MAR 7 1983

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

36480-76

BEST DOCUMENT AVAILABLE

BEST DOCUMENT AVAILABLE



beans, Potatoes, Carrots And Sugar Beets

ACCEPTED
40.0% COMMENTS
60.0% COMMENTS
TOTAL 100.0%
EPA Letter Dated:

tin hydroxide per gallon

OF CHILDREN

GER

TICAL TREATMENT

or 2 glasses of water and induce vomiting by touching back of
mouth to an unconscious person. See a physician immediately.
After and get medical attention.
ing, including shoes, and wash skin with soap and plenty of

PRECAUTIONARY STATEMENTS.

STATEMENTS

Domestic Animals

in. May cause skin irritation. Do not breathe spray mist. May
of spray with skin or eyes, wear goggles or face shield when
ing. Change to clean clothing daily.

IL HAZARDS

areas may be killed. Do not apply directly to water. Do not apply
contaminate water by cleaning of equipment or disposal of

ID DISPOSAL

or cleaning of equipment

ording to label instructions must be disposed of according to

being, or disposal of in a sanitary landfill, or by incineration if

ts: EPA Est. No.

ity:

MIDAMERICA CHEMICAL CO., INC.
402 South Fifth Street
Lawrenceville, Kansas 66044

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

TRIPLE TIN 4L Fungicide can be applied as a ground or aerial spray to control fungicidal infestations on listed crops. Application rates are for general use. The state agricultural extension or agricultural experiment station specialists should be consulted for specific applications and timing recommendations. With any spray application thorough coverage is essential for good control.

Apply this product only as specified on this label.

PRECAUTIONS:

TRIPLE TIN 4L Fungicide can be applied in combinations with Flowable Sulfur for additional fungicidal control. Do not add surfactants, spreaders, or stickers. Do not graze dairy or meat animals in treated areas.

GENERAL INSTRUCTIONS FOR APPLICATION

SUGAR BEETS: Cercospora Leafspot

Ground application, apply 3.8 - 9.5 fluid ounces of TRIPLE TIN 4L per acre in at least 15 gallons of water. Full coverage of the foliage is necessary for best results. Aerial application, (helicopter or fixed wing aircraft) apply 3.8 - 9.5 fluid ounces of TRIPLE TIN 4L per acre in 5 - 10 gallons of water. Diluted spray should be directed uniformly to all parts of the plant. Use lower gallonage when plants are small and increase volume with plant size. Use the lower rate for protective sprays and the higher rates later in the season or during high infestation periods. Application should begin when leafspot conditions appear or when the disease is in the area and repeated at 10 to 14 day intervals. Do not treat within 14 days of harvest. Do not graze or feed beet tops to livestock.

POTATOES: Early and Late Blight

Ground application, apply 4.7 - 9.5 fluid ounces of TRIPLE TIN 4L per acre in at least 15 gallons of water. Full coverage of the foliage is necessary for best results. A spray pressure of less than 200 pounds is recommended. Aerial application, apply 4.7 - 9.5 fluid ounces per acre in 3 - 10 gallons of water. For helicopter application, fly high enough so as not to whip the vines. Diluted spray should be directed uniformly to all parts of the plant and the gallonage increased according to the size of the plants. Application should begin with the appearance of blight weather conditions and continue on a 7 day schedule. Do not treat within 7 days of harvest when applied by ground or aerial application equipment. The lower rate of application should be used early in the season and the high rate mid to late season when blight infection is in the area.

PEANUTS: Cercospora Leafspot

Ground application, apply 4.7 - 7.6 fluid ounces per acre in 10-100 gallons of water. Aerial application, apply 4.7 - 7.6 fluid ounces per acre in 3 - 10 gallons of water. Use lower rate when leafspot is light, increasing as disease pressure increases.

Apply in sufficient water for uniform, full coverage of foliage. Use lower gallonage in early season when plants are small and increase spray volume as plant size increases. Application should begin approximately six (6) weeks after planting or as soon as first sign of leafspot appear. Spray application should continue on a 10-14 day schedule. Do not apply within 14 days of harvest. Do not allow hogs to feed on peanuts in treated fields. Hulls from treated peanuts may be used in feed for livestock. Do not use vines for feed.

CARROTS: Cercospora Leafspot and Alternaria Blight

Apply 3.8 - 7.6 fluid ounces per acre. Use lower rate when diseases are light, increasing rate when disease pressure increases. Apply in sufficient water for uniform, full coverage of foliage 25 - 100 gallons per acre. Use lower volumes in early season when plants are small and increase spray volume as plant size increases. Applications should begin 6 weeks after planting or as soon as first signs of disease appear. Spray applications should continue at 7 day intervals or as needed for continued control. Do not apply within 14 days of harvest. Do not use tops as feed for livestock.

PECANS: Scab, Brown Leafspot, Downy Spot, Powdery Mildew, Liver Spot, Sooty Mold and Leaf Blotch.

Aerial application, apply 11.4 - 22.8 fluid ounces per acre in a minimum 20 gallons of water. Ground application, mix 3 - 9 fluid ounces of TRIPLE TIN 4L for each 100 gallons of water, 15.2 - 45.6 fluid ounces per 500 gallon tank and apply as a full coverage spray. Diluted spray should be directed to all parts of the tree. Applications should begin at pre-pollination stages when the young leaves are unfolding, and a second application made when the small nuts are forming. Repeat spray applications at 2 - 4 week intervals as needed to maintain control. Use the lower rate for the first two applications or until the disease becomes severe, or during severe scab, powdery mildew, or other disease infections. Do not apply after shucks have started to open.

WARRANTY - CONDITION OF SALE:

DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and should be followed carefully. It is however impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of Mid America Chemical Company or the Seller of this product, such things as crop injury, ineffectiveness or other unintended consequences may result.

ALL SUCH RISKS ARE ASSUMED BY THE BUYER.

MACCO warrants that this product conforms to the chemical description of the label and is reasonably fit for the purposes referred to in the directions for use. MACCO makes no other warranties, express or implied, including FITNESS OR MERCHANTABILITY. In no case shall MACCO or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. The foregoing is a condition of sale by Mid America Chemical Company and is accepted as such by the Buyer.

BEST DOCUMENT AVAILABLE

U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (WH-567) WASHINGTON, D.C. 20460	EPA REGISTRATION NO.	DATE OF ISSUANCE <div style="text-align: center;">MAR 7 1983</div>
NOTICE OF PESTICIDE: <input type="checkbox"/> REGISTRATION <input checked="" type="checkbox"/> REREGISTRATION <i>(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)</i>	TERM OF ISSUANCE	
NAME OF PESTICIDE PRODUCT		
NAME AND ADDRESS OF REGISTRANT (Include ZIP code)		
<div style="text-align: right; font-family: monospace; font-size: 1.2em;"> 16/ 90367 18/ 1 </div>		
<p>NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.</p> <p>On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.</p> <p>A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.</p> <p>Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p>		
<input type="checkbox"/> ATTACHMENT IS APPLICABLE		
SIGNATURE OF APPROVING OFFICIAL		DATE

-2-

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA Sec. 5(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Henry A. Jacoby *H.A.J.*
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (TS-767)

Enclosure