

EPA Reg. No. 36488-25
Label Notification Action
Revised April 7, 2005
Page 1 of 4
[FRONT PANEL- Page 1 of 1]

RINGER® VEGETABLE INSECT ATTACK®

- For Organic Gardening {OPTIONAL LOGO}
- OMRI™ Listed, Organic Materials Review Institute {OPTIONAL LOGO}
- Ready To Use
- Contains *Bacillus thuringiensis* (B.T.)
- Biological insecticide for Control of Tomato Hornworm, Looper, Webworm, Armyworm, Cutworm, Leaf Roller, Tomato Fruitworm, and Other Caterpillars on Vegetables, Fruits, Ornamentals, Flowers and Lawns

ACTIVE INGREDIENTS	BY WT.
<i>Bacillus thuringiensis</i> , subsp. <i>kurstaki</i> solids, spores, and insecticidal toxins	00.4365%
OTHER INGREDIENTS	<u>99.5635%</u>
TOTAL	100.0000%


The percent active ingredient does not indicate product performance and potency rates are not federally standardized.

EPA Reg. No. 36488-25 EPA Est. No. 69208-MO-001

KEEP OUT OF REACH OF CHILDREN

CAUTION(See back panel for additional precautionary statements)

NET CONTENTS 1 LB (0.45 kg)



EPA Reg. No. 36488-25
 Label Notification Action
 Revised April 7, 2005
 Page 2 of 4
 [BACK PANEL- Page 1 of 3]

PRECAUTIONARY STATEMENTS:

Hazards to Humans and Domestic Animals

CAUTION: Harmful if inhaled. Avoid breathing dust. Remove contaminated clothing and wash before reuse. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

FIRST AID	
If Inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.
If on Skin or Clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in Eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for medical treatment. You may also contact 1-800-858-7378 for emergency medical treatment advice.</p>	

ENVIRONMENTAL HAZARDS: Do not apply directly to water. Do not contaminate water when disposing of equipment washwater or rinsate.

EPA Reg. No. 36488-25
 Label Notification Action
 Revised April 7, 2005
 Page 3 of 4
 [BACK PANEL- Page 2 of 3]

GENERAL INFORMATION:

Ringer® Vegetable Insect Attack®, in its easy-to-use shaker canister, is a specially formulated dust to provide best results when caterpillars and worms are newly hatched. The active ingredient, Bacillus thuringiensis is a naturally occurring component of the environment. Ringer® Vegetable Insect Attack® is a highly selective insecticide for use against listed caterpillar pests. Honey bees foraging in treated areas are not harmed when Ringer® Vegetable Insect Attack® is used according to directions.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Application Rates:

Apply Vegetable Insect Attack® when pests are first noticed and repeat every 7 days to 10 days or as necessary. Caterpillar egg hatch can occur over an extended time period. Monitor treated areas for the presence of newly hatched pests and retreat if necessary. Thorough leaf coverage is necessary for effective control. After ingesting a treated portion of the leaf, caterpillars stop feeding within a few hours. Death occurs in a few days.

Apply to foliage of listed crops at a rate of 2 to 3 ounces per 50 feet of row plantings.

Use on these crops:

Broccoli	Peppers
Cauliflower	Lentils
Turnips	Cucumbers
Celery	Squash
Cabbage	Collard Lettuce
Chinese Cabbage	Beets
Kohlrabi	Beans
Tomatoes	Endive
Escarole	Peas
Carrots	Sweet Corn
Melons	Eggplant
Spinach	Potato
Kale	Tobacco
Turnip Greens	
Mustard	
Brussels Sprouts	
Sweet Potato	

To control these insects:

Armyworms
 Diamondback Moth
 Green Cloverworm
 Hornworms
 Imported Cabbageworm
 Loopers
 Melonworm
 Pickleworm
 Mimosa Webworm
 Salt Marsh Caterpillar
 Tomato Fruitworm
 Corn Earworm
 Bollworm
 Tobacco Budworm

EPA Reg. No. 36488-25
Label Notification Action
Revised April 7, 2005
Page 4 of 4
[BACK PANEL- Page 3 of 3]

Use on these fruits:

Blackberries
Dewberries
Strawberries
Blueberries
Grapes
Currants
Raspberries

To Control these insects:

Armyworms
Grape Leafroller/Leafroller
Loopers
Grapeleaf Skeletonizer
Hornworms
Salt Marsh Caterpillar

Use on Lawns

To Control this insect:

Sod Webworm

Use on Flowers and Ornamentals

To Control these insects:

Loopers
Tobacco Budworm
Omnivorous Loopers
Omnivorous Leaf Roller
Diamondback Moth
Armyworms
Ello Moth (Hornworm)
Io Moth
Oleander Moth

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

STORAGE: Store in a cool dry area away from heat or open flame and inaccessible to children.

DISPOSAL:

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Manufactured by:
Woodstream Corporation
69 N. Locust St.
Lititz, PA 17543
(717) 626-2125

www.saferbrand.com

Look for other Safer® Brand products at www.saferbrand.com

Dipel® is a registered trademark of Valent BioSciences.