

U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDE REGISTRATION WASHINGTON, D.C. 20460  <b>NOTICE OF PESTICIDE</b> <small>Registration</small> <small>of Pesticides</small>	EPA REGISTRATION NO. _____ DATE OF REGISTRATION _____
	TERM OF REGISTRATION _____ NAME OF REGISTRANT _____

SEP 19 1984

NAME AND ADDRESS OF REGISTRANT \_\_\_\_\_  
 \_\_\_\_\_  
 \_\_\_\_\_  
 \_\_\_\_\_  
 \_\_\_\_\_

**NOTE:** Changes in address, formula, labeling, or other information that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/Registered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Registration is returned here with \_\_\_\_\_

Registration is in no way to be construed as an endorsement or approval of this product by this Agency. In order to protect health and the environment, the Agency retains the right to suspend or cancel the registration of a pesticide at any time with the Act. It is a violation of any national connection with the registration of a pesticide under this Act to use the name of the registrant or the registration number in any deceptive use of the name of the product has been used by others.

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPLICANT/REGISTRANT \_\_\_\_\_ DATE \_\_\_\_\_

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Timothy A. Gardner  
Product Manager (1)  
Insecticide-Rodenticide Branch  
Registration Division (TS-767)

**Enclosures**

RD:W.E.Nelson:DCR-45294:WANG-0574K:KIM.Raven:479-1013:9/14/84:Del.9/30,84

FRONT  
PRODUCT #165 W.P.

REUTER

SCL WEBWORM ATTACK

ACCEPTED  
DATE: 08/18/84  
BY: [Signature]

SEP 18 1984

(Picture)

[Faint illegible text]  
55435-2

CONTAINS  
BT  
FAST ACTING

- . NATURAL CONTROL
- . EFFECTIVE
- . NO CHEMICALS

Caution: Do not let children play with this product.

ACTIVE INGREDIENT: Bacillus thuringiensis Berliner, var. kurstaki, primary powder  
 fermentation product, potency of 10,000 International Units per milligram of product  
 (equivalent to 7.3 Billion International Units per pound of product).....20%

INERT INGREDIENTS:.....80%

36488-24

CONTAINS  
BT  
FAST ACTING

- . NATURAL CONTROL
- . EFFECTIVE
- . NO CHEMICALS

Caution: Do not let children play with this product.

ACTIVE INGREDIENT: Bacillus thuringiensis Berliner, var. kurstaki, primary powder, fermentation product, potency of 16,000 International Units per milligram of product (equivalent to 7.3 billion International Units per pound of product).....20%

INERT INGREDIENTS:.....80%

EPA Reg. No. 36488-  
EPA Est. No. 36488-VA-1

Net Wt. 4 oz.

NATURAL PEST CONTROL THAT WORKS

REUFER  
SOD WEBWORM ATTACK

Sod Webworm Attack is a natural, effective control for sod webworms. Unlike certain chemical pesticides, this product is environmentally beneficial and has shown no harmful effects on people, pets, or plants. Sod webworms feeding on treated grass quickly become sick, stop feeding and die.

DIRECTIONS FOR USE:

This product is specially formulated for use on lawns as a spray. Application is best with a tank or hose-end sprayer.

Sod webworms emerge in spring and begin feeding on grass blades just above the soil surface. Damage appears as small brown patches. Using a flashlight, they are easily found at night. In mid summer, adult moths lay eggs randomly in the lawn. This creates the most damaging second generation which appears in late summer.

For best results, apply Sod Webworm Attack in the evening when worms begin to feed. Thorough coverage of the grass blades is important to achieve effective control.

Sod Webworm Attack also controls most other worms and caterpillars in the lawn. This product has no effect on earthworms.

APPLICATION RATES:

This package will cover 5,000 sq. ft. of lawn area. Mix two (2) tablespoons per gallon of water. Simply mix the proper amount of wettable powder with a similar amount of water. Mix into a thin paste. Then add the balance of water required. Do not store diluted spray for more than 12 hours. Mix thoroughly and agitate frequently during application.

<u>Tablespoons</u>	<u>Gallons of Water</u>	<u>Sq. Ft. of Coverage</u>
5	1	1,000
10	2	2,000
20	4	4,000
25	5	5,000

One application will effectively control the sod webworm population that is present at the time of application. Second or third generations may appear. This requires a separate application.

STORAGE AND DISPOSAL:

Reclose container and store in a cool, dry place. Discard empty container according to local refuse regulations.

NOTICE: Reuter Laboratories makes no warranty, express or implied, for any particular purpose concerning this material, other than those which are contained on this label.

Sod webworms emerge in spring and begin feeding on grass blades just above the soil surface. Damage appears as small brown patches. Using a flashlight, they are easily found at night. In mid summer, adult moths lay eggs randomly in the lawn. This creates the most damaging second generation which appears in late summer.

For best results, apply Sod Webworm Attack in the evening when worms begin to feed. Thorough coverage of the grass blades is important to achieve effective control.

Sod Webworm Attack also controls most other worms and caterpillars in the lawn. This product has no effect on earthworms.

#### APPLICATION RATES:

This package will cover 5,000 sq. ft. of lawn area. Mix two (2) tablespoons per gallon of water. Simply mix the proper amount of wettable powder with a similar amount of water. Mix into a thin paste. Then add the balance of water required. Do not store diluted spray for more than 12 hours. Mix thoroughly and agitate frequently during application.

<u>Tablespoons</u>	<u>Gallons of Water</u>	<u>Sq. Ft. of Coverage</u>
5	1	1,000
10	2	2,000
20	4	4,000
25	5	5,000

One application will effectively control the sod webworm population that is present at the time of application. Second or third generations may appear. This requires a separate application.

#### STORAGE AND DISPOSAL:

Reclose container and store in a cool, dry place. Discard empty container according to local refuse regulations.

NOTICE: Reuter Laboratories makes no warranty, express or implied, for any particular purpose concerning this material, other than those which are contained on this label.

MANUFACTURED IN USA BY: Reuter Laboratories, Inc.  
14540 John Marshall Highway  
1 (800) 368-2244 Gainesville, VA 22065

|||  
UPC CODE  
79347-00165

NATURAL PEST CONTROL THAT WORKS