

ACCEPTED
JUL 0 2 1987
 Under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 36488-31

FRESH SCENTED (starburst)
 with "CITRUSOAP"

Concentrate
 Side 1
 Revised 6/24/87
 as per letters of
 22 April 1986,
 26 March 1987, and
 12 May 1987

An effective way to kill aphids, mites and mealybugs.

R
 Aphid-Mite Attack

LOGO Reuter

Concentrate

ACTIVE INGREDIENTS: Potassium Salts of Fatty Acid .. 25.00%*

INERT INGREDIENTS:

Citrus Aromatics	0.50%
Other Inerts	74.50%
TOTAL	100.00%

*Includes 18.19% as potassium oleate and 6.81% as potassium salts of other fatty acids:


Keep out of reach of children.

WARNING

(See back panel for additional precautionary statements.)

EPA REG. NO. 36488-GR
 EPA EST. NO. 36488-VA-01

16 fl.oz.
 (473 ml)



Concentrate
Side 2
Rev. 6/24/87

R
Aphid-Mite Attack

Aphid-Mite Attack by Reuter is an effective spray that kills aphids, mites and mealybugs.

Aphid-Mite Attack is a specially formulated combination of fresh scented aromatics and insecticidal soaps. This product kills pests and provides a pleasant, fresh fragrance.

Aphid-Mite Attack can be used on fruits, vegetables, flowers, shade, nut and fruit trees, ornamentals, and indoor and greenhouse plants.

EFFECTIVE AGAINST: All Mites, Aphids, Crickets, Earwigs, Fungus Gnats, Lace Bugs, Leaf-Feeding Caterpillars and Beetles, Leafhoppers, Mealybugs, Plant Bugs, Scales and Scale Crawlers, Thrips, Whiteflies.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Aphid-Mite Attack provides best control when insects are newly hatched. It should be applied at first sighting of insects. Thorough coverage of both the tops and bottoms of leaves is important. Spray leaves to the point of run off. Do not spray foliage in full sun; spray in early morning or in the evening when temperatures are below 85°F.

Apply every 5 to 7 days when insects are present.

APHID-MITE ATTACK MAY BE APPLIED UP TO THE DAY OF HARVEST.

Shake well before use. Use with soft water whenever possible. Keep container tightly sealed to prevent evaporation.

BEST AVAILABLE COPY

Concentrate
Side 3
Rev. 6/24/87

APPLICATION RATE:

Aphid-Mite Attack can be applied with all types of hand or power sprayers. Read and carefully follow the instructions for the type of sprayer you use.

For insects on most plants mix 5 oz. (10 Tbsp.) per gallon of water. On delicate ferns or tender new growth, use at 1/2 rate (5 Tbsp./gal.). Do not use on Nasturtium, Sweetpea, Violet, Mountain Ash, Japanese Maple, Horse Chestnut, or newly planted cuttings.

Do not store diluted spray material. Rinse empty sprayer thoroughly after use.

PRECAUTIONARY STATEMENTS: Hazards to Humans and Domestic Animals

WARNING: Concentrate causes severe eye and skin irritation. Do not get in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT:

If in eyes: flush immediately with plenty of fresh water; get medical attention.

If on skin: flush immediately with plenty of fresh water; get medical attention if irritation persists.

Physical and Chemical Hazards: Do not use or store near open flames.

Environmental Hazards: Keep out of lakes, streams and ponds.

STORAGE AND DISPOSAL

STORAGE: Store securely under lock and key in a cool dry place.
DISPOSAL: Do not reuse empty containers. Rinse original container thoroughly and securely wrap in several layers of newspaper and discard in trash.

NOTICE: Reuter Laboratories makes no warranty, expressed or implied for any purpose other than those contained on this label.

Manufactured in USA by Reuter Laboratories
3450 Natural Way
Manassas Park, VA 22111