

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

May 10, 2022

Georgia Anastasiou Agent Woodstream Corporation c/o Lewis & Harrison 2461 S. Clark St., Suite 710 Arlington, VA 22202

Subject: Notification per PRN 98-10 – Adding Alternate Brand Names

Product Name: Bromethalin 0.01% Pellet III

EPA Registration Number: 36488-71

Application Date: 04/05/2022 Decision Number: 584536

Dear Ms. Anatasiou:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped "Notification" and will be placed in our records.

The following alternate brand names have been added to the product record:

- Victor V® Fast-Kill® Brand Pellets III
- Victor V® Fast-Kill® Brand Bromethalin Pellets III
- Victor V® Fast-Kill® Brand Rodent Killer Pellets III

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Page 2 of 2 EPA Reg. No. 36488-71 Decision No. 584536

If you have any questions, you may contact Jasmin Jackson at 202-566-2797 or by email at Jackson.Jasmin@epa.gov.

Sincerely,

Gene Benbow, Product Manager 7 Invertebrate & Vertebrate Branch 3 Registration Division (7505P) Office of Pesticide Programs MASTER LABEL - OUTER CONTAINER FRONT PANEL (Page 1 of 1)

BROMETHALIN 0.01% PELLET III

Alternate Brand Names:

Victor V® Fast-Kill® Brand Pellets III
Victor V® Fast-Kill® Brand Bromethalin Pellets III
Victor V® Fast-Kill® Brand Rodent Killer Pellets III

This product may only be used inside and within 100 feet of buildings or inside of transport vehicles (ships, trains, or aircraft).

ACTIVE INGREDIENT:	BY WT.
Bromethalin	0.01%
OTHER INGREDIENTS:	<u>99.99%</u>
TOTAL:	100.000%

KEEP OUT OF REACH OF CHILDREN

CAUTION: (See side [back] panel for additional precautionary statements)

NET CONTENTS:

4.0 lb/64.0 oz [to] 50.0 lb/800.0 oz bait pellets

NOTIFICATION

36488-71

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

05/10/2022

MASTER LABEL – OUTER CONTAINER BACK PANEL (Page 1 of 4)

	FIRST AID
HAVE LABEL WHEN OBTAINING TREATMENT ADVICE.	
	- Call a poison control center, doctor or 1-800-858-7378
If swallowed:	immediately for treatment advice.
	- Have person sip a glass of water if able to swallow.
	- Do not induce vomiting unless told to do so by the poison
	control center or doctor.
	- Do not give anything by mouth to an unconscious person.
TREATMENT FOR PET POISONING	
	If an animal eats this bait, call a veterinarian at once.

NOTE TO PHYSICIAN OR VETERINARIAN

Contains the nerve poison Bromethalin. This is not an anticoagulant rodenticide. For humans or animals ingesting bait and/or showing poisoning signs (such as muscle tremors, loss of hind limb use, or seizures for animals), limit absorption by either emesis or gastric lavage. Sublethal symptoms, if present, would be the result of cerebral edema and should be treated accordingly through administration of an osmotic diuretic and corticosteroid.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Keep away from children, domestic animals and pets. All handlers (including applicators) must wear shoes plus socks, and gloves. Any person who retrieves carcasses or unused bait following application of this product must wear gloves.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing them. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish, birds and other wildlife. Dogs and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply this product directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark.

MASTER LABEL – OUTER CONTAINER BACK PANEL (Page 2 of 4)

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL! Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

IMPORTANT: Do not expose children, pets, or nontarget animals to rodenticides. To help prevent accidents:

- 1. Store product not in use out of reach of children and pets.
- 2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six (6) years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears or other potentially destructive animals, or in areas prone to vandalism.
- 3. Dispose of product container and unused, spoiled or unconsumed bait as specified on this label.

Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used wherever children, pets, non-target mammals, or birds may have access to the bait placement location.

USE RESTRICTIONS: This product may only be used to control House Mice (*Mus musculus*), Norway Rats (*Rattus norvegicus*), Roof Rats (*Rattus rattus*), Polynesian Rats (*Rattus exulans*)*, Cotton Rats (*Sigmodon hispidus*)*, Eastern Harvest Mice (*Reithrodontomys humuli*)*, Golden Mice (*Ochrotomys nuttalli*)*, Meadow Voles (*Microtus pennsylvanicus*)*, White-throated Woodrats (*Neotoma albigula*)*, Southern Plains Woodrats (*Neotoma micropus*)*, and Mexican Woodrats (*Neotoma mexicana*)*. This product must be used *in and within* 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port or terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting, beyond 100 feet from a structure, as defined above, is prohibited. This product must not be applied directly to food or feed crops. Burrow baiting with this product is prohibited.

Do not sell this product in individual containers holding less than 4 pounds of bait.

SELECTION OF TREATMENT AREAS: Determine where rats and mice will most likely find an opening and consume the bait. Generally, these areas are along walls, by gnawed openings or rub marks, in or beside burrows, in corners and concealed places, between floors and walls, or in locations

MASTER LABEL – OUTER CONTAINER BACK PANEL (Page 3 of 4)

where signs of rodents have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

Norway Rats, Roof Rats, Polynesian Rats*, Cotton Rats*, White-throated Woodrats*, Southern Plains Woodrats*, and Mexican Woodrats*: Use 1 to 6 ounces of bait per placement. Make placements at intervals of approximately 15 to 30 feet at appropriate locations in infested area. Adjust the amount of bait provided according to the level of rat feeding expected at each placement location.

House Mice, Eastern Harvest Mice*, Golden Mice*, and Meadow Voles*: Use $\frac{1}{4}$ to $\frac{1}{2}$ ounce (7-14 grams) of bait per placement. Make placements at intervals of approximately 8 to 12 feet at appropriate locations in infested area. Up to 3 ounces may be placed at points of extremely high house mouse activity.

Baiting Strategy (Rats and Mice): Highest bait consumption is expected to occur on the first day or two after treatment begins. After several days, inspect the placement and replenish bait at sites where there is evidence of heavy feeding or where there is continued evidence of rodent activity. Normally, the volume of bait consumed will be significantly less with bromethalin baits than with anticoagulant baits. Although one night's feeding will usually provide a lethal dose, maintain bait for at least one week to provide all rodents a chance to feed on the bait. Continue baiting until all signs of feeding have ceased.

Follow-Up: Replace contaminated or spoiled bait immediately. Using gloves, collect and properly dispose of all dead animals and leftover bait. To discourage reinfestation, limit sources of rodent food, water and harborage and use rodent-proof structures as much as possible. If reinfestation does occur, repeat treatment. When a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

*Not for use in California.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

PESTICIDE DISPOSAL: Dispose of wastes resulting from the use of this product in trash or at an approved waste disposal facility.

CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container.

Plastic Containers: Triple rinse (or equivalent) then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or incineration.

Paper Containers: Dispose of empty container by placing in trash, at an approved waste disposal facility or by incineration.

EPA Reg. No. 36488-71 Draft Master Label April 5, 2022 Page 5 of 6

MASTER LABEL – OUTER CONTAINER BACK PANEL (Page 4 of 4)

Note: Seller warrants that this product complies with the specifications expressed in this label. To the extent consistent with applicable law, Seller makes no other warranties, and disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for the intended purpose. To the extent consistent with applicable law, Seller's liability or default, breach or failure under this label shall be limited to the amount of the purchase price. To the extent consistent with applicable law, Seller shall have no liability for consequential damages.

Manufactured For: Woodstream Corporation 69 N. Locust St Lititz, PA 17543 For Information Call: 1-800-800-1819

EPA Reg. No. 36488-71

EPA Est. No. 47629-PA-01

© Copyright 2011 Woodstream Corp. Made In USA

MASTER LABEL – OPTIONAL CLAIMS (Page 1 of 1)

[Optional Marketing/Advertising Claims]

- Palatable [bait] formulation
- Contains Bromethalin
- 0.01% Bromethalin pellets
- Ready to use
- Kills Anticoagulant Resistant Norway Rats, Roof Rats and House Mice
- Norway Rats, Roof Rats, and House Mice usually consume a lethal dose in a single night's feeding, but it may take two or more days from the time of bait consumption for these rodents to die. Rodents cease feeding after consuming a toxic dose. This product kills anticoagulant-resistant Norway Rats, Roof Rats and House Mice
- [Kills] [Controls] House Mice, Norway Rats, Roof Rats, Polynesian Rats*, Cotton Rats*, Eastern Harvest Mice*, Golden Mice*, Meadow Voles*, White-throated Woodrats*, Southern Plains Woodrats*, and Mexican Woodrats*
- May be used in and within 100 feet of buildings
- For use in [Can be used in] Attics, Kitchens, Basements, Garages, consistent with all use restrictions and label precautions and limitations.
- Formulated for House Mouse, Norway Rat, and Roof Rat control
- Demonstrated control of Norway Rats, Roof Rats, and House Mice
- [Controls][Kills] House Mice, Norway Rats, Roof Rats, Polynesian Rats*, Cotton Rats*, Eastern Harvest Mice*, Golden Mice*, Meadow Voles*, White-throated Woodrats*, Southern Plains Woodrats*, and Mexican Woodrats*
- [Controls][Kills] anticoagulant resistant Norway Rats, Roof Rats, and House Mice.
- [Kills][Controls] House Mice, Norway Rats, Roof Rats, Polynesian Rats*, Cotton Rats*, Eastern Harvest Mice*, Golden Mice*, Meadow Voles*, White-throated Woodrats*, Southern Plains Woodrats*, and Mexican Woodrats* in and around (within 100 feet of) the home
- Can be used Indoors or Outdoors (within 100 feet of buildings) for control of House Mice, Norway Rats, Roof Rats, Polynesian Rats*, Cotton Rats*, Eastern Harvest Mice*, Golden Mice*, Meadow Voles*, White-throated Woodrats*, Southern Plains Woodrats*, and Mexican Woodrats*
- For Indoor and Outdoor (within 100 feet of buildings) control of House Mice, Norway Rats, Roof Rats, Polynesian Rats*, Cotton Rats*, Eastern Harvest Mice*, Golden Mice*, Meadow Voles*, White-throated Woodrats*, Southern Plains Woodrats*, and Mexican Woodrats*
- For indoor and outdoor (within 100 feet of buildings) control of House Mice, Norway Rats, Roof Rats, Polynesian Rats*, Cotton Rats*, Eastern Harvest Mice*, Golden Mice*, Meadow Voles*, White-throated Woodrats*, Southern Plains Woodrats*, and Mexican Woodrats*

^{*}Not for use in California.