




36638-1



## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS ( & DOMESTIC ANIMALS )

### CAUTION

#### STATEMENT OF PRACTICAL TREATMENT

Avoid contact with eyes. In case of eye contact immediately flush with water. Get medical attention if irritation persists. Wash hands after handling.

#### ENVIRONMENTAL HAZARDS

Do not contaminate water by cleaning of equipment or disposal of waste.

#### PHYSICAL OR CHEMICAL HAZARDS

Extremely flammable. Contents under pressure. Keep away from fire, sparks and heated surfaces. Do not puncture or incinerate. Exposure to temperature above 130°F. (54°C.) may cause bursting.

#### DIRECTIONS FOR USE

##### GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

##### STORAGE AND DISPOSAL

**Storage:** Store in a cool place until used. Long term storage should be at 72°F. (22°C.) or below. Unused NoMate PBW is flammable and normal precautions to prevent exposing it to sources of ignition should be taken.

**Disposal:** Product that cannot be used ac-

cording to label instructions must be disposed of in compliance with applicable Federal, State or local procedures. Do not reuse container. Wrap empty container and put in trash receptacle.

**Instructions for Use—**NoMate PBW is intended for use in controlling pink bollworm damage in cotton by preventing male moths from finding and mating females. Its use results in reduced crop damage by suppression of female egg laying and larval infestation.

#### Rate of Application


NoMate PBW is used with Bio-Tac® adhesive (adjuvant) alone or with Bio-Tac and an insecticide in a tank mix. Ranges of acceptable application rates and recommended application rates for NoMate are:

|                     | Application<br>Rate Acre | Recommended<br>Rate Acre |
|---------------------|--------------------------|--------------------------|
| NoMate PBW Tank Mix | 5-15 grams               | 7.5 grams                |
| NoMate PBW Alone    | 15-30 grams              | 15.0 grams               |

**Tank Mix—Other than California:** Guidance on acceptable insecticides for use with NoMate PBW and complete tank mix instructions are available from your Albany International representative.

**Tank Mix—California:** Use of the NoMate PBW tank mix in California is regulated under a Section 24(c) Special Local Needs Registration (EPA SLN No. CA 810045). Refer to the Section 24(c) Registration Bulletin for complete tank mix instructions.

# NoMate pbw



For the  
Attract 'N Kill™ System

## For Pink Bollworm Control

A BEHAVIOR MODIFYING CHEMICAL FOR  
PINK BOLLWORM (PECTINOPHORA GOSYPIELLA)

### ACTIVE INGREDIENTS:

(Z,Z)-7,11-Hexadecadien-1-ol-acetate 3.8%

(Z,E)-7,11-Hexadecadien-1-ol-acetate 3.8%

### INERT INGREDIENTS:

92.4%

TOTAL 100.0%

### CAUTION

KEEP OUT OF REACH OF CHILDREN  
SEE SIDE PANEL FOR  
PRECAUTIONARY STATEMENTS

**ALBANY**  
INTERNATIONAL

110 A Street  
Needham Heights, MA 02194

ESTABLISHMENT NO. 36638-MA-1  
EPA REGISTRATION NO. 36638-1

NET WEIGHT 600 grams (1 lb. 5 oz.)

# NoMate<sup>™</sup> pbw



## Time of Application

1. Monitoring to detect the time of pest emergence should begin when linids are planted using "Danka" traps baited with Albany International lures. During the season, monitoring should continue on a frequent basis (every 1-3 days is suggested) to help you schedule NoMate PBW application for most effective control.
2. The first application should be made prior to the time when the number of adult moths present is large enough to produce a damaging larval infestation.
  - A. In many areas, particularly those with a history of larval infestations early in the season, start applications shortly before plants produce the first set of buds (squares).
  - B. If overwintering survival of pink bollworm is low, as indicated by field monitoring and history of infestation, start applications at or just prior to first blooms.
3. Second and subsequent applications should be made before the effect of the previous application significantly diminishes. Do not allow periods between applications of NoMate PBW when mating suppression is no longer effective. Reapply based on the following factors or combinations of them:
  - A. When traps begin to catch moths again, some male moths may be able to find females and another application may be needed. Zero or low trap catches are an indication that the last application of NoMate PBW is working.
  - B. NoMate PBW is designed to last up to 14 days at temperatures of 100°-105°F. Lower temperatures may extend longevity; higher temperatures can reduce longevity. During low temperature conditions, the time between applications should not exceed 21 days.
  - C. Pink bollworm infestations in a field will vary from time to time, usually in cycles from comparatively low to high. When trap catches (and other indicators) show that the population is approaching a high level (peak), it is especially important to be sure that the last application of NoMate PBW is still effective or to make another application.
  - D. Severe weather conditions such as heavy wind and rain can cause "fibers" to be knocked off the plants and thus reduce the longevity of a treatment. Fields should be checked soon after such weather to determine if another application is needed.

## Methods of Field Application

Applications of NoMate PBW by air require specially designed equipment which is available from Albany International. Local custom applicators have this equipment and are prepared to provide application services.

1. With air application equipment, applications can be made by flying with rows or across rows and fiber distribution should be as even as possible. A flying altitude of 15-50 feet is recommended with the higher altitude being flown when ground winds are still or moderate and lower altitudes flown as necessary to compensate for high ground wind conditions.
2. Bio-Tac, an inert sticker, is available from your NoMate PBW supplier and is used to hold individual fibers on cotton foliage. For best results, before mixing with Bio-Tac, keep cans of NoMate PBW sealed and in a cool place (72°-77°F [22°C] or below). See Bio-Tac label for instructions about mixing ratios and other important information. If insecticide is used, it should be added to Bio-Tac, then add NoMate PBW fibers to the mixture. Mix only the amount to be used immediately. The mixture should be applied according to application machinery requirements.
3. Storing partially used cans of NoMate PBW is NOT RECOMMENDED. NoMate PBW is available in both 600 gram and 150 gram cans. When preparing for an application, a combination of these sizes containing the amount needed for the application (or slightly more) should be used.
4. Applications can be made at any time of day.
5. Applications should only be made by a qualified person to insure proper rate and method of application.

## NOTE:

Limiting the use of conventional insecticides may help to maintain the population of predators and parasites to assist in the biological control of other pests in the field.

## Limited Warranty

Albany International warrants that this product, as of the time of its sale by Albany International, conforms to the statements on the label. Except as so warranted, the product is sold AS IS. ALBANY INTERNATIONAL MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED OF FITNESS OR MERCHANTABILITY. In no case shall Albany International be liable for CONSEQUENTIAL or SPECIAL or INDIRECT DAMAGES resulting from the use or handling of this product. This warranty may be varied only by agreement in writing signed by a duly authorized representative of Albany International.