

NOMATE SHOOTGARD

RATE OF APPLICATION

The rate of application should be approximately 37 grams per acre.

TIME OF APPLICATION

One application is needed per season and it should be done just before the moth flight which usually occurs shortly after snow melts and the ground thaws in the area to be treated. It is suggested that those using this product who are not familiar with the biology of the western pine shoot borer seek the guidance of an entomologist or others who can provide advice on the probable timing of moth emergence.

APPLICATION METHOD

Air application is the method of choice and aircraft should be equipped with special application equipment available from Albany International and/or its distributors. Applications should be as uniform as possible. They can be made at any time of day. Nomate Shootgard should be mixed with a suitable grade of Bio-Tap next sticker material to adhere this product to the trees. Bio-Tap is available from Albany International and/or its distributors. See Bio-Tap instructions for use for mixing ratios and other important information. For additional information and/or advice on application contact Albany International.

LIMITED WARRANTY

Albany International Controlled Release Division warrants that the product is as described on the label. Albany International and its distributors do not warrant the product to control the pest. NO OTHER WARRANTY EXPRESSES OR IMPLIED, INCLUDING MERCHANTABILITY, shall be given by Albany International. CONSEQUENT DAMAGE AND LOSS OF PROFITS AND OTHER DAMAGES SHALL BE THE RESPONSIBILITY OF THE USER. THE USER SHALL BE RESPONSIBLE FOR THE PROPER USE OF THE PRODUCT.

NOMATE



SHOOTGARD

WESTERN PINE SHOOT BORER SUPPRESSANT

Active Ingredients	
(E)-9-dodecenyl acetate	2.2%
(Z)-9-dodecenyl acetate	9.1%
Inert Ingredients	88.7%
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE SIDE PANEL FOR PRECAUTIONARY STATEMENTS

Mfg by

ALBANY
INTERNATIONAL
Controlled Release
Division

110 West Street • Fitchburg, MA 01424

ESTD. 1961 • 36638-MA-1
EPA Registration No. 36638-1

NET WEIGHT 1.76 Lb (800 grams)

SHOOTGARD

FLAME SUPPRESSANT



PRECAUTIONARY STATEMENTS

Hazards to Humans (& Domestic Animals)

CAUTION

STATEMENT OF PRACTICAL TREATMENT

Avoid contact with eyes. In case of eye contact immediately flush with water. Get medical attention if irritation persists. Wash hands after handling.

ENVIRONMENTAL HAZARDS

Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Flammable. Keep away from fire, sparks and heated surfaces. Exposure to temperatures above 130° F (54° C) may cause bursting.

INSTRUCTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with labeling.

STORAGE AND DISPOSAL

Keep container in original place until used. Long term storage should be at 50° F (10° C) or below. Unused Nomate Shootgard is flammable and normal precautions to prevent exposing it to fire or heat should be taken.

Dispose of in original container and put in trash receptacle.

Nomate Shootgard is intended for use in protected areas to protect against pine bud scale from forest damage caused by Western pine shoot borer. It is also used against Kauri dieback. This is accomplished by preventing contact between adult weevils and the pine bud scale. The mating process, both egg laying and subsequent fertilization, is thereby reduced.

2.2%
9.1%
80.7%
100.0%

IF CHILDREN

CAUTIONARY

Y
AL
ise

Jan 1984

38-MA
38-1

(800 grams)