

PM 18 36638-20

TPW SPIRAL

NoMate® TPW Spiral is a pheromone communication disruptant for control of tomato pinworm (*Keiferia lycopersicella*).

Active Ingredients:

(E)-4-Tridecen-1-y1 Acetate7.40%

(Z)-4-Tridecen-1-y1 Acetate0.23%

Inert Ingredients92.37%

Total.....100.00%

CAUTION

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

STATEMENT OF PRACTICAL TREATMENT


Avoid prolonged contact with skin. Wash hands after handling.

ENVIRONMENTAL HAZARDS

Keep out of lakes, ponds and streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



EPA Reg. No. 36638-20

EPA Reg. No. 36638 MT-01

Net Contents: 315 grams (equivalent to 0.120 gram active ingredient/spiral)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a cool place until used. Prolonged storage should be at or less than 72°F (22°C).

Pesticide Disposal: After use, used spirals should be retrieved if practical. Discard any retrieved spirals in trash.

Container Disposal: Do not reuse empty pouch. Discard pouch in trash.

INSTRUCTIONS FOR USE

NoMate® TPW Spiral is intended to prevent damage to tomatoes and other crops infested by tomato pinworm. The product functions by disrupting the mating communication between the sexes of the adult moths. By interfering with the natural mating process, fertile egg laying and subsequent larval infestation can be suppressed, thus reducing the need for conventional insecticides. Reduction in conventional insecticide use nurtures the populations of parasites and predators which assist in the biological control of other pests.

RATE OF APPLICATION

The recommended rate of application is 200-400 spirals per acre (500-1,000 spirals per hectare). Use the spirals in a single application per year.

METHOD OF APPLICATION

NoMate® TPW Spiral is applied directly by wrapping the spiral around stems, branches, stakes, wires and strings. Open the container only immediately prior to application, and re-seal any unused spirals. Apply the product at the level of the crop canopy, and, for maximum effectiveness a uniform field distribution is necessary. Take steps to ensure that dispensers do not enter harvest and shipping containers.

SCENTRA
A DIVISION OF ECOGEN INC.

2005 Cabot Blvd. West, Langhorne, PA 19047
215/757-1590 or 800/220-2135

BEST AVAILABLE COPY

monitor, using the wing style traps and Scentry TPW pheromone lures, to detect the time of application. Monitoring should begin when fields are planted. During the season, monitoring should continue to help assess treatment effectiveness. The application of NoMate® TPW Spiral should take place no later than two weeks after transplant (in tomatoes) or if traps capture 2-5 tomato pinworm moths per day. If thorough field monitoring indicates a low level of pinworm infestation, the application may be delayed until economic thresholds are reached.

Ecogen warrants that this product conforms to the description of this label and is reasonably fit for the purposes stated on this label when used in accordance with the directions on this label under normal conditions of use.

ECOGEN MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

If this product is defective, Buyer's exclusive remedy shall be the replacement of the product, or if replacement is impracticable refund of the purchase price. In no case will Ecogen be liable for incidental, consequential or special damages resulting from the handling, storage or use of this product.

BEST AVAILABLE COPY