

day-baldwin

Hexachloride Cream 1% U.S.P.

NET WT. 60 g. (2.11 oz.)

EPA Reg. No. 36988-1
EPA Est No. 36988-NJ-1

Contains Gamma benzene hexachloride 1%, Isopropyl palmitate, stearyl alcohol, cetyl alcohol, sodium lauryl sulfate, glycerin, triethanolamine, water, methyl paraben 0.2%, propyl paraben 0.1% and Cis Isomer of 1-(3 chlorallyl)-3,5,7-triaza-1-azoniaadamantane chloride 0.1%, scented.

FOR EXTERNAL USE ONLY — STORE IN A COOL PLACE

Warning: Keep out reach of children

Usual dose — see package insert.

CAUTION: Federal law prohibits dispensing without prescription.

To remove label, peel off. Lot Number in crimp of tube.

DAY-BALDWIN
Hillside, N.J. 07205

**DO NOT RE-USE CONTAINER.
DESTROY WHEN EMPTY.**

ACCEPTED
OCT 17 1975
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 36988-1

GAMMA BENZENE HEXACHLORIDE CREAM 1% U.S.P.

Description: Gamma benzene hexachloride cream 1% U.S.P. contains gamma benzene hexachloride 1% U.S.P. in a white water washable base made up of isopropyl palmitate, stearyl alcohol, cetyl alcohol, sodium lauryl sulfate, glycerin, triethanolamine, water, methyl paraben 0.2%, propyl paraben 0.1% and cis isomer of 1-(3-chloroallyl)-3, 5, 7-triaza-1-azoniaadamantane chloride 0.1%, scented.

actions: Gamma benzene hexachloride is an ectoparasiticide, toxic to the parasites of scabies and pediculosis.

Indications: For the treatment of infestations with *Sarcoptes scabiei* (scabies), *Pediculus capitis* (head louse) and *Phthirus pubis* (crab louse).

Contraindications: Gamma benzene hexachloride should not be used on persons with a history of hypersensitivity to any of the product components, or those who manifest a primary skin irritation response.

Warnings: Do not apply to face. Avoid getting in eyes. Frequently, repeated applications may cause dermatitis. Not for internal use as this drug produces toxic results similar to those from DDT. Fatal cases of aplastic anemia have resulted from prolonged exposure to vaporized gamma benzene hexachloride.

(OVER)

day baldwin

Cream 1% U.S.P.

NET WT. 60 g. (2.11 oz.)

EPA Reg. No. 36988-1
EPA Est No. 36988-NJ-1

Contains Gamma benzene hexachloride 1%, isopropyl palmitate, stearyl alcohol, cetyl alcohol, sodium lauryl sulfate, glycerin, triethanolamine, water, methyl paraben 0.2%, propyl paraben 0.1% and Cis Isomer of 1-(3 chlorallyl)-3,5,7-triaza-1-azoniaadamantane chloride 0.1%, scented.

FOR EXTERNAL USE ONLY — STORE IN A COOL PLACE

Warning: Keep out reach of children

Usual dose — see package insert.

CAUTION: Federal law prohibits dispensing without prescription.

To remove label, peel off. Lot Number in crimp of tube.

DAY-BALDWIN
Hillside, N.J. 07205

**DO NOT RE-USE CONTAINER.
DESTROY WHEN EMPTY.**

Precautions: In the presence of secondary infections, the use of appropriate treatment should be instituted. Acutely inflamed skin with raw, weeping surfaces should receive appropriate care before applying gamma benzene hexachloride.

Adverse Reactions: Allergic sensitivity reactions and primary skin irritation responses have been reported following applications to the skin.

Dosage and Administration: For Scabies: After a soap and water bath to remove any scaling or crusted detritus, apply a thin layer uniformly to all body skin surfaces from neck to toes (not to face). Leave on for 24 hours and then take another bath to remove the drug. Follow this bath with a complete change of all clothing, bed linen, towels, etc. Retreatment should be avoided unless new clinical lesions are noted, but if required, a second application may be made after one week. **NOTE:** To guard against reinfection, others in the family, or all potential cases, should be treated concurrently. For pediculosis capitis or pubis: Rub the cream into the affected hairy areas (avoid getting in eyes). Leave on for 8 to 12 hours and wash out thoroughly with soap and water. Provide a complete change of head apparel, clothing and bed linens. If necessary, treatment may be repeated after one week. Others in the family, or close contacts, should be treated concurrently to prevent reinfection.

How Supplied: 16 ozs. (one lb.)

60 g (2.11 oz. tube)