

①

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified applicator's certification.



ACTIVE INGREDIENT:
N-[[4-Chlorophenyl] amino] carbonyl]-2, 6-difluorobenzamide 25.0%
INERT INGREDIENTS 75.0%
100.0%
U.S. Patent Numbers 3,748,356, 3,833,903, 3,989,842, 4,013,717, 4,110,469 and other patents pending
A RESEARCH DISCOVERY OF DUPHAR B.V. AMSTERDAM, HOLLAND

Keep Out of Reach of Children

CAUTION

See side/back panels for additional precautionary statements.

EPA Reg. No. 37100-1

EPA Est. No.

NET WEIGHT

POUNDS (kilograms)

RECEIVED duphar

Crop Protection Division, Int.
Apollolaan 161
1077 AR Amsterdam, Holland

JUL 19 1982

REGISTRATION DIVISION
(TS-267)

②

**PRECAUTIONARY
STATEMENTS
HAZARDS TO HUMANS
CAUTION**

Harmful if swallowed. Avoid breathing of dust. Avoid contact with eyes or skin. Avoid contamination of feed and foodstuffs.

**ENVIRONMENTAL
HAZARDS**

Avoid application to lakes, streams, or ponds. Do not apply when weather conditions favor significant or extensive drift from target area. Do not apply on crops.

Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: Dispose of in an incinerator or landfill approved for pesticide containers.

GENERAL: Consult Federal, State or local disposal authorities for approved alternative procedures.

3

**DIRECTIONS FOR USE
RESTRICTED USE PESTICIDE
IT IS A VIOLATION OF FEDERAL
LAW TO USE THIS PRODUCT
IN A MANNER INCONSISTENT
WITH ITS LABELING.**

GYPSY MOTH:

To be used to protect trees and associated plants not used for food or feed. To control larvae and to reduce egg mass deposition in infested areas. For use in forested areas sparsely inhabited by and infrequently used by humans. In campground and other recreational areas, application should be made during periods of minimal use and only after notification of persons using such facilities.

DIMILIN W-25 Insecticide is effective against the larval stages of the gypsy moth. Application should be made prior to full leaf expansion when larvae are in the first, second or third instar.

DIMILIN W-25 applied at label rates does not directly affect bees.

RECEIVED

JUL 19 1992

REGISTRATION DIVISION
(TS-267)

4

AERIAL APPLICATION:

Apply 1 to 4 ounces of DIMILIN W-25 in sufficient water volume (0.5 to 2.5 gals. per acre) to achieve uniform coverage of foliage.

GROUND APPLICATION:

For mist blower equipment, apply the recommended amount in sufficient water volume (1.5 to 10.0 gals. per acre) to achieve uniform coverage of foliage. Do not make more than one application per year.

QUARANTINE PROGRAMS:

For use by United States Department of Agriculture, Animal Plant Health and Inspection Service, Plant Protection and Quarantine personnel and State co-operators involved in quarantine programs. For use in eradication of isolated infestation, two applications of 1 to 2 ounces of DIMILIN W-25 per acre seven to fourteen days apart.

For use in quarantine programs involving the movement of nursery stock from infested to non-infested areas. Two applications of 1 to 2 ounces of DIMILIN W-25 per acre seven to fourteen days apart on nursery stock.

ACCEP
11/1/82
E

CO11 1382

11/1/82
11/1/82
11/1/82

2: 7
(5)

Douglas Fir Tussock Moth: To protect foliage from attack by the larval stage, apply 4 ounces of Dimilin® W-25 per acre by aerial or ground application equipment. Apply after egg hatch when insect larvae are in the 1st and 2nd instar stages of development (approximately 5 days after full hatch).

Aerial application: Apply 4 ounces of Dimilin® W-25 in 0.5 to 2 gallons of water per acre. Uniform coverage of the foliage is essential.

Ground application: Apply 4 ounces of Dimilin® W-25 per acre in sufficient water for uniform, full coverage of foliage (10 to 200 gallons per acre).

Larval kill with Dimilin® W-25 is slower than with some other insecticides. In most instances, several days may be needed before full effect can be seen.

Forest Tent Caterpillar: Apply 4 ounces of Dimilin® W-25 per acre in the early instar stages of development and/or before foliage is expanded.

Aerial application: Apply a recommended amount in 0.5 to 2 gallons of water per acre. Uniform coverage of foliage is essential.

Ground application: Apply recommended amount in sufficient water for uniform, full coverage of foliage (10 to 200 gallons per acre).

Larval kill with Dimilin® W-25 is slower than with some other insecticides. In most instances, several days may be needed before full effect can be seen.

To be used to protect trees and associated plants not used for food or feed. To control larvae and to reduce egg mass deposition in infested areas. For use in forested areas sparsely inhabited by and infrequently used by humans. In campground and other recreational areas, application should be made during periods of minimal use and only after notification of persons using such facilities.

Nantucket Pine Tipmoth: Apply 4 ounces of Dimilin® W-25 per acre in the early instar stages of development, preferably at the beginning of the second tipmoth generation at egg hatch.

Aerial application: Apply recommended amount in 0.5 to 2 gallons of water per acre for uniform coverage.

Ground application: Apply recommended amount in sufficient water for uniform, full coverage of foliage (10 to 200 gallons per acre).

For use in Christmas tree plantations, nurseries, and forested areas during periods of minimal use by workers and others, and only after notification of the treatment.