

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified applicator's certification.



Active ingredient
N-[[[4-Chlorophenyl] amino] carbonyl]-2,6-difluorobenzamide 47.37
Inert ingredients 52.63
100.00

U.S. patent numbers 3,740,388; 3,833,883; 3,888,847; 4,013,717; 4,118,488 and other patents pending.
A research discovery of DUPHAR B.V. Amsterdam, Holland.

Keep Out of Reach of Children
CAUTION
(See side/back panels for additional
precautionary statements).

EPA Reg No 37100-17

EPA Est No 37100-

NET WEIGHT

DUPHAR
Crop Protection Division, Inc.
Appelsteun 151
1077 AR Amsterdam, Holland

BEST DOCUMENT AVAILABLE

AC
37100-17
MAR
1974
EPA
3.2.84

6

39100-17

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS CAUTION

Harmful if swallowed. Avoid breathing of dust. Avoid contact with eyes or skin.

Wear one piece overall which have long sleeves and long pants. Avoid contact of body exposed parts. As specified in the USDA/ EPA Guide for Commercial Applicators during mixing, handling or application of DDT in ODC. Remove contaminated clothing and wash before reuse.

Wear a protective respirator properly approved by the Mining, Quarantine and Safety Administration (formerly the U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health while mixing, loading or making ground application of DDT in ODC.

Avoid consumption of food and beverages.

ENVIRONMENTAL HAZARDS

Avoid application to lakes, streams, or ponds. Do not apply the product in the Gulf States within the coastal zone. This zone includes: 1. The area between the coastline and a parallel line 3 miles from the coastline including dry bays and estuaries which are contiguous to the sea; 2. The area within 3 miles of tidally influenced waters including coastal saline marshes and fringes.

Do not apply when weather conditions favor drift from target area. Do not apply DDT in ODC near any person or dwelling. It is recommended that the product be applied at least 100 feet from dwellings when they are dispersed. If dwellings are not far enough for marking the spray swath or pattern, workers protected at all times should be substituted with the spray swath. Do not contaminate water by leaking of equipment or disposal of wastes. Do not spray or treat treated water bodies.

STORAGE AND DISPOSAL

Preventions: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Pesticide disposal: Pesticide spray mixture or residue that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Hazardous Conservation and Recovery Act.

Container disposal: Dispose of in an incinerator or landfill approved for pesticide containers.

General: Consult Federal, State or local disposal authorities for approved alternative procedures.

DIRECTIONS FOR USE

RESTRICTED USE PESTICIDE
IT IS A VIOLATION OF FEDERAL
LAW TO USE THIS PRODUCT
IN A MANNER INCONSISTENT
WITH ITS LABELING

BEST DOCUMENT AVAILABLE

37100-17

3

COTTON: To control boll weevil through reproduction suppression, do not apply more than 2 oz. (1/8 pint) of DIMILIN ODC per acre per treatment in combination with 2 to 4 quarts of paraffinic crop oil such as DIMOIL™ or a vegetable oil such as cottonseed oil.

MIXING:

1. Fill tank with oil required.
2. Begin agitation and add required amount of Dimilin.
3. Continue agitation while mixing and applying.

For aerial application apply 2 to 4 qts. total volume per acre. For ground application apply 2 to 4 qts. per acre to give uniform coverage.

For maximum control of weevil reproduction, make first application at pinhead square stage of cotton growth when overwintering weevils are entering the fields. Repeat treatments should allow at least 5 to 7 day intervals between applications. Do not make more than six applications and do not apply more than 12 oz. (3/4 pint) of DIMILIN ODC per acre during one growing season. Do not apply after bolls begin to open. Do not rotate crops other than cotton until six (6) months following last application of DIMILIN.

DIMILIN does not kill the adult weevils, however, eggs deposited by treated female weevils will not hatch, thus limiting reproduction. The control of egg hatch and larval development within the square prevents shedding and will also control leaf-tying. After initial control, 7-10 days are required before weevils eggs are laid.

In use for boll weevil control, DIMILIN allows normal build-up of beneficial insects which may aid in control of bollworm and other insects. Emulsifiable concentrate formulations used in tank mixes, in the presence of oil, may result in phytotoxicity. Care should be taken where such mixtures are used.

IMPORTANT NOTICE - Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose express or implied extends to the use of this product, contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

BEST DOCUMENT AVAILABLE