

# STRIKE BAC GENERAL PURPOSE AQUEOUS INSECTICIDE

A concentrated pyrenone micro Emulsion - clear, stable, leaves no oily residues.

Effective, fast acting Insecticide.

For use in Homes, Institutions, Restaurants, Food Processing Plants, Dairies, Livestock and Outdoors.

For Mosquito control apply with conventional mechanical or compressed air equipment.

**ACTIVE INGREDIENTS:**

Pyrethrins	0.1%
Piperonyl Butoxide, Technical	1.0%
Petroleum Distillate	0.4%
<b>INERT INGREDIENTS</b>	98.5%
	100.00%

Equivalent to 0.8% (butyl-carbityl (6-propylpiperonyl) ether and 0.2% related compounds

PYRENONE - Fairfield American Trademark

**CAUTION**

**KEEP OUT OF REACH OF CHILDREN**

See side panel for additional cautions

NET CONTENTS: 1 Gallon, 3.785 Liters

ACCEPTED

OCT 1 1981

37265-15

EPA Reg. No. 37265-15  
General Chemical Inc.  
3502 W. Carnegie Dr.  
Santa Ana, CA 92704

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush skin and eyes with plenty of water. Obtain medical attention if irritation persists.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in food processing plants, wash all equipment, benches, shelving, etc., where exposed food will be handled, with potable water. In home all food processing surfaces and utensils should be covered during treatment, or thoroughly washed before use.

### ENVIRONMENTAL HAZARDS

Remove pets, birds, and cover fish aquariums before spraying. This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as directed. Do not reuse empty containers. Destroy by perforating or crushing and burying in a safe place.

### DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

### DISPOSAL STATEMENTS FOR PESTICIDES

**PESTICIDE DISPOSAL (all except household)**  
Pesticide spray mixture or residue that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

#### CONTAINER DISPOSAL

1. Metal (all sizes)  
Triple rinse (or equivalent) and offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by other approved State and local procedures.

2. Plastic (all)  
Triple rinse (or equivalent) and offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incineration if permitted by State and local authorities.

### DIRECTIONS FOR USE (Indoor Application)

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. The use of this product in food processing establishments should be confined to times periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Vacate treated areas and ventilate before reoccupying.

For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

**SURFACE SPRAYING:** The control roaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, sawtoothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, meal worms, grain mites and cadellies. Use a good sprayer adjusted to coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control carpet beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets, and infested areas of shelving. Repeat treatment as needed.

To control bedbug: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, mouldings and floors. Repeat treatment as needed.

To control fleas and brown dog ticks: Thoroughly spray infested areas, pet beds resting quarters, nearby cracks and crevices, along and behind baseboards, mouldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

To control clothes moths: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and others storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

**SPACE SPRAYING:** Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of flies, houseflies, fruit flies, gnats, mosquitoes, skipper flies, weeps, house flies, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths. Direct spray at an ounces per 100 cubic feet of space.

**FOGGING:** To kill exposed or accessible stages of the above named crawling insects, use a rate of one gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly whenever possible.

**ON ANIMALS:** To protect cattle and horses from horn flies, house flies, mosquitoes and gnats, apply a light mist sufficient to wet the surface of the hair. To control stable flies, horseflies, and deer flies, apply at a rate of 2 ounces per adult cow or horse, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control blood-sucking lice, apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 2 to 3 weeks if required.

**OUTDOOR ADULT MOSQUITO CONTROL:** Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest, around low marshy places, stagnant pools or ponds, and shore lines. Apply at a rate of 1 1/2 gallons per acre (4 1/2 ounces per 1000 square feet).

Using truck-mounted conventional mixing equipment such as Rotomat models SIC, 91 or 100HT traveling at 5 m.p.h. to treat a 300 foot swath calibrate equipment to deliver 4.5 gallons per minute.

### WARRANTY STATEMENT

This is a quality tested product. Any claim is limited to the purchase price of the product. No other warranty is either given or implied.

PTED

1981

37265-15