

WARNINGS: If swallowed, wheeled or absorbed through the skin. Avoid breathing spray mist. Avoid contact with eyes, skin or clothing. Remove pets, birds, and cover fish aquariums before spraying. In case of contact, use immediately with soap and water. Wash arms, hands, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse. Keep all unprotected persons out of operational area while applying the insecticide or working in treated areas during posted period. Avoid contamination of food and foodstuffs. **WARNINGS:** Do not use in nurseries or rooms where infants, or aged persons are confined. **DO NOT** install directly over or within 12 feet of any food handling or dispensing area. Large drafts of forced ventilation may reduce or eliminate the insecticide efficacy of these treatments.

STATEMENT OF PRACTICAL TREATMENT
If Swallowed: Do not induce vomiting because of aspiration hazard. Contact a physician immediately.
If in Eyes: Flush with plenty of water. Get medical attention if irritation persists.
If on Skin: Wash with plenty of water. Get medical attention if irritation persists.

STORAGE AND DISPOSAL
Storage: Store in cool dry place. Keep container closed.
Disposal: Do not reuse empty container. Wrap container
 and put in trash collection.



is especially designed to kill small flying, biting, annoying insects and to act as a repellent as well.

Used in the Micro-Blast[®] Automatic Dispenser the special wave is actuated automatically, emitting 27 mg doses each 15 minutes. Each dose contains millions of tiny particles which float in the air, killing a high proportion of flies, mosquitoes, gnats, moths, and certain other small flying insects. Usually, the effectiveness of Micro-Blast[®] Bug Killer will be noticeable within one hour after the dispenser is actuated.

If less than 24 hour control is required, the dispenser should be set to start one hour before control is desired and run one hour after control is established. Factors such as air change rate, size, and shape of space treated, humidity and temperature affect insect activity and control. Therefore, one unit per 3,000 cubic feet should be considered as an average need with adjustments made depending on specific conditions at place of usage. For best results, mount the dispenser as high as possible and in a location where moving air will carry the spray throughout the desired area. This 2.7 oz. can of Micro-Mist® Bug Killer contains approximately 2,800 "sprays" sufficient to sustain operation for a total 30 days on an around the clock basis and 60 days on a 12 hour basis.

To achieve control on a 24 hour basis it is suggested that this container be changed once a month (once every 2 months on a 12 hour basis).

Cynthia A. Lewis
Maden Enterprises, Inc.
P.O. Box 4599
Columbia, SC 29240

Dear Mr. Lewis:

Subject: Q-Mist Bug Killer II, Micro-Mist Bug Killer
EPA Registration No. 37299-5
Your submission of July 1, 1983

Your pre-approved final printed labelings are acceptable. We have updated
our files to reflect such, and a stamped copy is enclosed for your records.

Sincerely yours,

Timothy A. Gardner
Product Manager (17)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure

TS-767:nelson:DCR-10945:WANG-U216C:bje:Raven:479-2013:7/28/83

CONCURRENCES							
SYMBOL							
SURNAME							
DATE							