

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS**

CAUTION

Harmful if inhaled or absorbed through skin. Avoid breathing dust. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before re-use.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from treated areas.

PHYSICAL AND CHEMICAL HAZARDS

Sulfur dust suspended in air ignites easily. Keep away from heat, sparks, and flame. Do not smoke while applying this product.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT: Do not re-enter treated crop area prior to 24 hours after application.

GENERAL WORKER PROTECTION STATEMENTS: Do not apply this product in such a manner as to directly, or through drift, expose workers or other persons. The area being treated must be vacated by unprotected persons.

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warning must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. (Indicate specific oral warnings which inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure.) When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written warnings must include the following information: "CAUTION". "Area treated with SULFUR on (date of application)." "Do not enter without appropriate protective clothing for 24 hours." "In case of accidental exposure wash with plenty of soap and water, flush eyes with water for at least 15 minutes, get medical attention."

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

STORAGE: Store in a cool, dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site, or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration; or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

ACCEPTED

APR 21 1988

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under
EPA Reg. No. 37568-2

SUNGRO

COMBO SULFUR

DUSTING & SPRAYING

50 Lbs. Net Weight

ACTIVE INGREDIENTS:

Sulfur 90.0%

INERT INGREDIENTS: 10.0%

100.0%

Sulfur Fineness: 93% Passing No. 325 U.S. Sieve

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENTS OF PRACTICAL TREATMENT

If on skin: Wash with plenty of soap and water. Get medical attention.

If in Eyes: Flush with water for at least 15 minutes and get medical attention.

See side panel for additional precautionary statement.

Manufactured By



LYKES



AGRI SALES

P. O. Box 1685
Eaton Park, Florida 33840

EPA Reg. No. 37568 - 2

EPA Est. No. 37568 - FL -002

**OBSERVE THE FOLLOWING
USE PRECAUTIONS**

Sulfur may cause severe fruit and foliage injury to certain crops. Use only if local use has proved that sulfur does not damage crops in your area. Sulfur may burn foliage when temperature is high. Do not apply if temperature during or within 3 days after application is expected to exceed 90° F.

Do not use within 2 weeks of an oil spray treatment or on citrus within 60 days. When growing crops for processing consult the processor before applying sulfur.

APPLES, PEARS: Scab, Powdery Mildew:
As a dust: Apply 30 to 40 lbs. per acre.

As a spray: Use 6 to 8 lbs. per 100 gals. Reduce Sulfur to 2 lbs. if hot weather threatens. Make first application in Green Tip Stage. Repeat application in Pink Bud Stage.

CITRUS: Rust Mites:

As a dust: Apply 80 lbs. per acre.

As a spray: Use 80 lbs. per acre in sufficient water to cover. Begin application at first sign of damage and repeat as necessary. Rust Mites should be controlled at all times of the year to prevent russetting of fruit and greasy spot on leaves.

PEACH: Brown rot blossom blight, Fruit brown rot, Scab:

As a dust: Apply 30 to 40 lbs. per acre.

As a spray: Use 6 to 8 lbs. per 100 gallons in Pink, cover, and pre-harvest sprays.

BEANS, PEAS: Powdery Mildew, Rust:

As a dust: Apply 6 to 8 lbs. per acre.

As a spray: Use 6 to 8 lbs. per acre in sufficient water to cover. Start application as soon as the first true leaves form. Repeat every 8 to 14 days as necessary.

IMPORTANT: Use higher dosage rates during periods of heavy rainfall or rapid growth. Keep new growth covered with light uniform coating.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.