

MALATHION SPRAY

For Lawns And Gardens

ACTIVE INGREDIENT:

Malathion: 0,0-dimethyl phosphorodithioate of
diethyl mercaptosuccinate 57.0%

INERT INGREDIENTS: 43.0%
TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching finger to back of throat. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF INHALED: Remove victim to fresh air. Apply respiration if indicated.

IF IN EYES: Flush eyes for at least 15 minutes with water. Call a physician immediately

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water.


See Elsewhere On Label For Additional Precautionary Statements

EPA Reg. No. 37686-10

EPA Est. No. 51036-6A-1

Manufactured For
CHEM NUT INC
P.O. Box 3706
Albany, Georgia 31706

NET CONTENTS. _____



Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.

PRECAUTIONARY STATEMENTS
Hazards To Humans And Domestic Animals
CAUTION

May be harmful if swallowed. Do not take internally. Avoid contact with eyes, skin and clothing. Avoid breathing vapors, dust, or spray mist. Wash thoroughly after handling. Wash contaminated clothing before reuse. Flush contaminated eyes with plenty of water and get medical attention if irritation persists. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

NOTE TO PHYSICIAN: Malathion is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.
Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

ENVIRONMENTAL HAZARDS

OUTDOOR USE ON NONAQUATIC SITES.

This pesticide is toxic to fish, aquatic invertebrates, and aquatic life stages to amphibians. Do not apply directly to water or wetlands (i.e., swamps, bogs, marshes, and potholes). Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

INDOOR USE.

Use only with adequate ventilation. After using this product indoors, ventilate thoroughly before occupying enclosed spaces. Do not allow contact with treated surface until sprays have dried.

CHEMIGATION PROHIBITION

Do not apply through any type of irrigation system.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

RE-ENTRY STATEMENT



Do not enter treated areas without protective clothing until sprays have dried.

Protective clothing means, at least, a hat or other suitable head covering, a long sleeved shirt and long legged trousers or a coverall type garment (all of sloely woven fabric covering the body, including the arms and legs), shoes and socks

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Do not apply this product in such a manner as to directly or through drift expose workers or others persons. The area being treated must be vacated by unprotected persons.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: **WARNING!** Area treated with Malathion on (date of application). Do not enter without appropriate protective clothing until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

STORAGE. Store in cool, dry place. In case of spill, remove spillage with an industrial type vacuum cleaner and wash area with household bleach.

PESTICIDE DISPOSAL. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

RECOMMENLATIONS

VEGETABLES.

Beans	Tomatoes	Squash	Onions
Cabbage	Potatoes	Melons	Spinach
Turnips	Cucumbers	Lettuce	Mustard

ATTACKING:

Mexican Bean Beetles	Tomato Russet Mites	Mealy Bugs
Leaf Hoppers	Aphids	Cabbage Loopers
Mites	Lace Bugs	Thrips
Spider Mites	White Fly	Imported Cabbage Worms


Use 1 1/2 to 2 teaspoonfuls to one gallon of water, Spray upper and lower sides of leaves thoroughly. Repeat application at 5 to 7 day intervals, as necessary for control. Do not use on Cucurbits unless plants are dry. Do not apply to green onions. Do not apply to cabbage, spinach, mustard or head lettuce within 7 days of harvest. Do not apply to leaf lettuce within 14 days of harvest. Do not use on any other vegetables named within 3 days of harvest.

FRUITS. For control of Aphids, Red Spider Mites, Mealy Bugs, Woolly Aphids, White Fly, Thrips, Tarnished Plant Bug, Pear Psyllid, Scale Crawlers, Plum Curculio attacking Apples, Pears, Grapes, Peaches and Plums.

Use 1 1/2 to 2 teaspoonfuls per one gallon of water. Repeat applications as necessary. Do not use on Fruits within 3 to 7 days of harvest. Malathion may cause injury to McIntosh and Cortland varieties of Apples and Ribier Grapes. Do not apply to peaches within 7 days of harvest. Do not apply to apples or plums within 3 days of harvest or to pears within 1 day of harvest. Do not apply to grapes within 3 days of harvest. Some varieties of grapes are not tolerant to Malathion emulsion and injury may occur to the foliage.

ORNAMENTALS: For the control of Aphids, Red Spider Mites, Woolly Aphids, White Fly, Thrips, Rose Leafhopper attacking Roses, Camellias, Azaleas and evergreens. Injury to certain ferns including Boston, Maidenhair and Pteris, as well as some species of Crassula.

Use 1 1/2 to 2 teaspoonfuls per one gallon of water.



FLIES AND MOSQUITOES: Spray around House, Barns, Outhouses, Stables, Dog Kennels, Manure Piles and other Breeding Places. Mix one part of Malathion to 20 parts of water. Use one gallon of diluted spray to cover 1000 sq. ft. Do not contaminate milk, milking equipment, feed or drinking water.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. THE MANUFACTURER MAKES NO WARRANTIES, EXPRESSED OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USE, WHICH EXTEND BEYOND THE DESCRIPTION ON THE LABEL. Read all directions carefully.

