

**Ray W. Hawksley Co., Inc!**

ACCEPTED

**MICROBIOCIDES SERIES 301**

**WATER TECHNOLOGY**



220 Cutting Blvd. Richmond, Ca 94804 - (415) 235-5780

**ACTIVE INGREDIENT**

Poly [oxyethylene (dimethyliminio) ethylene-  
(dimethyliminio) ethylene dichloride] . . . . . **6.0** %

**INERT INGREDIENT** . . . . . **94.0** %

This product contains **0.5** lb. of active ingredient per gallon and weighs **8.44** lb. per gallon.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes, or clothing.

**FIRST AID:** If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

**ENVIRONMENTAL HAZARDS:** This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Permits may be required for discharges containing this pesticide into lakes, streams, ponds or public water. For guidance, contact the regional office of the Environmental Protection Agency.

**DIRECTIONS FOR USE  
GENERAL CLASSIFICATION:**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**MICROBIOCIDES SERIES 301** is used to control algae, bacteria, and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove algal growth, microbiological slime, and other deposits.

An initial slug addition of **10.1** to **25.25** fluid ounces of **MICROBIOCIDES SERIES 301** per 1000 gallons of water to provide a concentration of **80** to **200** parts per million of **MICROBIOCIDES SERIES 301**, based on the total weight of water in the system is recommended. Repeat initial dosage until control is evident.

Subsequent slug additions of **2.52** to **25.25** fluid ounces of **MICROBIOCIDES SERIES 301** per 1000 gallons of water (**20** to **200** parts per million of **MICROBIOCIDES SERIES 301**) should be employed every 2 to 5 days or as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the microbiological problem. Slug additions should be made in the sump of water cooling towers.

**STORAGE AND DISPOSAL:** Keep container closed when not in use. Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Do not reuse empty container. Triple rinse (or equivalent) and dispose of container in an incinerator or in an approved landfill, or bury in a safe place. Rinsate that cannot be used or reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Open dumping is prohibited.

ACCEPTED

37785-2

29 DEC 1978

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER  
-- UNDER NO. -- SUBJECT  
TO ATTACHED COMMENTS.

Manufactured by:  
Ray W. Hawksley Co., Inc.  
220 Cutting Blvd.  
Richmond, CA 94804

EPA REG. NO. 37785-2 EPA EST. NO. 37785-CA-01

NET CONTENTS: 55 GALLONS