

Ray W. Hawksley Co., Inc.

MICROBIOCIDES SERIES 302

WATER TECHNOLOGY
220 Cutting Blvd. Richmond, Ca 94804 (415) 235-5780

ACCEPTED
37785-3
MAY 9, 1979
UNDER THE FEDERAL INSECTICIDE,
FUNGICIDE AND ROENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
NO.

DIRECTIONS FOR USE.
GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

MICROBIOCIDES SERIES 302: is used to control algae, bacteria, and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove algal growth, microbiological slime, and other deposits.

An initial slug addition of 5.0 to 12.5 fluid ounces of **MICROBIOCIDES SERIES 302** per 1000 gallons of water to provide a concentration of 40 to 100 parts per million of **MICROBIOCIDES SERIES 302**, based on the total weight of water in the system is recommended. Repeat initial dosage until control is evident.

Subsequent slug additions of 1.20 to 12.5 fluid ounces of **MICROBIOCIDES SERIES 302** per 1000 gallons of water (10 to 100 parts per million of **MICROBIOCIDES SERIES 302**) should be employed every 2 to 5 days or as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the microbiological problem. Slug additions should be made in the sump of water cooling towers.

ACTIVE INGREDIENT

Poly (oxyethylene (dimethyliminio) ethylene-
(dimethyliminio) ethylene dichloride) 12.0 %

INERT INGREDIENT 88.0 %

This product contains 1.02 lb. of active ingredient per gallon and weighs 8.53 lb. per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes, or clothing.

FIRST AID: If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Permits may be required for discharges containing this pesticide into lakes, streams, ponds or public water. For guidance, contact the regional office of the Environmental Protection Agency.

BEST DOCUMENT AVAILABLE

STORAGE AND DISPOSAL: Keep container closed when not in use. Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Do not reuse empty container. Triple rinse (or equivalent) and dispose of container in an incinerator or in an approved landfill, or bury in a safe place. Rinse that cannot be used or reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Open dumping is prohibited.

Manufactured by:

Ray W. Hawksley Co., Inc.
220 Cutting Blvd.
Richmond, CA 94804

EPA REG. NO. 37785-3 EPA EST. NO. 37785-CA-01

NET CONTENTS: 55 GALLONS

Ray W. Hawksley Co., Inc.

MICROBIOCIDES SERIES 302

WATER TECHNOLOGY
220 Cutting Blvd. Richmond, Ca 94804 - (415) 235-5780

ACCEPTED
37785-3
MAY 9, 1979
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
EPA UNDER NO.

DIRECTIONS FOR USE
GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

MICROBIOCIDES SERIES 302 is used to control algae, bacteria and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove all growth, micro biological slime, and other deposits.

An initial slug addition of 5.0 to 12.5 fluid ounces of MICROBIOCIDES SERIES 302 per 1000 gallons of water provide a concentration of 40 to 100 parts per million. MICROBIOCIDES SERIES 302, based on the total weight of water in the system is recommended. Repeat initial dosage until control is evident.

Subsequent slug additions of 1.20 to 12.5 fluid ounces of MICROBIOCIDES SERIES 302 per 1000 gallons of water (10 to 100 parts per million of MICROBIOCIDES SERIES 302) should be employed every 2 to 5 days or as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the micro biological problem. Slug additions should be made in the sump of water cooling towers.

ACTIVE INGREDIENT

Poly (oxyethylene (dimethyliminio) ethylene (dimethyliminio) ethylene dichloride) 12.0 %

INERT INGREDIENT 88.0 %

This product contains 1.02 lb. of active ingredient per gallon and weighs 8.53 lb. per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes, or clothing.

FIRST AID: If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Permits may be required for discharges containing this pesticide into lakes, streams, ponds or public water. For guidance, contact the regional office of the Environmental Protection Agency.

BEST DOCUMENT AVAILABLE

STORAGE AND DISPOSAL: Keep container closed when not in use. Do not contaminate water, food, or feed by storage, disposal, cleaning of equipment. Do not reuse empty container. Triple rinse (or equivalent) and dispose of container in an incinerator or in an approved landfill, or bury in a safe place. Rinsate that cannot be reused or reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Open dumping is prohibited.

Manufactured by:

Ray W. Hawksley Co., Inc.
220 Cutting Blvd.
Richmond, CA 94804

EPA REG. NO. 37785-3 EPA EST. NO. 37785-CA-

NET CONTENTS: 55 GALLONS

Ray W. Hawksley Co., Inc.

ACCEPTED
37785-3

MAY 9, 1979

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
EPA REGISTRATION NO.

MICROBIOCIDE SERIES 302

WATER TECHNOLOGY 
220 Cutting Blvd., Richmond, Ca 94804 - (415) 239-5780

INGREDIENT

ethylene (dimethyliminio) ethylene-
(nio) ethylene dichloride] 12.0 %

INGREDIENT 88.0 %

It contains 1.02 lb. of active ingredient per gallon and
53 lb. per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. Avoid breathing vapors. Avoid
skin, eyes, or clothing.

If swallowed, drink promptly a large quantity of milk,
gelatin solution, or, if these are not available, drink large
of water. Avoid alcohol. Call a physician immediately.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep
streams, or ponds. Permits may be required for discharges
this pesticide into lakes, streams, ponds or public water. For
contact the regional office of the Environmental Protection

BEST DOCUMENT AVAILABLE

DIRECTIONS FOR USE.
GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner
inconsistent with its labeling.

MICROBIOCIDE SERIES 302 is used to control algae, bacteria,
and fungi in recirculating commercial and industrial water cooling
towers. Prior to its use, systems must be cleaned to remove algal
growth, microbiological slime, and other deposits.

An initial slug addition of 5.0 to 12.5 fluid ounces of
MICROBIOCIDE SERIES 302 per 1000 gallons of water to
provide a concentration of 40 to 100 parts per million of
MICROBIOCIDE SERIES 302, based on the total weight of water
in the system is recommended. Repeat initial dosage until control is
evident.

Subsequent slug additions of 1.20 to 12.5 fluid ounces
of MICROBIOCIDE SERIES 302 per 1000 gallons of water
(10 to 100 parts per million of MICROBIOCIDE
SERIES 302) should be employed every 2 to 5 days or as needed.
The frequency of addition depends upon the relative amount of
bleedoff and the severity of the microbiological problem. Slug additions
should be made in the sump of water cooling towers.

STORAGE AND DISPOSAL: Keep container closed when not in use.
Do not contaminate water, food, or feed by storage, disposal, or
cleaning of equipment. Do not reuse empty container. Triple rinse (or
equivalent) and dispose of container in an incinerator or in an approved
landfill, or bury in a safe place. Rinsate that cannot be used or
reprocessed should be disposed of in a landfill approved for pesticides
or buried in a safe place away from water supplies. Open dumping is
prohibited.

Manufactured by:

Ray W. Hawksley Co., Inc.
220 Cutting Blvd.
Richmond, CA 94804

EPA REG. NO. 37785-3 EPA EST. NO. 37785-CA-01

NET CONTENTS: 55 GALLONS