

Ray W. Hawksley Co., Inc.

MICROBIOCIDES SERIES 304

WATER TECHNOLOGY

220 Cutting Blvd. Richmond, Ca 94804 (415) 231-5780

ACCEPTED
37785-5
27 NOV 1978

UNDER THE FEDERAL INSECTICIDE AND FUNGICIDE ACT OF 1947 FOR ECONOMIC PEST CONTROL UNDER NO. 37785-5 SUBJECT TO ATTACHED COMMENTS

DIRECTIONS FOR USE
GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

MICROBIOCIDES SERIES 304 is used to control algae, bacteria, and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove algal growth, microbiological slime, and other deposits.

An initial slug addition of 1.90 to 4.75 fluid ounces of MICROBIOCIDES SERIES 304 per 1000 gallons of water to provide a concentration of 16 to 40 parts per million of MICROBIOCIDES SERIES 304, based on the total weight of water in the system is recommended. Repeat initial dosage until control is evident.

Subsequent slug additions of 0.47 to 4 fluid ounces of MICROBIOCIDES SERIES 304 per 1000 gallons of water (4 to 40 parts per million of MICROBIOCIDES SERIES 304) should be employed every 2 to 5 days or as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the microbiological problem. Slug additions should be made in the sump of water cooling towers.

ACTIVE INGREDIENT

Poly [oxyethylene (dimethyliminio) ethylene- (dimethyliminio) ethylene dichloride] 30.0 %

INERT INGREDIENT 70.0 %

This product contains 2.67 lb. of active ingredient per gallon and weighs 8.90 lb. per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes, or clothing.

FIRST AID: If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Permits may be required for discharges containing this pesticide into lakes, streams, ponds or public water. For guidance, contact the regional office of the Environmental Protection Agency.

BEST DOCUMENT AVAILABLE

STORAGE AND DISPOSAL: Keep container closed when not in use. Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Do not reuse empty container. Triple rinse (or equivalent) and dispose of container in an incinerator or in an approved landfill, or bury in a safe place. Rinsate that cannot be used or reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Open dumping is prohibited.

Manufactured by:

Ray W. Hawksley Co., Inc.
220 Cutting Blvd.
Richmond, CA 94804

EPA REG. NO. 37785-5 EPA EST. NO. 37785-CA-01

NET CONTENTS: 55 GALLONS

Ray W. Hawksley Co., Inc.

MICROBIOCIDE SERIES 304

WATER TECHNOLOGY 
220 Cutting Blvd. Richmond Ca 94804 - (415) 235-5780

ACCEPTED
37785-5
27 NOV 1978
UNDER THE FEDERAL
FUNGICIDE AND
FOR ECONOMIC PROSEC
ED UNDER NO.
ATTACHED COM-ENIS

DIRECTIONS FOR USE
GENERAL CLASSIFICATION

It is a violation of Federal law to use this product inconsistent with its labeling.

MICROBIOCIDE SERIES 304 is used to control algae and fungi in recirculating commercial and industrial water towers. Prior to its use, systems must be cleaned to remove growth, microbiological slime, and other deposits.

An initial slug addition of 1.90 to 4.75 (fluoride) MICROBIOCIDE SERIES 304 per 1000 gallons provide a concentration of 16 to 40 parts per million MICROBIOCIDE SERIES 304, based on the total weight in the system is recommended. Repeat initial dosage until no more is evident.

Subsequent slug additions of 0.47 to 4.75 of MICROBIOCIDE SERIES 304 per 1000 gallons (4 to 40 parts per million of MICROBIOCIDE SERIES 304) should be employed every 2 to 5 days. The frequency of addition depends upon the relative bleedoff and the severity of the microbiological problem. Slugs should be made in the sump of water cooling towers.

ACTIVE INGREDIENT

Poly (oxyethylene (dimethyliminio) ethylene (dimethyliminio) ethylene dichloride) 30.0 %

INERT INGREDIENT 70.0 %

This product contains 2.67 lb. of active ingredient per gallon and weighs 8.90 lb. per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes, or clothing.

FIRST AID: If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Permits may be required for discharges containing this pesticide into lakes, streams, ponds or public water. For guidance, contact the regional office of the Environmental Protection Agency.

BEST DOCUMENT AVAILABLE

STORAGE AND DISPOSAL: Keep container closed when not in use. Do not contaminate water, food, or feed by storage, spillage, or cleaning of equipment. Do not reuse empty container. Transfer to equivalent) and dispose of container in an incinerator or in a landfill, or bury in a safe place. Rinseate that cannot be reprocessed should be disposed of in a landfill approved for hazardous waste or buried in a safe place away from water supplies. Open flames prohibited.

Manufactured by:

Ray W. Hawksley Co., Inc.
220 Cutting Blvd.
Richmond, CA 94804

EPA REG. NO. 37785-5 EPA EST. NO. 37785-5

NET CONTENTS: 55 GALLONS

Ray W. Hawksley Co., Inc.

ACCEPTED
37785-5
27 NOV 1978

DIRECTIONS FOR USE
GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

MICROBIOCIDAL SERIES 304

UNDER THE FEDERAL
FUNGICIDE AND RO
FOR ECONOMIC P
ED UNDER NO.
ATTACHED COM

MICROBIOCIDAL SERIES 304 is used to control algae, bacteria, and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove algal growth, microbiological slime, and other deposits.

An initial slug addition of 1.90 to 4.75 fluid ounces of **MICROBIOCIDAL SERIES 304** per 1000 gallons of water to provide a concentration of 16 to 40 parts per million of **MICROBIOCIDAL SERIES 304**, based on the total weight of water in the system is recommended. Repeat initial dosage until control is evident.

Subsequent slug additions of 0.47 to 4.75 fluid ounces of **MICROBIOCIDAL SERIES 304** per 1000 gallons of water (4 to 40 parts per million of **MICROBIOCIDAL SERIES 304**) should be employed every 2 to 5 days or as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the microbiological problem. Slug additions should be made in the sump of water cooling towers.

WATER TECHNOLOGY 
Cutting Blvd. Richmond, Ca 94804 (415) 235 5780

INGREDIENTS

(dimethyliminio) ethylene-
(ethylene dichloride) 30.0 %

Water 70.0 %

Contains 2.67 lb. of active ingredient per gallon and
1.00 lb. per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS
DANGERS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. Avoid breathing vapors. Avoid contact with eyes, or clothing.

If swallowed, drink promptly a large quantity of milk, water, or fruit juice, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

ADDITIONAL HAZARDS: This pesticide is toxic to fish. Keep away from streams, or ponds. Permits may be required for discharges of this pesticide into lakes, streams, ponds or public water. For more information, contact the regional office of the Environmental Protection Agency.

BEST DOCUMENT AVAILABLE

STORAGE AND DISPOSAL: Keep container closed when not in use. Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Do not reuse empty container. Triple rinse (or equivalent) and dispose of container in an incinerator or in an approved landfill, or bury in a safe place. Rinsate that cannot be used or reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Open dumping is prohibited.

Manufactured by:

Ray W. Hawksley Co., Inc.
220 Cutting Blvd.
Richmond, CA 94804

EPA REG. NO. 37785-5 EPA EST. NO. 37785-CA-01

NET CONTENTS: 55 GALLONS