Ray W. Hawksley Co., Inc.

MICROBIOCIDE SERIES 303

WATER TECHNOLOGY 12 CONTROL STATE ST

ACTIVE INGREDIENT Poly[oxyethylene (dimethyliminio) ethylene-(dimethyliminio) ethylene dichloride] 20.0 %

INERT INGREDIENT 80.0 #

this product contains 1.74 lb, of active ingredient per gallon and weighs 8.69 1b. per gallon.

ACCEPTED 37785-9

CAUTION

KEEP OUT OF REACH OF CHIUDREN THE FEDERAL DISECTIONE FUNGICIDE AND HODENTICIDE ACT FUR ECUNOMIC POISON REGISTER ED UNDER NO. SURJECT TO ATTACHED COMMENTS.

PRECAUTIONARY STATEMENTS HAZARDS TO HUHANS AND DONESTIC ANIMALS

(AU'10N: Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes, or clothing.

11th AlD: If swallowed, drink promptly a large quantity or milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid Alcohol. Call a physician immediately.

FINTHORMENTAL MAZARDS: This pesticide is toxic to fish. keep out of lakes, streams, or ponds. Permits may be required for discharges containing this pesticide into lake, dreams, ponds or public water. For quidance, control the regional affice of the Environmental Perform spency,

PM 32 REG# 37785-9 DIRECTIONS FOR USE GENERAL CLASSIFICATION:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

MICROBIOCIDE SERIES 303 is used to control algae, bacteria, and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove algal growth, microbiological slame, and other deposits.

An initial slug addition of 7.9 to 7.3 fluid ounces of MICPOBIOCIDE SERIES 303 per 1000 gallons of water to provide a concentration of 24 to 60 parts per million of MICROBIO-CIDE SERIES 303, based on the total weight of water in the system, is recommended. Repeat-initial dosage until control is evident.

Subsequent slug additions of 0.73 to 8.3 fluid ounces of MICROBIOCIDE SERIES 303 per 1000 gallons of water (6 to 60 parts per million of !!ICROBIOCIDE SERIES 303) should be employed every 2 to 5 days or as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the microbiological problem. Slug additions should be made in the sump of water cooling towers.



STORAGE AND DISPOSAL

Keep container closed when not in use. Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Do not reuse empty container. Triple rinse (or equivalent) and dispose of container in an incinerator or in an approved landfill, or bury in a safe place. Rinsate that cannot be used or reprocessed should be disposed of in a landfill approved for pesticides or buriet in a safe place away from water supplies. Open dumping is prohibited.

MANUFACTURED BY:

Ray !! Hawksley Co., Inc. 220 Cutting Blvd. Richmond, Ca. 94804 EPA REG. 110. 37785 CPA EST. 40. 37785- CA-01

HET CONTENTS: 30 Gallons