

Ms. Kim Davis
Agent for Professional Supply, Inc.
RegWest
30856 Rocky Road
Greely, CO 80631-9375

DEC 27 2001

Dear Ms. Davis:

Subject: Amendment - Comply with 2000 Diazinon MOA;
Change Signal Word;
Update Storage and Disposal statements; and
Change PHI's;
Basic Confidential Statement Of Formula
Professional brand Pest Control Formula DC-500
EPA Registration Number 37915-6
Your submission dated August 8, 2001

The amendment referred to above, submitted in connection with registration under FIFRA section 3(c)(7)(a), is acceptable provided that you:

1. Submit and/or cite all data or other material required for registration/reregistration of your product under FIFRA section 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:
 - a. On the front panel change "For Use In or Around the Home Vegetable Garden, Home Flower Garden and Home Lawn" to "For outdoor use on the home lawn and the home garden only".
 - b. Under DIRECTIONS FOR USE change "Reformulation or repackaging this product is prohibited." to "Reformulation or repackaging of this product is prohibited.".
 - c. Under General Information add "Do not apply in greenhouses".
 - d. Under Precautionary Statements replace "May be fatal if swallowed. May be absorbed through the skin." with "Harmful is swallowed, inhaled or absorbed through the skin.". Also change "Do not breathe spray mist." to "Do not breathe vapor or spray mist.".
 - e. Under Environmental Hazards change "For lawns, where irrigation (watering) is recommended..." to "For lawns, where watering is recommended...".
 - f. Under Environmental Hazards change "Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark." to "Do not apply directly to water."

3. Submit three (3) copies of your final printed labeling before you release the product for shipment.
4. Comply with all applicable provisions of the 2000 Memorandum of Agreement Regarding Pesticide Products Containing Diazinon (i.e., the Diazinon MOA). In accordance with the Diazinon MOA you have requested the following revisions to the conditions of registration for this diazinon product with outdoor non-agricultural (i.e., home and garden) uses:
 - a. You have requested under section 6(f) of FIFRA that this registration be voluntarily canceled on a date not later than December 31, 2004.
 - b. You may not manufacture, label, or release for shipment this product after June 30, 2003.
 - c. You may not sell or distribute this product after August 31, 2003.
 - d. The registration for this product shall expire and be canceled on December 31, 2004.
 - e. You will send to every person who directly purchases this product (i.e., to every seller) a notification of Product Recovery Program to be conducted by Syngenta and Makhteshim after December 31, 2004.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

The Basic Confidential Statement of Formula (CSF) dated June 20, 2001 is acceptable and has been added to this product's registration file.

Sincerely,

George T. LaRocca
Product Manager 13
Insecticide Branch
Registration Division (7505C)

3/9

This product must not be used on golf courses and sod farms.

Professional Brand Pest Control Formula DC-500

Lawn and Garden Insect Control

For Use In or Around the Home Vegetable Garden, Home Flower Garden and Home Lawn

For Control of Certain Insects on Fruits, Vegetables, Lawns and Ornamentals

(Do Not Use in the Home)

Active Ingredient:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 48.0%

Other Ingredients* 52.0%

*Contains xylene range aromatic solvent Total 100.0%

Contains 4 lbs. Diazinon per gallon.

ACCEPTED
with COMMENTS
In EPA Letter Dated:

DEC 27 2001

37915-6

Keep Out of Reach of Children

CAUTION

See back panel for additional precautionary statements.

Net Contents: _____ fl. oz.

Professional Supply, Inc.

4403 East Speedway

Tucson, AZ 85712

EPA Reg. No. 37915-6

EPA Est. 37915-AZ-1

{Back Panel}

DIRECTIONS FOR USE and CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire *Directions for Use* and *Conditions of Sale and Warranty* before using this product. If terms are not acceptable, return the unopened product container at once.

Conditions of Sale and Warranty

The *Directions for Use* of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Plant injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of Professional Supply, Inc. or the Seller. All such risks shall be assumed by the Buyer.

Professional Supply, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the *Directions for Use* subject to the inherent risks referred to above. Professional Supply, Inc. makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall Professional Supply, Inc. or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.

Professional Supply, Inc. and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing *Conditions of Sale and Warranty*, which may be varied only by agreement in writing signed by a duly authorized representative of Professional Supply, Inc.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Reformulation or repackaging this product is prohibited.

Failure to follow the *Directions for Use* and precautions on this label may result in poor insect control, plant injury or illegal residues.

General Information

DC-500 is an emulsifiable solution which, when diluted with water according to the directions, is intended for the control of outdoor insects which destroy lawns, fruit trees, vegetable plants, flowers, trees and shrubs or annoy pets and people. Contains diazinon insecticide which provides positive control of many "hard-to-kill" sucking and chewing insects. DC-500 works fast and remains effective. Do not use in barns or dwellings. Do not use on livestock.

In the ***Directions for Use*** which follow, best control is obtained when application is made at first sign of pests. Application should be repeated only as directed to maintain control. To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, since plant injury may result. Not for use on plants being grown for sale or other commercial use, or for research purposes. For use on plants intended for aesthetic purposes or climactic modification and being grown in ornamental gardens or lawns and grounds. Observe days interval between last application and harvest indicated by number in () following crop. Do not apply within indicated interval or illegal residues may occur.

Fruits

Unless otherwise indicated, only five (5) applications per season permitted. To protect bees, do not apply to fruit trees when trees or substantial numbers of weeds in the orchard are in bloom.

Crop	Pest(S)	Rate	Timing/Special Directions
Cherries (21)	Fruit Tree Leafrollers	1 tbsp. in 3 gals. of water	Apply in cover sprays as infestations occur beginning at petal fall. Repeat application after 7 days if necessary. Do not apply more than 2 tbsp. of diazinon per 1000 sq. ft. per application or make more than 4 applications per season.
	Black Cherry Aphids	1 tbsp. in 3 gals. of water	Apply as a cover spray when aphids appear. Repeat application after 7 days if necessary. Do not apply more than 2 tbsp. of diazinon per 1000 sq. ft. per application.
	San Jose Scale Crawlers	1 tbsp. in 3 gals. of water	Apply during the pre-pink stage or when crawlers first appear. Do not apply more than 2 tbsp. of diazinon per 1000 sq. ft. per application.
	Cherry Rust Mites	½ - 1 tbsp. in 3 gals. of water	Apply to trees as a single supplemental spray after harvest. Do not apply more than 2 tbsp. of diazinon per 1000 sq. ft. per application.
Grapes (28)	Pacific Spider Mites, Grape Berry Moths, Grape Leafrollers, Omnivorous Leafrollers, Aphids and Spider Mites	1 - 1½ tbsp. per 1000 sq. ft.	Apply as pests occur as a thorough cover spray. Repeat application after 7 days if necessary. Do not make more than 5 applications per season. Do not apply within 28 days of harvest or illegal residues may occur.
Peaches (21)	Aphids, Brown Mites (Clover Mites), Two-spotted Spider Mites, Olive Scale Crawlers and San Jose Scale Crawlers	1 tbsp. in 3 gals. of water	Apply as infestations occur. Repeat application after 14 days if necessary. Do not apply more than 2 tbsp. of diazinon per 1000 sq. ft. per application or make more than 4 applications per season.
Plums (21)	Brown Mites (Clover Mites), European Red Mites	1 tbsp. in 3 gals. of water	Apply as needed every 7 days up to a maximum of 4 applications. Do not apply more than 2 tbsp. of diazinon per 1000 sq. ft. per application.
	Leafcurl Plum Aphids, Mealy Plum Aphids and Thistle Aphids	½ - 1 tbsp. in 3 gals. of water	See above.

Crop	Pest(S)	Rate	Timing/Special Directions
Strawberries (5)	Aphids and Two-spotted Spider Mites	1 tbsp. in 3 gals. of water	Apply as pests occur. Repeat application if necessary every 7 days, but do not make more than 3 applications per season.
	Cyclamen Mites	2 tbsp. in 3 gals. of water	Direct spray to the plant crowns. Plants should be sufficiently agitated to assure thorough coverage of the foliage. Repeat applications after 7 days if necessary. Application may be made to plants after harvest. Do not make more than 3 foliar applications per season.
	Strawberry Leafrollers	$\frac{3}{4}$ - 1 tbsp. in 3 gals. of water	Apply $4\frac{1}{2}$ gallons of spray per 1000 sq. ft. when blossoms show color.

Vegetables

For foliar or soil applications, DC-500 should always be applied as a spray in sufficient water to assure thorough coverage of the foliage or soil. **Soil Incorporation:** Following application of this product to control vegetable soil insects, immediately incorporate the product into the soil to the recommended depth (See **Timing/Special Directions**). For control of surface cutworms incorporate 2-3 inches; for subterranean cutworms incorporate 3-6 inches. **To Protect Bees:** Foliar application of this product should not be made to peas or beans if these vegetables or weeds in the treatment area are in bloom, or to corn during the pollen shed period.

For foliar applications, observe days interval between last application and harvest indicated by number in () following crop. Do not apply within indicated interval or illegal residues may occur. Observe maximum numbers of applications per season.

Crop	Pest	Rate	Timing/Special Directions
Beans: Lima and Snap [succulent varieties only] (7)	Cutworms	3 - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil.
	Wireworms	$4\frac{1}{2}$ - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
Beets, Red [Table] (14)	Cutworms	3 - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil.
	Mole Crickets	$1\frac{1}{2}$ tbsp. per 1000 sq. ft.	Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	$4\frac{1}{2}$ - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
Cabbage (21)	Root Maggots	3 - $4\frac{1}{2}$ tbsp. per 1000 sq. ft.	Apply to plant beds or fields just before sowing or seed or transplanting and immediately incorporate into the top 3-4 inches of soil.
		$\frac{1}{2}$ - 1 tbsp. in 3 gals. of water	In transplant water as a drench application when $4\frac{1}{2}$ -7 gals. of water are used per 1000 sq. ft. apply $\frac{1}{2}$ tbsp. per plant by hand to direct spray to the base of the plant. Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.
	Cutworms	3 - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil.
	Mole Crickets	$1\frac{1}{2}$ tbsp. per 1000 sq. ft.	Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	$4\frac{1}{2}$ - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids	$\frac{1}{2}$ - 1 tbsp. per 1000 sq. ft.	Apply as insect occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.

Crop	Pest	Rate	Timing/Special Directions
Cantaloupes (3)	Cutworms	3 - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil.
	Wireworms	4½ - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Striped Cucumber Beetles, Melonworms, Leafhoppers and Spider Mites	½ - 1 tbsp. per 1000 sq. ft.	Apply as insects occur. Repeat application, as necessary, ever 7 days. Do not make more than 5 applications per season.
Carrots (14)	Cutworms	3 - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil.
	Mole Crickets	1½ tbsp. per 1000 sq. ft.	Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	4½ - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids	1 tbsp. per 1000 sq. ft.	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.
Cucumbers (7)	Cutworms	3 - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil.
	Wireworms	4½ - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Striped Cucumber Beetles	1 tbsp. per 1000 sq. ft.	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.
Lettuce (14)	Cutworms	3 - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil.
	Mole Crickets	1½ tbsp. per 1000 sq. ft.	Apply to plant beds or fields 1-2 days before sowing or seed or transplanting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	4½ - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
Onions: Green and Bulb (14)	Onion Maggots	3 - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and mix into the top 3-4 inches of soil by disking, harrowing or other suitable means. Note: Diazinon will not control organophosphate-resistant onion maggots.
	Onion Thrips	1 tbsp. per 1000 sq. ft.	Apply as thrips appear. Repeat application, as necessary, every 7 days. Do not make more than 3 applications per season.
	Wireworms	4½ - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into top 4-8 inches of soil.
Peas [succulent only] (7)	Cutworms	3 - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil.
	Wireworms	4½ - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids and Leafminers	¾ - 1 tbsp. per 1000 sq. ft.	Apply as insects occur. Repeat application, as necessary, every 5 days. Do not make more than 3 applications per season. Peas may be picked immediately after application. Notes: 1) Diazinon will not control organophosphate-resistant leafminers; 2) to protect bees, do not apply foliarly if the crop or weeds in the treatment area are in bloom.
Peppers (5)	Aphids and Serpentine Leafminers	½ - 1 tbsp. per 1000 sq. ft.	Apply as insects appear. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.

Crop	Pest	Rate	Timing/Special Directions
Potatoes (35)	Cutworms (surface and subterranean)	3 - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil.
	Wireworms	4½ - 6 tbsp. per 1000 sq. ft.	Broadcast before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Colorado Potato Beetles and Dipterous Leafminers	½ - ¾ tbsp. per 1000 sq. ft.	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Note: Diazinon will not control organophosphate-resistant leafminers.
	Banded Cucumber Beetles	¾ - 1 tbsp. per 1000 sq. ft.	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.
Radishes (14)	Cutworms	3 - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil.
	Mole Crickets	1½ tbsp. per 1000 sq. ft.	Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	4½ - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
Sweet Corn (7)	Cutworms	3 - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil.
	Seed Corn Maggots	3 - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 2 inches of soil.
	Wireworms	4½ - 6 tbsp. per 1000 sq. ft.	Broadcast before planting and immediately incorporate into the top 4-8 inches of soil.
	Note: To protect bees, do not apply foliarly to corn during the pollen shed period.		
Tomatoes (1)	Cutworms	3 - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil.
	Mole Crickets	1½ tbsp. per 1000 sq. ft.	Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	4½ - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids and Dipterous Leafminers	½ tbsp. per 1000 sq. ft.	Apply as insects appear. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.
	Banded Cucumber Beetles	¾ - 1 tbsp. per 1000 sq. ft.	Apply as insect occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.
Turnips and Turnip Tops (14)	Cutworms	3 - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the soil.
	Mole Crickets	1½ tbsp. per 1000 sq. ft.	Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	4½ - 6 tbsp. per 1000 sq. ft.	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.

Ornamental Insect Control

To control certain insects on Azaleas, Carnations, Chrysanthemums, Gladioli, Roses, Boxwood, Holly, Locust, Maple, Oak, Pine, Spruce and Willow, apply the recommended rate as given below. Try to spray the underside of leaves and penetrate dense foliage. **Precaution:** Do not use on Ferns, Poinsettia, Hibiscus, papaya, Pilea and Gardenia since injury to plants may occur.

Use 1 tablespoon in 3 gallons of water to control the following insects: Aphids, Bagworms, Dipterous Leafminers, Leafhoppers, Mites (Carnation Bud, Carnation Shoot, Clover, Cyclamen, European Red, privet and Two-spotted Spider), Scale Crawlers (Cottoncushion, Lecanium, Pine Needle, San Jose, Soft Scale) and Thrips.

Lawn Pest Control

Excluding Golf Courses and Sod Farms

Precautions: Birds, especially waterfowl, feeding or drinking on treated areas, may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, NY between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. Where irrigation (watering) is recommended in the directions, or if waterfowl, i.e., ducks or geese, can be expected in the treatment area after treatment (except spot treatment), water lawn with at least ¼ inch of water immediately after spraying this product; however, stop watering before puddling occurs.

Use on Bent, Bermuda, Bluegrass, Dichondra, Fescue, Merion and St. Augustinegrass

To control pests listed below, apply recommended rates.

Pests	Rate Per 1000 Square Feet		Remarks
	DC-500	Water	
Lawn Cinch Bugs	4 - 6 tbsp.	3 gals.	Use higher rate for longer residual control and in lawns of dense growth, such as St. Augustine lawns.
Ants, Bermudagrass Mites, Crickets, Earwigs, Lawn Billbugs, Sod Webworms (Lawn Moths)	4 - 6 tbsp.	3 gals.	Spot spray ant hills and wasp ground nest openings. For billbug control, apply when activity is first observed or when chewed or brown grass is present, indicating damage from the insect.
Brown Dog Ticks, Chiggers, Fleas	6 tbsp.	3 gals.	Note: Do not apply to animals. For brown dog ticks, spray grass and under shrubbery, particularly near house.
Millipedes	6 tbsp.	3 gals.	Thoroughly wet grass a few hours before applying.
Rhodesgrass Scales	6 tbsp.	3 gals.	Apply when crawlers first emerge.

Nuisance Pests Outside the Home

Ants, Boxelder Bugs, Brown Dog Ticks, Chiggers, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Flies and Spiders	Mix 3 fl. oz. in 3 gallons of water and thoroughly spray window and door screens, sills, foundations, porch and patio walls and ceilings, entranceways, walks, garbage cans, tree trunks and into cracks and other places where insects can hide. To prevent entry into houses, spray a five-foot band of soil around the house next to the foundation and spray the foundation of the house to a height of 2-3 feet. Repeat application as necessary. Do not treat animals with product. Do not treat plants (other than tree trunks) at this nuisance pest rate.
--	--

Storage and Disposal

Storage: Store in a cool, dry place (preferably locked) inaccessible to children and pets and away from heat and open flame. **Pesticide Disposal:** To avoid harming aquatic organisms in rivers and other surface waters, do not pour the concentrate, spray mixture or rinse water into sanitary drains (for example, toilets, floor drains and sinks) or into storm water sewers (for example, street drains). **Container Disposal:** Securely wrap original container in several layers of newspaper and discard in trash.

PRECAUTIONARY STATEMENTS**Hazards to Humans & Domestic Animals**

CAUTION: May be fatal if swallowed. May be absorbed through the skin. Do not breathe spray mist. Causes substantial but temporary eye damage. Do not get in eyes, on skin or on clothing. Wear goggles or safety glasses. Wash thoroughly after handling and before eating or smoking. Avoid contamination of food and feed. Keep out of reach of domestic animals. Do not use on humans, pets or livestock. Do not contaminate ornamental fish ponds. Do not permit children or pets to go onto sprayed grass until spray has completely dried. Food utensils such as measuring cups should not be used for food purposes after use in measuring pesticides.

First Aid

This product contains diazinon, an organophosphate insecticide that inhibits cholinesterase.

If Swallowed:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious or convulsing person.
If in Eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on Skin or Clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Immediately rinse skin with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If Inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. [You may also call 1-800-xxx-xxxx for emergency medical treatment advice.]

Note to Physician: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. This product contains xylene range aromatic solvent which may present as aspiration hazard. Gastric lavage may be indicated if product was taken internally.

{Note: The first aid statements' grid format will be used if market label space permits; otherwise a paragraph format will be used.}

Environmental Hazards

This product is highly toxic to birds, fish and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas, may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, NY between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. For lawns, where irrigation (watering) is recommended in the directions, or if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), water lawn with at least ¼ inch of water immediately after spraying this product; however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Physical or Chemical Hazards

Do not use, pour, spill or store near heat or open flame.

{ } Denotes alternate/optional language

{ } Denotes language that does not appear on the market labeling