

FLY-CURB 7.76% PREMIX

(Contains RABON Oral Larvicide)

Contains RABON[®] to prevent the development of Horn Flies, Face Flies, House Flies and Stable Flies in the manure of treated cattle, House Flies in the manure of treated swine.

Active Ingredient

Tetrachlorvinphos: 2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate 7.76% (35g/lb.)

Inert Ingredients Roughage product, calcium carbonate and mineral oil.

Keep Out of Reach of Children

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION: Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, get medical attention. Wear long-sleeved shirt and pants; chemical resistant gloves; shoes and socks for protection when handling.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a dry place in original container.

Container Disposal: Completely empty container into application (mixer/feeder) equipment. Then dispose of empty bag in a sanitary landfill or by incineration or if allowed, by State and local authorities by burning. If burned, stay out of smoke. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Cattle containing this product may be fed up to slaughter and to lactating dairy cows without withholding the milk from market during or after treatment.

Start feeding RABON[®] larvicidal feeds early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall until cold weather restricts fly activity.

When fed, this product passes through the digestive system into the animal's manure where it kills fly larvae on contact shortly after fly eggs hatch.

It prevents the development of fly larvae in the manure of treated animals, but is not effective against existing adult flies.

In order to achieve optimum fly control, this product should be used in conjunction with other good management and sanitation practices.

This product will mix uniformly in feeds when standard mixing procedures are followed. Thus, usual problems that are common to all feed preparation and which cause stratification, such as excessive free fall or excessive handling, are to be avoided. It is recommended that appropriate preblending techniques be employed to assure adequate distribution throughout the feed mix. The premix should be preblended with ground grain, protein supplements, mineral mixes, etc. before being added to roughages, such as chopped hay or silage. Mixing time should be adequate to assure uniform dispersion. Optimum performance can be assured only if this product is dispersed uniformly with the feedstuff at the recommended level. Common feed mixing equipment (i.e. vertical mixers, horizontal blenders, mixer/feeder truck) may be used to prepare formulated feeds.

Feeds prepared with this product should not be pelleted, nor be mixed with feeds containing predominantly pellets. Further, this product should not be mixed in liquid feed supplements.

Microfeed

CATTLE

In a Concentrate Feed — Roughage Fed Separately

This product can be used to prepare concentrate feeds that will provide 70 mg. of RABON[®] per 100 pounds of body weight daily.

To prepare a larvicidal concentrate feed, mix this product according to the amount of concentrate to be fed per animal per day. Use the following table as a guide for determining the proper mixing rate.

MIXING DIRECTIONS

Pounds of Concentrate Consumed per Animal per Day	RABON [®] in the Concentrate mg	lb. %	Pounds of RABON [®] 7.7% Oral Larvicide Premix per ton of Concentrate
0.5	1584	0.35	90.0
1.0	792	0.18	45.0
1.5	528	0.12	30.0
2.0	396	0.09	22.5
5.0	159	0.035	9.0
10.0	79	0.018	4.5
15.0	53	0.012	3.0
20.0	39	0.0087	2.3
25.0	32	0.0069	1.8
30.0	26	0.0059	1.5

Feed the appropriate larvicidal concentrate indicated to cattle weighing between 400 and 1200 pounds. For larger cattle weighing between 1200 and 1700 pounds, increase the amount of premix per ton of concentrate to 1½ times that indicated.

CATTLE

In a Complete Ration — No Other Roughage Fed

This product can be used to prepare rations that contain 26.4 mg. of RABON[®] per pound of complete ration.

To prepare a larvicidal ration, mix 1.5 pounds of this product per ton of complete mixed ration containing both grain and roughage.

Full-feed this larvicidal complete ration to feeder cattle weighing from 400 to 1400 pounds or to dairy cattle at a rate to sustain milk production, but not less than 2.6 pounds of the ration per 100 pounds of body weight daily.

SWINE

All swine should be treated.

Pigs (weaners to market weight): Mix 1.3 pounds of this product per ton of meal-type feed and offer free choice. This is equivalent to 22.7 mg. of RABON[®] per pound of feed.

SOWS, BOARS and BREEDING GILTS: Mix 2.6 pounds of this product per ton of meal-type feed and offer 4 to 6 pounds of feed per animal per day. This is equivalent to 45.4 mg. of RABON[®] per pound of feed.

EPA Est. No.: 38092-NB-001

Code No.: 911

EPA Reg. No.: 38092-3

NET WEIGHT: 45 Lbs.