

PM03

38092-3



12/9/99

1081

FLY-CURB 7.76% PREMIX

Contains **RABON® ORAL LARVICIDE** to prevent the development of Horn Flies, Face Flies, House Flies and Stable Flies in the manure of treated cattle, House Flies in the manure of treated swine.

ACTIVE INGREDIENT

Tetrachlorvinphos: (z-2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate).....7.76% (35.2 g/lb)

INERT INGREDIENTS

Roughage product, calcium carbonate and mineral oil.....92.24% (418.4 g/lb)

TOTAL.....100.00% (453.6 g/lb)

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

FIRST AID

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if available by administering syrup of ipecac. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF ON EYES: Flush with plenty of water. Call a physician if irritation persists.

NOTE TO PHYSICIANS AND VETERINARIANS:

Poisoning Symptoms: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps. **Treatment:** Tetrachlorvinphos is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. Atropine is antidotal only if symptoms of cholinesterase inhibition are present. For additional first aid information on RABON Oral Larvicide call 1-800-821-7467 and ask for Boehringer Ingelheim Vetmedica, Inc. technical service department. For additional information on FLY-CURB 7.76% PREMIX call 1-800-228-2022.

Do not use this product on animals simultaneously or within a few days before or after treatment with cholinesterase inhibiting drugs, pesticides or chemicals.

HAZARD TO HUMANS

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction to some individuals. Wear Personal Protective Equipment (PPE) such as long sleeved shirt, long pants, shoes and socks and chemical resistant gloves when handling.

excessive handling, are to be avoided. It is recommended that appropriate preblending techniques be employed to assure adequate distribution throughout the feed mix. The premix should be preblended with ground grain, protein supplements, mineral mixes, etc. before being added to roughages such as chopped hay or silage. Mixing time should be adequate to assure uniform dispersion. Optimum performance can be assured only if this product is dispersed uniformly with the feedstuff at the recommended level. Common feed mixing equipment (i.e., vertical mixers, horizontal blenders, mixer/feeder truck) may be used to prepare formulated feeds.

Feeds prepared with this product should not be pelleted nor be mixed with feeds containing predominantly pellets. Further, this product should not be mixed in liquid feed supplements.

CATTLE

In a Concentrate Feed — Roughage Fed Separately

This product can be used to prepare concentrate feeds that will provide 70 mg. of RABON® per 100 pounds of body weight daily.

To prepare a larvicidal concentrate feed, mix this product according to the amount of concentrate to be fed per animal per day. Use the following table as a guide for determining the proper mixing rate.

MIXING DIRECTIONS

Pounds of Concentrate Consumed per Animal per Day	RABON® in the Concentrate mg/lb.	%	Pounds of Fly-Curb 7.76% Premix per ton of Concentrate
0.5	1584	0.35	90.0
1.0	792	0.18	45.0
1.5	528	0.12	30.0
2.0	396	0.087	22.5
2.5	317	0.063	18.0