

38110-4

4-16-1979

1/2

50 LBS. NET WEIGHT

**MAX-MIN  
FLY CONTROL MINERAL #2  
WITH RABON (R) ORAL LARVICIDE**

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle.

ACTIVE INGREDIENTS:	BY WEIGHT
2-Chloro-1-(2, 4, 6-trichlorophenyl) vinyl dimethylphosphate +	1.00%
INERT INGREDIENTS: ++	99.00%
TOTAL	100.00%

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**  
 SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS**  
**CAUTION**  
 Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

**ENVIRONMENTAL HAZARDS**  
 This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

+ Each pound contains 4.65g RABON (R)  
 ++ Refers only to ingredients which are not larvicidal.  
 EPA REG. NO. 38110-4  
 EPA EST. NO. 38110GA01

**GUARANTEED ANALYSIS:**

Calcium (Ca) Max .....	23.00%
Calcium (Ca) Min .....	20.00%
Phosphorus (P) Min .....	8.00%
Salt (NaCl) Min .....	23.00%
Salt (NaCl) Max .....	25.00%
Iron (Fe) Min .....	1.00%
Copper (Cu) Min .....	0.06%
Manganese (Mn) Min .....	0.20%
Zinc (Zn) Min .....	0.40%
Cobalt (Co) Min .....	0.01%
Iodine (I) Min .....	0.01%
Fluorine (F) Max .....	0.06%
Total Added Mineral Ingredients, Maximum .....	95.00%
Vitamin A USP Units per lb., Minimum .....	200,000
Vitamin D-3 USP Units per lb., Minimum .....	20,000

**FEED INGREDIENTS:**  
 Defluorinated Phosphate, Salt, Ground Limestone, Iron Oxide, Ferrous Carbonate, Cottonseed Meal, Cane Molasses, Vitamin A Palmitate (Stability Improved), D-Activated Animal Sterol, Manganese Oxide, Zinc Oxide, Copper Oxide, Cobalt Carbonate, Anise Oil and Calcium Iodate.

Manufactured by:  
**SOUTHEASTERN MINERALS, INC.**  
 Bainbridge, Georgia 31717 U.S.A.

**DIRECTIONS FOR USE:**  
**GENERAL CLASSIFICATION**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Feed MAX-MIN FLY CONTROL MINERAL #2 WITH RABON (R) ORAL LARVICIDE at the rate of 2.4 ounces per 1000 pounds of body weight per head per day by mixing it into the ground grain or complete daily ration.

Start feeding MAX-MIN FLY CONTROL MINERAL #2 WITH RABON (R) ORAL LARVICIDE early in the spring before flies begin to appear and continue feeding throughout the summer into the fall until cold weather restricts fly activity.

MAX-MIN FLY CONTROL MINERAL #2 WITH RABON (R) ORAL LARVICIDE may be fed up to slaughter and to lactating dairy cows.

MAX-MIN FLY CONTROL MINERAL #2 WITH RABON (R) ORAL LARVICIDE prevents the development of horn flies, face flies, house flies and stable flies on treated cattle but is not effective against house flies on the feeding program during the fly season. Additional measures to reduce the population of house flies may be needed in and around the manure but in other decaying vegetable matter on the premises.

In order to achieve optimum fly control, MAX-MIN FLY CONTROL MINERAL #2 WITH RABON (R) ORAL LARVICIDE should be used in conjunction with other good management and sanitation practices.

**STORAGE AND DISPOSAL**  
 Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

2/2

**ACCEPTED**

**APR 16 1979**

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 38110-4