

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (75-767) WASHINGTON, DC 20460 NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> REGISTRATION <input type="checkbox"/> Reregistration <i>(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)</i>	EPA REGISTRATION 3-110-a	DATE OF ISSUANCE JAN 29 1986
	TERM OF ISSUANCE	
	UNITED STATES DEPARTMENT OF AGRICULTURE	
	NAME OF PESTICIDE PRODUCT KENT-DAN FLY CONTROL Mineral with Baited GEAR Larvicide	

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Southeastern Minerals, Inc.
 P.O. Box 1860
 Birmingham, GA 31717

160 / 152630
18 / 1

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.


Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conventionally registered in accordance with FIFRA, Sec. 3(c)(7)(B) provided that you:

1. Remove the word "GENERAL USE" from the label, and delete the following language: "This product is registered for use on a wide range of crops and is suitable for use by the general public."
2. Change the labeling to read: "This product is registered for use on a wide range of crops and is suitable for use by the general public."
3. Add the words "EPA Reg. No. 3-110-a" to the label.
4. Change "Keep out of lakes, streams, or ponds" or "Do not apply directly to any body of water."
5. Change "Do not use in areas where water contains wild fish."
6. Delete "GENERAL CLASSIFICATION"

Any reference to 'general use' classification must be removed from (the product's name)'s labeling. 'General Use' as a classification is no longer deemed valid by the

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL: 

DATE: 1/29/86



Agency, even when products are not subject to restricted use as in the case of Max-Min Fly Control Mineral #6 With Rabon Oral Larvicide. If a product is classified for restricted use, this classification notifies individuals that the chemical may be applied only by or under the direct supervision of a certified applicator. The absence of a restricted use classification clearly implies that the aforementioned restriction does not apply to a particular pesticide or its uses. A second classification for general use is thereby unnecessary and should not appear on Max-Min Fly Control Mineral #6 With Rabon Oral Larvicide's labeling.

Mixing
in label

- e. Add "Feeds prepared with MAX-MIN FLY CONTROL MINERAL #6 WITH RABON ORAL LARVICIDE should not be pelleted, used with feeds containing predominately pellets, or mixed in liquid feed supplements."
 - f. See the enclosed sample label with mixing and feeding directions and table for determination of rations.
 - g. Please provide the bulk density on the Confidential Statement of Formula (CSF).
 - h. Please provide the chemical identity of the Dri Ani-Mol Ade 2X.
 - i. Please update the label Storage and Disposal statement per PR 83-3.
 - j. The lower Certified Limits for the active ingredient must not fall below the label claim. Please revise the limits.
3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosures

50 LBS. NET WEIGHT
 MAX-MIN
 FLY CONTROL MINERAL #6
 WITH RABON® ORAL LARVICIDE.

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle.

ACTIVE INGREDIENTS:	BY WEIGHT
2-Chloro-1-(2,4,5-trichlorophenyl) vinyl dimethylphosphate +	1.00%
INERT INGREDIENTS:++	99.00%
	TOTAL 100.00%

KEEP OUT OF REACH OF CHILDREN
 CAUTION
 SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

PRECAUTIONARY STATEMENTS
 HAZARDS TO HUMANS
 CAUTION

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

+Each pound contains 4.65g RABON®
 ++Refers only to ingredients which are not larvicidal.

EPA REG. NO. 38110-6
 EPA EST. NO. 38110GA01

ACCEPTED
 with COMMENTS
 EPA Letter Dated:

GUARANTEED ANALYSIS:

Calcium (Ca) Max.....	JAN 29 1986	22.00%
Calcium (Ca) Min.....		19.00%
Phosphorus (P) Min.....	Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.	6.00%
Salt (NaCl) Min.....		15.00%
Salt (NaCl) Max.....	<u>38110-6</u>	18.00%
Iron (Fe) Min.....		1.00%
Copper (Cu) Min.....		0.05%
Manganese (Mn) Min.....		0.20%
Zinc (Zn) Min.....		0.40%
Cobalt (Co) Min.....		0.005%
Iodine (I) Min.....		0.005%
Fluorine (F) Max.....		0.06%
Total Added Mineral Ingredients, Maximum.....		95.00%
Vitamin A SUP Units Per Lb., Minimum.....		200,000
Vitamin D-3 USP Units Per Lb., Minimum.....		20,000

FEED INGREDIENTS:

Defluorinated Phosphate, Salt, Calcium Carbonate, Iron Oxide, Ferrous Carbonate, Rice Mill Byproduct, Cane Molasses, Vitamin A Palmitate (stability improved), D-Activated Animal Sterol, Manganous Oxide, Zinc Oxide, Copper Oxide, Cobalt Carbonate, Anise Oil and Calcium Iodate.

401

**DIRECTIONS FOR USE:
GENERAL CLASSIFICATION**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Feed MAX-MIN FLY CONTROL MINERAL #6 WITH RABON® ORAL LARVICIDE at a rate of 2.4 ounces per 1000 pounds of body weight per head per day by mixing it into the ground grain or complete daily ration.

Start feeding MAX-MIN FLY CONTROL MINERAL #6 WITH RABON® ORAL LARVICIDE early in the spring before flies begin to appear and continue feeding throughout the summer into the fall until cold weather restricts fly activity.

MAX-MIN FLY CONTROL MINERAL #6 WITH RABON® ORAL LARVICIDE MAY BE FED UP TO SLAUGHTER AND TO LACTATING DAIRY COWS.

MAX-MIN FLY CONTROL MINERAL #6 WITH RABON® ORAL LARVICIDE PREVENTS THE DEVELOPMENT OF HORNFLIES, FACE FLIES, HOUSE FLIES AND STABLE FLIES IN THE MANURE OF TREATED CATTLE BUT IS NOT EFFECTIVE AGAINST EXISTING ADULT FLIES. When starting a feed program during the fly season, it is desirable to use other control measures to reduce the population of existing adult flies.

In some cases, supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter of silage on the premises.

In order to achieve optimum fly control, MAX-MIN FLY CONTROL MINERAL #6 WITH RABON® ORAL LARVICIDE should be used in conjunction with other good management and sanitation practices.

STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

50 LBS. NET WEIGHT

MAX-MIN FLY CONTROL MINERAL¹⁶ WITH RABON (R) ORAL LARVICIDE

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle.

ACTIVE INGREDIENTS:	BY WEIGHT
2-Chloro-1-(2, 4, 6-trichlorophenyl) vinyl dimethyl phosphate	1.00%
INERT INGREDIENTS	99.00%
TOTAL	100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

*Each pound contains 4.65g RABON (R)
**Refers only to ingredients which are not toxic.
EPA REG. NO. 38110-6
EPA EST. NO. 38110GA01

GUARANTEED ANALYSIS:

Calcium (Ca) Max	22.0%
Calcium (Ca) Min	19.00%
Phosphorus (P) Min	8.00%
Salt (NaCl) Min	15.00%
Salt (NaCl) Max	18.00%
Iron (Fe)	1.00%
Copper (Cu) Min	0.05%
Manganese (Mn) Min	0.20%
Zinc (Zn) Min	0.40%
Cobalt (Co) Min	0.005%
Iodine (I) Min	0.005%
Fluorine (F) Max	0.06%
Total Added Mineral Ingredients, Maximum	95.00%
Vitamin A SUP Units Per Lb., Minimum	200,000
Vitamin D-3 USP Units Per Lb., Minimum	20,000

FEED INGREDIENTS:

Delloninized Phosphate, Salt, Calcium Carbonate, Iron Oxide, Ferric Carbonate, Zinc Oxide, Calcium Hydroxide, Calcium Hydroxide (stability improved), D-Acivated Animal Sterol, Manganese Oxide, Zinc Oxide, Copper Oxide, Cobalt Carbonate, Anise Oil and Calcium Iodate.

DIRECTIONS FOR USE:

GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Feed MAX-MIN FLY CONTROL MINERAL¹⁶ WITH RABON (R) ORAL LARVICIDE at the rate of 24 ounces per 1000 pounds of body weight per head per day by mixing it into the ground grain or complete daily ration.

Start feeding MAX-MIN FLY CONTROL MINERAL¹⁶ WITH RABON (R) ORAL LARVICIDE early in the spring before flies begin to appear and continue feeding throughout the summer into the fall until cold weather restricts fly activity.

MAX-MIN FLY CONTROL MINERAL¹⁶ WITH RABON (R) ORAL LARVICIDE may be fed up to slaughter and to lactating dairy cows.

MAX-MIN FLY CONTROL MINERAL¹⁶ WITH RABON (R) ORAL LARVICIDE prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle but is not effective against existing adult flies. When starting a feeding program during the fly season, it is desirable to use other control measures to reduce the population of existing adult flies.

In some cases, supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or sludge on the premises.

In order to achieve optimum fly control, MAX-MIN FLY CONTROL MINERAL¹⁶ WITH RABON (R) ORAL LARVICIDE should be used in conjunction with other good management and sanitation practices.

STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

BEST AVAILABLE COPY