



QUALI-TECH

Rabon Fly Killer Premix S-3

(Contains Rabon® Oral Larvicide)

To prevent the development of House Flies in the manure from treated swine.

ACTIVE INGREDIENT	BY WEIGHT	* Refers only to ingredients which are not larvicidal.
2-Chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate.....	100%**	** RABON insecticide-Contains 4.54 grams of RABON per pound.
INERT INGREDIENTS	99.00%*	RABON is the registered trademark of Shell Oil Company.
TOTAL	100.00%	U.S. Patent No. 3,102,542.

KEEP OUT OF REACH OF CHILDREN
CAUTION

See below for additional precautionary statements

ACCEPTED
AUG 5 1983
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 38119-7

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
CAUTION

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE
GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

No pre-slaughter withholding period is required.

Pigs (weaners to market weight): Mix 10 lbs. of Swine Oral Larvicide per ton of meal type feed and offer free choice. This is equivalent to 22.7 mg. of Rabon per lb. of feed.

Sows, Boars, and Breeding Gilts: Mix 20 lbs. of Swine Oral Larvicide per ton of meal type feed and offer 4-8 lbs. of feed per animal per day. This is equivalent to 45.4 mg. of Rabon per lb. of feed.

Start feeding Swine Oral Larvicide early in the spring before flies begin to appear and continue feeding program throughout the summer and into the fall until cold weather restricts fly activity.

All swine should be treated. Swine Oral Larvicide prevents development of house flies in the manure from treated swine but is not effective against existing adult flies.

In some cases, supplemental fly control measures may be needed in and around the area to control adult flies which move in from other areas or which breed on decaying vegetable matter elsewhere on the premises.

In order to achieve optimum fly control, Swine Oral Larvicide should be used on conjunction with other good management and sanitation practices.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal: Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, State or Local procedures.

Container Disposal: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State or Local authorities.

WARRANTY AND LIMITATION OF DAMAGES

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions.

Seller makes no other express or implied warranty of fitness or of merchantability, and no agent of Seller is authorized to do so except in writing and with specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which claim is made.

Manufactured By:
QUALI-TECH PRODUCTS, INC.
318 LAKE HAZELTINE DRIVE
CHASKA, MINNESOTA 55318

EPA Reg. No. 38119-7
EPA Est. No. 38119-MN-1

NET WEIGHT: 10 POUNDS

PC
Lab. Rev. 011783
LOT NO.