

BLADEX MSMA HERBICIDE

ACCEPTED
SEP 17 1982
Under the Federal Insecticide, Fungicide, and Rodenticide Act
of 1947 and under the
pesticide
EPA Reg. No. _____

ACTIVE INGREDIENTS	% BY WEIGHT
2-4-chloro-6-(ethylamino)-s-triazin-2-yl amino-2-methylpropionitrile	17.21%
Monosodium Acid Methanearsonate	34.35%
INERT INGREDIENTS	48.44%
TOTAL	100.00%

Total arsenic, all in water soluble form, expressed as elemental 15.9%
One gallon of Bladex-MSMA Herbicide contains two (2) pounds of Bladex and four (4) pounds of MSMA.

KEEP OUT OF REACH OF CHILDREN
WARNING
See Additional Precautions on Side panel

EPA Reg. No. _____ Net Contents _____
EPA Est. No. _____

Manufactured By Setre Chemical Company
Memphis, Tn. 38137

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

May be fatal if swallowed, inhaled, or absorbed through the skin. Do not breathe vapors, or spray mist. Do not get in eyes, on skin or on clothing.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Give large quantities of water and induce vomiting by touching back of throat with fingers. Never give anything by mouth to an unconscious person.

If in Eyes: Flush with water for at least 15 minutes, get medical attention.

If on Skin: Wash thoroughly with soap and water, get medical attention.

ENVIRONMENTAL HAZARDS

Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

BEST DOCUMENT AVAILABLE

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Storage: Do not store near feed, food or water.

Pesticide Disposal. Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state or local procedures.

Container Disposal: Triple rinse/or equivalent and offer for recycling, or reconditioning, or dispose of in a sanitary landfill, or by other approved, State and local procedures.

APPLICATION DIRECTIONS

Setre Bladex-MSMA herbicide is effective as a directed postemergence treatment for the control of certain broadleaf weeds. Bladex-MSMA herbicide may be used in Alabama, Arkansas, Georgia, Louisiana, Mississippi, Missouri, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Gulf Coast, Coastal Bend and Blacklands only.

Bladex-MSMA herbicide may be applied postemergence following a preemergence application of Bladex. For coarse soils, apply no more than one postemergence and one preemergence application to the same crop in any one year. For medium and fine soils, apply no more than two postemergence and one preemergence application to the same crop in any one year.

MIXING AND SPRAYING

Application equipment should be thoroughly cleaned if another pesticide has been previously used.

Sufficient agitation should be provided during the filling operation and application to insure uniform mixing. Fill the spray tank with one-half (1/2) full of clean water and while pump and agitator are running, add the recommended amount of Bladex-MSMA herbicide. Fill with water to desired level.

Allow the spray mixture to agitate for several minutes prior to application. If the mixture is allowed to stand in a spray tank or nurse tank for an extended period of time (several hours to overnight), agitate for several minutes prior to application.

Apply with flat fan nozzles (even distribution flat fan for band application) with 50 mesh nozzle and in-line screens. Apply at 20-40 psi and avoid drift. Apply in properly calibrated spray equipment.

BEST DOCUMENT AVAILABLE

COTTON: Directed Postemergence: Apply Setre Bladex-MSMA herbicide after cotton is 6 inches tall but before first bloom, and before weeds are 2 inches tall. The spray should be directed to the soil around the base of the cotton plants, taking care to prevent spraying of cotton foliage, as injury will occur. Maximum coverage of uses is important for good control. Apply 1.2-2.0 quarts broadcast, 0.4-0.6 quarts on a 12 inch band and 0.6-1.0 quarts on a 19 inch band (38 inch rows) in 15-30 gallons of spray solution per sprayed acre. Addition of an approved (40 CFR .180 .1001) nonionic spreader surfactant at a rate of 1-2 quarts per 100 gallons of water (as directed by manufacture) is recommended.

Weeds Controlled: Cocklebur, prickly sida (leaved), annual morninglory (tall, smallflower, small white, ivyleaf), pigweed (redroot, sping), Johnsongrass, nutgrass, watergrass, goosegrass, dallisgrass and spotted spurge.

Do not graze or feed foliage from treated areas to livestock.

NOTICE OF WARRANTY

Setre Chemical Company warrant that this material conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the Directions for Use and Conditions of Sale, subject to the inherent risks referred to therein.

Setre Chemical Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of Chemical Company is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and not include consequential commercial damages such as loss of profit or value, etc.

BEST DOCUMENT AVAILABLE



F