

STOCKADE NO FLY RABON(r) 7.76 ORAL LARVICIDE PREMIX

To prevent the development of Horn Flies, Face Flies, House Flies and Stable Flies in the manure of treated cattle.

ACTIVE INGREDIENT	BY WEIGHT
2-chloro-1-(2,4,5-trichlorophenyl)vinyl dimethyl phosphate	7.76%
INERT INGREDIENTS	92.24%
	Total 100.00%

SEE REVERSE SIDE OF LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS


PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
CAUTION
 Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS
 This product is harmful to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.



When fed to cattle STOCKADE NO FLY RABON(r) ORAL LARVICIDE passes through the digestive system into the animal's manure where it kills fly larvae on contact shortly after fly eggs hatch. By preventing larval development, STOCKADE NO FLY RABON(r) ORAL LARVICIDE helps to control horn flies and face flies on pasture and house flies and stable flies in and around cattle lots.

STOCKADE NO FLY RABON(r) 7.76 ORAL LARVICIDE PREMIX will mix uniformly in cattle feeds when standard mixing procedures are followed. Thus, usual problems that are common to all feed preparation and which cause stratification, such as excessive free-fall or excessive handling, are to be avoided. It is recommended that appropriate techniques be employed to assure adequate distribution of STOCKADE NO FLY RABON(r) 7.76 ORAL LARVICIDE PREMIX throughout the feed mix. The premix should be preblended with ground grain, protein supplements, mineral mixes, etc., before being added to roughages, such as chopped hay or silage. Mixing time should be adequate to assure uniform dispersion. Optimum performance can be assured only if STOCKADE NO FLY RABON(r) 7.76 ORAL LARVICIDE PREMIX is dispersed uniformly with the feedstuff at the recommended level. Common feed mixing equipment (i.e., vertical mixers, horizontal blenders, mixer/folder truck) may be used to prepare formulated feeds.

Feeds prepared with STOCKADE NO FLY RABON(r) 7.76 ORAL LARVICIDE PREMIX should not be pelleted, nor be mixed with feeds containing predominantly pellets. Further, STOCKADE NO FLY RABON(r) 7.76 ORAL LARVICIDE PREMIX should not be mixed in liquid feed supplements.

SEE REVERSE SIDE OF LABEL FOR FURTHER FEEDING DIRECTIONS

Manufactured By
 VERN LIVESTOCK SUPPLEMENT CO., INC.
 A Harvest Industries Company
 P.O. Box 2056 (905) 327-3216 E.P. Reg. No. 3227-1
 130 Industrial Street
 Palmdale, California 93307

FEEDING DIRECTIONS

IN A COMPLETE RATION -- NO OTHER POUCHAGE FEED

STOCKADE NO FLY BARON(r) 7.76 ORAL LARVICIDE PREMIX can be used to prepare rations that contain 26.4 mg of Babon per pound of complete ration.

To prepare a larvicidal ration, mix 1.5 pounds of STOCKADE NO FLY BARON(r) 7.76 ORAL LARVICIDE PREMIX per ton of complete mixed ration containing both grain and roughage.

Full-feed this larvicidal complete ration to feeder cattle weighing from 400 to 1400 pounds, or to dairy cattle at a rate to sustain milk production, but not less than 2.6 pounds of the ration per 100 pounds of live weight daily.

IN A CONCENTRATED FEED -- POUCHAGE FEED SEPARATELY

STOCKADE NO FLY BARON(r) 7.76 ORAL LARVICIDE PREMIX can be used to prepare concentrate feeds that will provide 792 mg of Babon per animal per day.

To prepare a larvicidal concentrate feed, mix STOCKADE NO FLY BARON(r) 7.76 ORAL LARVICIDE PREMIX according to the amount of concentrate to be fed per animal per day. Use the following table as a guide for determining the proper mixing rate.

MIXING DIRECTIONS

Pounds of Concentrate Consumed per animal per day	Babon in the concentrate mg/lb	%	Pounds of STOCKADE NO FLY BARON(r) 7.76 ORAL LARVICIDE PREMIX per ton of concentrate
.5	1584	0.35	90.0
1.0	792	0.17	45.0
1.5	528	0.12	30.0
2.0	396	0.097	22.5
5.0	158	0.035	9.0
10.0	79	0.018	4.5
15.0	53	0.012	3.0
20.0	39	0.0097	2.3
25.0	32	0.0080	1.9
30.0	26	0.0058	1.5

Feed the appropriate larvicidal concentrate to cattle weighing between 400 and 1200 pounds. For larger cattle weighing between 1200 and 1700 pounds, increase the amount of concentrate fed to 1½ times that recommended.

Start feeding Babon larvicidal feeds early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall until cold weather restricts fly activity.

Rations containing STOCKADE NO FLY BARON(r) ORAL LARVICIDE may be fed up to slaughter and to lactating dairy cows without withholding the milk from the market during or after treatment.

STOCKADE NO FLY BARON(r) ORAL LARVICIDE prevents the development of Horn flies, face flies, house flies and stable flies in the manure of treated cattle but is not effective against existing adult flies.

In some cases, supplemental fly control measures may be needed in and around cattle lots and barns to control adult flies and stable flies which can breed not only in cattle manure but other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, STOCKADE NO FLY BARON(r) 7.76 ORAL LARVICIDE PREMIX should be used in conjunction with other good management and sanitation practices.

STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal. Storage is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.