


NET CONTENTS



CORROSIVE

SODIUM HYPOCHLORITE SOLUTION

10% AVAILABLE CHLORINE

ACTIVE INGREDIENT:
SODIUM HYPOCHLORITE . . . 10.0%
INERT INGREDIENTS . . . 90.0%
and also used in size 100.0%

DANGER
KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED
STRONG OXIDANT, MAY PRODUCE CHEMICAL BURNS
Do not get on skin or in eyes. Do not use ammonia or other bowl cleaners with this product. To do so will release hazardous gases.
FIRST AID: In case of skin and eye contact, flush with plenty of water. If skin irritation persists, get medical attention. Internal—give milk, water, or egg whites. Call physician immediately.

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds, or public water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

WATER PURIFICATION
In your treatment of well water, there is a possibility of iron, manganese, and hydrogen sulfide. These may be used to remove iron and manganese from the water to make it potable. The application of one gallon of sodium hypochlorite (10% available chlorine) to 10,000 gallons of water or its equivalent will oxidize a maximum of 1.5 ppm of iron and 0.5 ppm of manganese. Apply the sodium hypochlorite solution through a common faucet and disperse the available chlorine approximately 20 minutes after treatment with a soluble chlorine test kit. Adjust chlorine level as necessary to give a chlorine residual of at least 0.2 ppm which indicates an effective dosage.

MILK PLANT AND DAIRY EQUIPMENT
All equipment involved in the dairy industry may be sanitized after thoroughly prewashing by rinsing with a solution of sodium hypochlorite containing 500 ppm of available chlorine. This solution may be obtained by adding 1/2 gallon of 10 percent sodium hypochlorite to 10 gallons of potable water. It is recommended that this diluted solution be used periodically with a chlorine test kit to check for remaining high chlorine residuals (50-600 ppm), and that the residual chlorine be diluted to about 10 ppm. Do not use a 10 ppm level of chlorine in milk cans or in milk bottles and do not allow it to remain in contact with rubber parts. Prepared by adding 1 cup of 10 percent sodium hypochlorite to 100 gallons of potable water. After sanitizing, the equipment should be rinsed thoroughly with potable water.

MEAT AND POULTRY PROCESSING PLANTS
Surfaces and equipment in the meat industry may be sanitized by prewashing thoroughly with water containing 500 ppm (0.5 per cent) available chlorine (10 gallons of 10 percent sodium hypochlorite solution to 100 gallons of potable water) may be used for sanitizing walls, floors, ceilings and other areas. Solutions containing 200 ppm (0.2 per cent) of 10 percent sodium hypochlorite solution to 100 gallons of potable water may be used on other product equipment and machinery. (NOTE: After sanitizing, do not allow contact with the product. If a contact time higher than 200 ppm available chlorine is used on other product equipment and machinery, apply product over the entire area.)

LAUNDRY BLEACH
To use 1 part of 10 percent sodium hypochlorite solution with 1 part of water to obtain an equivalent to the normal household bleach solution of 1.25 percent. For instance, combine 1 part of 10 percent sodium hypochlorite solution to 9 parts of water to make 10 parts of 1.25 percent solution. Use 1 ounce of the diluted solution to use 1 1/2 pounds of laundry.

SWIMMING POOL TREATMENT
For every 10,000 gallons of chlorine residual, add 1 ounce of 10% Sodium Hypochlorite Solution per 1000 gallons of pool water. Approximate dosage of 1.0 to 1.5 ppm is obtained as determined by the use of a chlorine test kit. Therefore, adjust the dosage according to your pool as indicated on the chlorine test kit or as needed to maintain 1.0 to 1.5 ppm chlorine residual at all times. For pool water supersaturated with chlorine test kit. Keep the pool at pH 7.2 to 7.4 as determined by a pH test kit.

GALLONS OF WATER IN POOL	SODIUM HYPOCHLORITE SOLUTION REQUIRED
500	5 ounces
1,000	10 ounces
2,000	20 cups
10,000	100 cups
20,000	200 cups
50,000	500 cups

(1 cup equals 16 ounces equals 8 ounces liquid)

To control the growth of algae, prevent bacteria or fungi each week by adding sodium hypochlorite solution at 4 to 5 times the normal rate. Apply after children have left the pool and (NO SWIM) after bathing with the chlorine residual has dropped to 1.5 ppm or less. Dilute the required amount of sodium hypochlorite to at least 10 volumes of water added to the solution. This diluted solution can then be added to the discharge of the circulation system while the circulation pump is in operation. In dry or hot weather, application of the diluted solution can be sprayed over the surface of the pool from the edge with a stream sprayer gun.

NOTE
Sodium Hypochlorite degrades with age, high temperatures, and sunlight exposure. Use a test kit and correct the amount of recommended dosage to obtain the required level of available chlorine.

STORAGE AND SHIPMENT
Store this product in a cool, dry area away from fire, heat, and acids. Do not store in plastic containers. Do not use a plastic container for storage. Do not use a plastic container for storage. Do not use a plastic container for storage. Do not use a plastic container for storage.

NOTICE
Safety glasses and protective clothing are recommended when using this product. Safety glasses should be worn at all times. Do not use in areas of fire and explosion. Do not use in areas of fire and explosion.

EPA Reg. No. 38542-7
EPA Est. No. 38542-AR-01
ORM-B NA-1791

ACCEPTED with COMMENTS
EPA Letter Date:

JUN 25 1984

38542-7



Manufactured By
Swift Chemical Company, Inc.
MANUFACTURERS OF QUALITY CHEMICAL PRODUCTS

201 South Arkansas • Rogers, Arkansas 72756
Phone (501) 636-0420

U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDE PROGRAM REGISTRATION DIVISION (HHS-707) WASHINGTON, D.C. 20460	EPA REGISTRATION NO.	DATE OF ISSUANCE
	TERM OF ISSUANCE	
NOTICE OF PESTICIDE: <small>REGISTRATION</small> <small>REREGISTRATION</small> <i>(Under the Federal Insecticide, Fungicide, and Rodenticide Act)</i>	NAME OF PESTICIDE PRODUCT	
	Pesticide Product Name	
NAME AND ADDRESS OF REGISTRANT (Include ZIP codes)		
[Faint, illegible text]		
BEST DOCUMENT AVAILABLE		
NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.		
On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.		
A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.		
Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.		
<input type="checkbox"/> ATTACHMENT IS APPLICABLE		
SIGNATURE OF APPROVING OFFICIAL		DATE