


ACCEPTED
 MAY 22 1973
 Unit _____, _____, _____, and
 Rochester, N.Y., is submitted for the
 possible registration under
 EPA Reg. No. 38542-7

STATEMENT OF PRACTICAL TREATMENT (FIRST AID)
 IF CONTACT WITH EYES OCCURS, flush with water for at least 15 minutes. Get prompt medical attention.
 IF CONTACT WITH SKIN OCCURS, wash with plenty of soap and water.
 IF SWALLOWED, drink large quantities of milk or gelatin solution. If these are not available, drink large quantities of water. DO NOT give vinegar or other acids. DO NOT induce vomiting. Get prompt medical attention.



EPA Reg No 385427
 EPA Est No 38542 AR01

DOT: NA1791 CORROSIVE

SODIUM HYPOCHLORITE SOLUTION

KEEP OUT OF REACH OF CHILDREN


DANGER

ACTIVE INGREDIENT
 SODIUM HYPOCHLORITE 10.0%
 INERT INGREDIENTS 90.0%

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER: Corrosive may cause severe skin and eye irritation or chemical burns to broken skin. Causes eye damage. Wear safety glasses or goggles and rubber gloves when handling this product. Wash after handling. Avoid breathing vapors. Vacate poorly ventilated areas as soon as possible. Do not return until strong odors have dissipated.
ENVIRONMENTAL HAZARDS
 This product is toxic to fish. Do not discharge into lakes, streams, ponds or public waterways unless in accordance with a NPDES permit. For guidance, contact the regional office of the U.S. Environmental Protection Agency.
PHYSICAL OR CHEMICAL HAZARDS
STRONG OXIDIZING AGENT: Mix only with water according to label directions. Mixing this product with chemicals (e.g. ammonia, acids, detergents, etc) or organic matter (e.g. urine, feces, etc) will release chlorine gas which is irritating to eyes, lungs and mucous membranes.

It is a violation of Federal law to use this product in a manner inconsistent with the labeling.

See other precautions on back panel.
 FOR SALE TO, USE AND STORAGE BY SERVICE PERSONS ONLY

NET CONTENTS  Manufactured By
Swift Chemical Company, Inc
 201 South American Boulevard, Atlanta, GA 30328
 Phone: (404) 521-0442

BEST AVAILABLE COPY

27

DISINFECTION OF SWIMMING Pools
PUBLIC SYSTEMS: Use a ratio of 1.3 oz. of this product to 100 gallons of water. Begin treating the pool with a hypochlorite until a free available chlorine residual of at least 0.3 ppm and no more than 0.8 ppm is obtained throughout the distribution system. Check water frequently with a chlorine test kit. Bacteriological sampling must be conducted at a frequency no less than that prescribed by the National Interstate Drinking Water Regulations. Contact your local health department for further details.

WALK PLAY AND BIKE EQUIPMENT
IMMERSIVE FROM CONTACT SURFACES:
CLEAN IN PLACE METHOD: Thoroughly clean equipment after use. Prepare a solution of a 200 ppm available chlorine sanitizing solution equal to 100% of volume capacity of the equipment by using this product in a ratio of 2 oz. of product with 10 gallons of water. Pump solution through the system until full flow is obtained at all connections. The system is completely filled with the sanitizer and will air be removed from the system. Close drain valves and hold under pressure for at least 10 minutes to insure contact with all internal surfaces. Remove pump cleaning solution from drain valve and test with a chlorine test kit. Repeat entire cleaning/sanitizing process if effluent contains less than 50 ppm available chlorine. Rinse drains with potable water prior to use.

WALK AND PLAY PROCESSING PLANTS
IMMERSIVE FROM CONTACT SURFACES, WALLS, FLOORS, CEILING, ETC.
EFFECTIVE METHOD: Prepare a sanitizing solution by thoroughly mixing 9 oz. of this product with 10 gallons of water to provide approximately 600 ppm available chlorine by weight. Clean surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution. Maintain contact with the sanitizer for at least 2 minutes. Do not rinse equipment with water after treatment and do not wash equipment overnight.

IMMERSIVE FROM CONTACT SURFACES EQUIPMENT AND FACILITIES:
EFFECTIVE METHOD: A solution of 100 ppm available chlorine may be used in the sanitizing solution if a chlorine test kit is available. Solutions containing an initial concentration of 100 ppm available chlorine must be tested and adjusted periodically to insure that the available chlorine does not drop below 50 ppm. Prepare a 100 ppm sanitizing solution by thoroughly mixing 1.3 oz. of this product with 10 gallons of water. If no test kit is available, prepare a sanitizing solution by thoroughly mixing 3 oz. of this product with 10 gallons of water to provide approximately 200 ppm available chlorine by weight.

Clean equipment prior to use in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution. Sanitizing contact time shall be maintained at a minimum of 2 minutes. If chlorine residue from the sanitizer is objectionable, it should be removed by a suitable test kit. Rinse drains with potable water after treatment and do not wash equipment overnight.

Sanitizing solution should not be used for general cleaning but may be used for sanitizing purposes.

Swimming Pool Water Disinfection
For a one pump or spring start up, superchlorinate with 54 to 106 oz. of product for each 10,000 gallons of water to provide 5 to 10 ppm available chlorine by weight. Check the level of available chlorine with a test kit. Adjust and maintain pool water pH to between 7.2 to 7.6. Adjust and maintain the alkalinity of the pool to between 50 to 100 ppm.

To maintain the pool and sanitize or by a feeder device 1.3 oz. of this product for each 10,000 gallons of water to provide one available chlorine residual between 0.5 and 1.0 ppm by weight. Sanitized pools should maintain a residual of 1.0 to 1.3 ppm available chlorine. Test the pH, available chlorine residual and alkalinity of the water frequently with appropriate test kits. Frequency of water treatment will depend upon temperature and number of swimmers.

Every 3 days, or as necessary, superchlorinate the pool with 54 to 106 oz. of product for each 10,000 gallons of water to provide 5 to 10 ppm available chlorine by weight. Check the level of available chlorine with a test kit. Do not reenter pool until the chlorine level is between 1.0 to 1.3 ppm.

At the end of the swimming season or when water is to be drained from the pool, chlorine that will be used to dissipate iron treated pool water before discharge. Do not chlorinate the pool water within 24 hours prior to discharge.

WATERING POOLS: Water in a still clear and clean, apply 4 oz. of product per 1000 gallons, until filter is running. To obtain a 3 ppm available chlorine residual, as determined by a suitable test kit. Cover pool, progress water, filter and heater components in winter by following manufacturers' instructions.

POOL EGG SANITIZATION
Thoroughly clean all eggs thoroughly with 3 oz. of this product with 10 gallons of water to produce a 200 ppm available chlorine solution. The sanitizer temperature should not exceed 120 deg F. Spray the water sanitizer so that the eggs are thoroughly wetted. Rinse the eggs to thoroughly dry before coating or breeding. Do not apply a potable water rinse. The solution should not be re-used to sanitize eggs.

NOTE:
Sodium hypochlorite degrades with age, high temperature and sunlight exposure. Use a chlorine test kit and increase the amount of recommended dosage to obtain the required level of available chlorine.

STORAGE AND DISPOSAL

Store this product in a cool, dry area away from direct sunlight and heat to avoid evaporation. In case of spill, flood, or other large quantities of water, product or container that cannot be used, it should be diluted with water and disposed in a suitable area. Do not reuse empty containers but triple rinse the container after use for recycling or reconditioning or puncture and disposal of the container, landfill, or incineration, or if allowed by state and local authorities, by burning. If other rules apply, use them.

NOTICE:
Do not exceed the maximum exposure or tolerance concerning the use of this product. It is not intended for use as a disinfectant. Do not use for purposes other than those listed on the label. Do not use for purposes other than those listed on the label. Do not use for purposes other than those listed on the label.

BEST AVAILABLE COPY

Handwritten mark or signature.