

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

DANGER: Corrosive, may cause severe skin irritation or chemical burns to broken skin. Causes eye damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling this product. Wash after handling. Avoid breathing vapors. Vacate poorly ventilated areas as soon as possible. Do not return until odors have dissipated.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams or ponds. Treated effluent may not be discharged into lakes, streams, ponds or public waters without a valid discharge permit. For guidance, contact the regional office of the Environmental Protection Agency.

PHYSICAL AND CHEMICAL HAZARDS: STRONG OXIDIZING AGENT. Mix only with water according to label directions. Mixing this product with gross filth such as feces, urine, etc., or with ammonia, acids, detergents or other chemicals may release hazardous gases irritating to eyes, lungs and mucous membranes.

manufactured by

**RIVIERA POOL SERVICE, INC.
4215 N. Federal Highway
Ft. Lauderdale, FL 33308**

**EPA REG. NO. 38547-1 EPA EST. NO.
NET CONTENTS**

JUL 3 1978

38547-1

AQUA PURE

BEST DOCUMENT AVAILABLE

**"FOR SWIMMING POOL CHLORINATION
AND SANITIZING."**

ACTIVE INGREDIENT
SODIUM HYPOCHLORITE 9.2%
INERT INGREDIENTS 90.8%

**KEEP OUT OF REACH OF CHILDREN
DANGER**

FIRST AID: If on skin, wash with plenty of soap and water. If in eyes, flush with water for at least 15 minutes. Get medical attention. If swallowed, drink large quantities of milk, or gelatin solution or, if these are not available, drink large quantities of water. Do NOT give vinegar or other acids. Do NOT induce vomiting. Get prompt medical attention.

See additional precautions on side panel

**DIRECTIONS FOR USE
GENERAL CLASSIFICATION**

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT
IN A MANNER INCONSISTENT WITH ITS LABELING**

DIRECTIONS FOR SWIMMING POOL CHLORINATION

For a new pool or for spring start-up: superchlorinate with one (1) pint (16 fl. ozs.) of this sodium hypochlorite solution for each 2500 gallons of water. This dosage is equivalent to 5 ppm available chlorine by weight.

For Pool Maintenance:

- A) Adjust pool water pH to 7.6 - 7.2 range and maintain.
B) Add manually or by a feeder device this sodium hypochlorite solution at a rate to maintain an available chlorine residual of 0.6 to 1.0 ppm. One-half pint (8 fl. ozs.) of this sodium hypochlorite solution for each 6000 gallons of water will provide 1.0 ppm available chlorine by weight. Frequency of additions to maintain 0.6 to 1.0 ppm available chlorine will depend upon temperature and number of swimmers. Use test kit to make certain the pH and chlorine residual are in the proper range.

**DIRECTIONS FOR DISINFECTION OF POTABLE WATER
FOR HOME WELL WATER SYSTEMS**

Dilute this sodium hypochlorite solution in the ratio of one part sodium hypochlorite solution to 9 parts softened water. Mix sodium hypochlorite solution and water thoroughly and begin feeding of solution with a hypochlorinator (metering pump). Maintain a free available chlorine residual of at least 0.2 ppm and no more than 0.6 ppm throughout the distribution system, as determined by a DPD chlorine test kit. Bacteriological sampling must be conducted at a frequency no less than that prescribed by the National Interim Primary Drinking Water Regulations. Check water frequently with a DPD chlorine test kit.

*Contact your local Health Department for further details.

DIRECTIONS FOR SANITIZING FOOD PROCESSING OR DAIRY EQUIPMENT

Clean equipment in the normal manner. Just before using, rinse all surfaces thoroughly with this sodium hypochlorite solution containing 200 ppm available chlorine. Maintain contact with disinfectant for a minimum of two minutes. Do not rinse with water after treatment with sodium hypochlorite solution. Do not soak overnight. Addition of one fluid ounce (2 tblsp.) of this sodium hypochlorite solution per 4 gallons of water will provide approximately 200 ppm available chlorine by weight.

NOTE: This product degrades with age. Use a chlorine test kit and increase dosage as necessary to obtain the required level of available chlorine.

STORAGE AND DISPOSAL: Store in a cool, dry area away from direct sunlight. In case of spill, flood area with large quantities of water. Rinse empty containers thoroughly with water and either return to manufacturer or discard by placing in trash collection or burying in an approved landfill. Product or rinseate that cannot be reused, should be diluted with water and disposed of in a sanitary sewer. Do not contaminate food or feed by storage, disposal or cleaning of equipment.

**CAUTIONARY STATEMENTS
HUMANS & DOMESTIC ANIMALS**

may cause severe skin irritation or chem-
ical damage. Causes eye damage. Do not get in eyes.
Wear goggles or face shield and rubber
gloves when handling this product. Wash after handling. Avoid
contact with poorly ventilated areas as soon as possible
until odors have dissipated.

HAZARDS: This pesticide is toxic to fish,
birds and ponds. Treated effluent may not be
used in streams, ponds or public waters without a
permit. For guidance, contact the regional office
of the Environmental Protection Agency.

CHEMICAL HAZARDS: STRONG OXIDIZING
agent. Do not mix with water according to label directions. Mix-
ing with gross filth such as feces, urine, etc., or with
other agents or other chemicals may release haz-
ardous fumes to eyes, lungs and mucous membranes.

Manufactured by

POOL SERVICE, INC.
1001 Federal Highway
Apopka, FL 33308

547-1 EPA EST. NO.

CONTENTS

JUL 3 1978
3854-7-1
AQUA PURE
BEST DOCUMENT AVAILABLE
"FOR SWIMMING POOL CHLORINATION
AND SANITIZING."
ACTIVE INGREDIENT
SODIUM HYPOCHLORITE 9.2%
INERT INGREDIENTS 90.8%
**KEEP OUT OF REACH OF CHILDREN
DANGER**
FIRST AID: If on skin, wash with plenty of soap and water. If in
eyes, flush with water for at least 15 minutes. Get medical atten-
tion. If swallowed, drink large quantities of milk, or gelatin solu-
tion or, if these are not available, drink large quantities of water.
Do NOT give vinegar or other acids. Do NOT induce vomiting.
Get prompt medical attention.
See additional precautions on side panel.

**DIRECTIONS FOR USE
GENERAL CLASSIFICATION**
IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT
IN A MANNER INCONSISTENT WITH ITS LABELING

DIRECTIONS FOR SWIMMING POOL CHLORINATION
For a new pool or for spring start-up, superchlorinate with one (1) pint (16 flu. ozs.) of
this sodium hypochlorite solution for each 2500 gallons of water. This dosage is equiv-
alent to 5 ppm available chlorine by weight.
For Pool Maintenance:
A) Adjust pool water pH to 7.0 - 7.2 range and maintain.
B) Add manually or by a feeder device this sodium hypochlorite solution at a rate to
maintain an available chlorine residual of 0.6 to 1.0 ppm. One half pint (8 flu. ozs.)
of this sodium hypochlorite solution for each 6000 gallons of water will provide
1.0 ppm available chlorine by weight. Frequency of additions to maintain 0.6 to 1.0
ppm available chlorine will depend upon temperature and number of swimmers.
Use test kit to make certain the pH and chlorine residual are in the proper range.

**DIRECTIONS FOR DISINFECTION OF POTABLE WATER
FOR HOME WELL WATER SYSTEMS**
Dilute this sodium hypochlorite solution in the ratio of one part sodium hypochlorite
solution to 9 parts softened water. Mix sodium hypochlorite solution and water thor-
oughly and begin feeding of solution with a hypochlorinator (metering pump). Main-
tain a free available chlorine residual of at least 0.2 ppm and no more than 0.6 ppm
throughout the distribution system as determined by a DPD chlorine test kit. Bacteri-
ological sampling must be conducted at a frequency no less than that prescribed by
the National Interim Primary Drinking Water Regulations. Check water frequently
with a DPD chlorine test kit.
*Contact your local Health Department for further details.

DIRECTIONS FOR SANITIZING FOOD PROCESSING OR DAIRY EQUIPMENT
Clean equipment in the normal manner. Just before using, rinse all surfaces thor-
oughly with this sodium hypochlorite solution containing 200 ppm available chlorine. Main-
tain contact with disinfectant for a minimum of two minutes. Do not rinse with water
after treatment with sodium hypochlorite solution. Do not soak overnight. Addition of
one fluid oz. (2 bisps.) of this sodium hypochlorite solution per 4 gallons of water will
provide approximately 200 ppm available chlorine by weight.
NOTE: This product degrades with age. Use a chlorine test kit and increase dosage
as necessary to obtain the required level of available chlorine.

STORAGE AND DISPOSAL: Store in a cool, dry area away
from direct sunlight. In case of spill, flood area with large quantities of water. Rinse
empty container thoroughly with water and either return to manufacturer or discard
by placing in trash collection or burying in an approved landfill. Product or rinse that
cannot be used should be diluted with water and disposed of in a sanitary sewer. Do
not contaminate food or feed by storage, disposal or cleaning of equipment.